

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TORSION NUMBERS OF AUGMENTED GROUPS WITH APPLICATIONS TO KNOTS AND LINKS
Autor: Silver, Daniel S. / Williams, Susan G.
Kapitel: 1. Introduction
DOI: <https://doi.org/10.5169/seals-66079>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

TORSION NUMBERS OF AUGMENTED GROUPS WITH APPLICATIONS TO KNOTS AND LINKS

by Daniel S. SILVER and Susan G. WILLIAMS *)

Dedicated to the memory of Arnold E. Ross

ABSTRACT. Torsion and Betti numbers for knots are special cases of more general invariants b_r and β_r , respectively, associated to a finitely generated group G and epimorphism $\chi: G \rightarrow \mathbf{Z}$. The sequence of Betti numbers is always periodic; under mild hypotheses about (G, χ) , the sequence b_r satisfies a linear homogeneous recurrence relation with constant coefficients. Generally, b_r exhibits exponential growth rate. However, again under mild hypotheses, the p -part of b_r has trivial growth for any prime p . Applications to branched cover homology for knots and links are presented.

1. INTRODUCTION

A *knot* is a simple closed curve in the 3-sphere S^3 . Knots are *equivalent* if there is an orientation-preserving homeomorphism of S^3 that carries one into the other. Equivalent knots are regarded as the same. An *invariant* is a well-defined quantity that depends only on a knot equivalence class. Two knots for which some invariant differs are necessarily distinct.

Associated to any knot k and natural number r there is a compact, oriented 3-manifold M_r , the r -fold cyclic cover of S^3 branched over k . A precise definition can be found in [Li97] or [Ro76], for example. Topological invariants of M_r are invariants of k . Two such invariants, the first Betti number β_r and the order b_r of the torsion subgroup of $H_1(M_r; \mathbf{Z})$, were first considered by J. Alexander and G. Briggs [Al28], [AB27] and by O. Zariski [Za32]. The continuing interest in these invariants is witnessed by numerous papers (e.g., [Go72], [Me80], [We80], [Ri90] and [GS91]). We call b_r the r^{th} *torsion*

*) First author partially supported by ÉGIDE at CMI, Université de Provence. Second author partially supported by CNRS at Institut de Mathématiques de Luminy. Both authors partially supported by NSF grant DMS-0071004.

number of k . We say that b_r is *pure* if the corresponding Betti number β_r vanishes (equivalently, $H_1(M_r; \mathbf{Z})$ is a pure torsion group).

Betti numbers are known to be periodic in r , and they are relatively easy to compute (see Proposition 2.2). A useful formula for pure torsion numbers was given by R. Fox in [Fo56]. Although the proof given by Fox was insufficient, a complete argument was given by C. Weber [We80]. Weber observed that the problem of computing non-pure torsion numbers is “... une question plus difficile”.

Torsion and Betti numbers for knots are a special case of a more general, algebraic construction that depends only on an *augmented group*, consisting of a finitely generated group G and a surjection $\chi: G \rightarrow \mathbf{Z}$. We define torsion and Betti numbers in this general context. For a large class of augmented groups, including those that correspond to knots, we provide a formula for all torsion numbers, generalizing the formula of Fox. We prove that the sequence of torsion numbers satisfies a linear recurrence relation.

Torsion numbers tend to grow quickly as their index r becomes large. F. González-Acuña and H. Short [GS91] and independently R. Riley [Ri90] proved that the sequence of pure torsion numbers of any knot k has exponential growth rate equal to the Mahler measure of the Alexander polynomial of k . We improved upon this in [SW00] by showing that the entire sequence b_r grows at this rate and generalizing the result in a natural way for links. The proofs in [SW00] use a deep result about algebraic dynamical systems due to D. Lind, K. Schmidt and T. Ward (Theorem 21.1 of [Sc95]). Here we extend such results for torsion numbers b_r associated to many augmented groups. In contrast, we prove under suitable hypotheses that for any prime number p the p -component of b_r (i.e., the largest power of p that divides b_r) grows subexponentially. The proof relies on a p -adic version of Jensen’s formula, proven by G.R. Everest and B. Ní Fhlathúin [EF96], [Ev99]. As a corollary we strengthen a theorem of C. Gordon [Go72] by proving that for any knot the sequence of torsion numbers either is periodic or else displays infinitely many prime numbers in the factorization of its terms.

In the final section we apply our techniques to the problem of computing homology groups of branched cyclic covering spaces associated to knots and links.

We are grateful to Dan Flath, Adam Sikora, Doug Lind and Hamish Short for useful discussions. The University of Maryland, the Centre de Mathématiques et Informatique in Marseille, and Institut de Mathématiques de Luminy provided kind hospitality during the period of this work. Finally, we thank the referees for helpful comments and suggestions.