

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TORSION NUMBERS OF AUGMENTED GROUPS WITH APPLICATIONS TO KNOTS AND LINKS
Autor: Silver, Daniel S. / Williams, Susan G.
Kapitel: 4. Prime parts of torsion numbers
DOI: <https://doi.org/10.5169/seals-66079>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

4. PRIME PARTS OF TORSION NUMBERS

We recall Jensen's formula, a short argument for which can be found in [Yo86].

LEMMA 4.1 [Jensen's formula]. *For any complex number α ,*

$$\int_0^1 \log |\alpha - e^{2\pi i \theta}| d\theta = \log \max\{1, |\alpha|\}.$$

By Lemma 4.1 the Mahler measure $M(f)$ of a nonzero polynomial with complex coefficients can be computed as

$$\exp \int_0^1 \log |f(e^{2\pi i \theta})| d\theta.$$

This observation motivated the definition of Mahler measure for polynomials in several variables. (See [Bo81] or [EW99], for example.)

In [EF96], [Ev99] G.R. Everest and B.Ní Fhlathúin proved a p -adic analogue of Jensen's formula, which we describe. Assume that α is an algebraic integer lying in a finite extension K of \mathbf{Q} . For every prime p there is a p -adic absolute value $|\cdot|_p$, the usual Archimedean absolute value corresponding to ∞ . We recall the definition (see [La65] for more details): If p is a prime number, then $|p^r m/n|_p = 1/p^r$, where r is an integer, and m, n are nonzero integers that are not divisible by p . By convention, $|0|_p = 0$. Each $|\cdot|_p$ extends to an absolute value $|\cdot|_v$ on K . Let Ω_v denote the smallest field which is algebraically closed and complete with respect to $|\cdot|_v$. Let \mathbf{T}_v denote the closure of the group of all roots of unity, which is in general locally compact. Note that if $p = \infty$, then $\Omega_v = \mathbf{C}$ and $\mathbf{T}_v = \mathbf{T}$. Everest and Fhlathúin define

$$M_{\mathbf{T}_v}(t - \alpha) = \exp \int_{\mathbf{T}_v} \log |t - \alpha|_v d\mu = \exp \lim_{r \rightarrow \infty} \frac{1}{r} \sum_{\zeta^r=1} \log |\zeta - \alpha|_v.$$

Here \int denotes the Shnirelman integral, given by the limit of sums at the right, where one skips over the undefined summands. The above integral exists even if $\alpha \in \mathbf{T}_v$, in which case it can be shown to be zero. Moreover, one has

$$(4.1) \quad \int_{\mathbf{T}_v} \log |t - \alpha|_v d\mu = \log \max\{1, |\alpha|_v\},$$

which Everest and Fhlathúin refer to as a p -adic analogue of Jensen's formula.

Recall that $b_r^{(p)}$ denotes the p -component of b_r , the largest power of p that divides b_r . The *content* of $f \in \mathbf{Z}[t]$ is the greatest common divisor of the coefficients. Using (4.1) we will prove

THEOREM 4.2. *Let (G, χ) be an augmented group, and let p be a prime.*

- (i) *If \mathcal{M} has a square matrix presentation and $\Delta(t) \neq 0$, then the sequence $\{b_{r_k}\}$ of pure torsion numbers satisfies*

$$\lim_{r_k \rightarrow \infty} (b_{r_k}^{(p)})^{1/r_k} = (\text{content } \Delta)^{(p)}.$$

- (ii) *If \mathcal{M} is a direct sum of cyclic modules, then the sequence of all torsion numbers satisfies*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = (\text{content } \Delta)^{(p)}.$$

- (iii) *If \mathcal{M} is torsion free as an abelian group, then*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = 1.$$

EXAMPLE 4.3. For any positive integer m , consider the augmented group (G, χ) where G is the Baumslag-Solitar group $\langle x, y \mid y^m x = x y^m \rangle$ and $\chi: G \rightarrow \mathbf{Z}$ maps $x \mapsto 1$ and $y \mapsto 0$. One verifies that $\mathcal{M} \cong \mathcal{R}_1/(m(t-1))$. The quotient module \mathcal{M}_r is isomorphic to $\mathbf{Z}^r/A_r \mathbf{Z}^r$, where

$$A_r = \begin{pmatrix} m & 0 & 0 & 0 & \cdots & -m \\ -m & m & 0 & \cdots & & 0 \\ 0 & -m & m & 0 & \cdots & 0 \\ & & \vdots & & & \\ 0 & 0 & & \cdots & -m & m \end{pmatrix}.$$

The matrix is equivalent by elementary row and column operations to the diagonal matrix

$$\begin{pmatrix} m & & & & \\ & \ddots & & & \\ & & m & & \\ & & & \ddots & \\ & & & & 0 \end{pmatrix}.$$

Hence $\mathcal{M}_r \cong \mathbf{Z} \oplus (\mathbf{Z}/m)^{r-1}$, and so $b_r = m^{r-1}$ for all r . Consequently,

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = m^{(p)}.$$

The Alexander polynomial of any knot is nonzero, and its coefficients are relatively prime. Hence the following corollary is immediate from Theorem 4.2 (iii).

COROLLARY 4.4. *For any knot k and prime p ,*

$$\lim_{r \rightarrow \infty} (b_r^{(p)})^{1/r} = 1.$$

Theorem 2.10 and Corollary 4.4 imply that whenever the Alexander polynomial of k has Mahler measure greater than 1, infinitely many distinct primes occur in the factorization of the torsion numbers b_r . In other words, the homology groups $H_1(M_r, \mathbf{Z})$ display nontrivial p -torsion for infinitely many primes p . Since the sequence $\{b_r\}$ is a division sequence, the number of prime factors of b_r is unbounded.

What about the case in which the Alexander polynomial of k has Mahler measure equal to 1? The argument of Section 5.7 of [Go72] shows that the number of prime factors remains unbounded as long as the Alexander polynomial does not divide $t^M - 1$ for any M . If it does divide, then the torsion numbers b_r are periodic by Section 5.3 of [Go72] (see also Corollary 2.2 of [SiWi00]). Hence we obtain

COROLLARY 4.5. *For any knot, either the torsion numbers b_r are periodic or else for any $N > 0$ there exists an r such that the factorization of b_r has at least N distinct primes.*

The proof of Theorem 4.2 requires the following lemma.

LEMMA 4.6. *If $f(t) = c_0 t^n + \cdots + c_{n-1} t + c_n$ is a nonzero polynomial in $\mathbf{Z}[t]$ with roots $\lambda_1, \dots, \lambda_n$ (not necessarily distinct) in Ω_v , then*

$$|c_0|_v \prod_{i=1}^n \max\{1, |\lambda_i|_v\} = |\text{content } f|_v.$$

Proof. The argument that we present is found in [LW88]. Set $a_j = c_j/c_0$ for $0 \leq j \leq n$, so $f(t) = c_0(t^n + a_1 t^{n-1} + \cdots + a_n)$. Each a_j is an elementary symmetric function of the roots λ_i , namely the sum of products of roots taken j at a time. Using the ultrametric property

$$|x + y|_v = \max\{|x|_v, |y|_v\},$$

we see that if exactly k values of $|\lambda_i|_v$ are greater than 1, then

$$\max_j |a_j|_v = |a_k|_v = \prod_{j=1}^k \max\{1, |\lambda_j|_v\}.$$

But

$$\max_j |a_j|_v = \max \left\{ 1, \left| \frac{c_1}{c_0} \right|_v, \dots, \left| \frac{c_n}{c_0} \right|_v \right\} = \frac{|\text{content } f|_v}{|c_0|_v}.$$

Hence the lemma is proved. \square

Proof of Theorem 4.2. In case (i), the pure torsion number b_{r_k} is equal to $\left| \prod_{\zeta^{r_k}=1} \Delta(\zeta) \right|_v$. We have

$$|b_{r_k}|_v = \left| \prod_{\zeta^{r_k}=1} \Delta(\zeta) \right|_v = |c_0|_v^{r_k} \prod_{\zeta^{r_k}=1} \prod_{j=1}^n |\zeta - \lambda_j|_v,$$

where c_0 is the leading coefficient of Δ and $\lambda_1, \dots, \lambda_n$ are its roots. Hence

$$\begin{aligned} |b_{r_k}|_v^{1/r_k} &= |c_0|_v \prod_{\zeta^{r_k}=1} \prod_{j=1}^n |\zeta - \lambda_j|_v^{1/r_k} \\ &= |c_0|_v \prod_{j=1}^n \exp \left(\frac{1}{r_k} \sum_{\zeta^{r_k}=1} \log |\zeta - \lambda_j| \right), \end{aligned}$$

so that

$$\lim_{r_k \rightarrow \infty} |b_{r_k}|_v^{1/r_k} = |c_0|_v \prod_{j=1}^n \exp \int_{\mathbf{T}_v} \log |t - \lambda_j|_v d\mu,$$

which by equation (4.1) is equal to

$$|c_0|_v \prod_{j=1}^n \max \{1, |\lambda_j|_v\}.$$

By Lemma 4.6 this is equal to $|\text{content } \Delta|_v$. But for integers n we have $n^{(p)} = |n|_v^{-1}$.

Now suppose \mathcal{M} is cyclic. As in the proof of Theorem 3.8, we let γ be the cyclotomic order of Δ and consider the subsequence of b_r with r in a fixed congruence class modulo γ . Then starting with Theorem 3.3 we may apply the argument above with Δ/Φ in place of Δ to show that the limit of $(|b_r|^{(p)})^{1/r}$ along this subsequence is the p -component of the content of Δ/Φ . But content is multiplicative and cyclotomic polynomials have content 1, so the limit along all congruence classes is $(\text{content } \Delta)^{(p)}$. The result is immediate for direct sums of cyclic modules.

Finally, we can extend the result when \mathcal{M} is torsion-free as an abelian group using Theorem 3.6. But for this case the content of Δ is 1. \square