

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE NONAMENABILITY OF SCHREIER GRAPHS FOR INFINITE INDEX QUASICONVEX SUBGROUPS OF HYPERBOLIC GROUPS
Autor: Kapovich, Ilya
Kurzfassung
DOI: <https://doi.org/10.5169/seals-66081>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE NONAMENABILITY OF SCHREIER GRAPHS
FOR INFINITE INDEX QUASICONVEX
SUBGROUPS OF HYPERBOLIC GROUPS

by Ilya KAPOVICH

ABSTRACT. We show that if H is a quasiconvex subgroup of infinite index in a nonelementary hyperbolic group G then the Schreier coset graph for G relative to H is nonamenable.

1. INTRODUCTION

A connected graph of bounded degree X is *nonamenable* if X has nonzero Cheeger constant or, equivalently, if the spectral radius of the simple random walk on X is less than one (see Section 2 below for more precise definitions). Nonamenable graphs play an increasingly important role in the study of various probabilistic phenomena, such as random walks, harmonic analysis, Brownian motion, and percolations on graphs and manifolds (see for example [2, 5, 6, 7, 15, 17, 18, 24, 30, 43, 44, 62, 71, 72]), as well as in the study of expander families of finite graphs (see for example [52, 66, 67]).

It is well-known that a finitely generated group G is nonamenable if and only if the Cayley graph of G with respect to some (any) finite generating set is nonamenable. The notion of a *word-hyperbolic group* was introduced by M. Gromov [40] and has played a central role in Geometric Group Theory for the last fifteen years. Word-hyperbolic groups are nonamenable unless they are virtually cyclic. Thus the Cayley graphs of word-hyperbolic groups provide a large and interesting class of nonamenable graphs. In this paper we investigate nonamenability of Schreier coset graphs corresponding to subgroups of hyperbolic groups.