Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 48 (2002)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE NONAMENABILITY OF SCHREIER GRAPHS FOR INFINITE INDEX QUASICONVEX SUBGROUPS OF HYPERBOLIC GROUPS

Autor: Kapovich, Ilya

Bibliographie

DOI: https://doi.org/10.5169/seals-66081

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 13.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

We can now obtain Corollary 1.4 stated in the Introduction.

COROLLARY 5.4. *Let $G = \langle x_1, \dots, x_k \mid r_1, \dots, r_m \rangle$ be a nonelementary word-hyperbolic group and let $H \leq G$ be a quasiconvex subgroup of infinite index. Let a_n be the number of freely reduced words in $A = \{x_1, \dots, x_k\}^{\pm 1}$ of length n that represent elements of H. Let b_n be the number of all words in A of length n that represent elements of H. Then*

$$\limsup_{n \to \infty} \sqrt[n]{a_n} < 2k - 1$$

and

$$\limsup_{n \to \infty} \sqrt[n]{b_n} < 2k .$$

Proof. Note that $k \geq 2$ since G is nonelementary. Put $A = \{x_1, \dots, x_k\}$ and $Y = \Gamma(G, H, A)$. We choose $x_0 := H1 \in VY$ as the base-vertex of Y. Note that Y is $2k$-regular by construction. Also, for any vertex x of Y and any word w in $A \cup A^{-1}$ there is a unique path in Y with label w and origin x. The definition of Schreier coset graphs also implies that a word w represents an element of H if and only if the unique path in Y with origin x_0 and label w terminates at x_0. Therefore $a_n(Y)$ equals the number of freely reduced words in the alphabet $A = \{x_1, \dots, x_k\}^{\pm 1}$ of length n that represent elements of H. Similarly, $b_n(Y)$ equals the number of all words in A of length n representing elements of H. By Theorem 1.2, Y is nonamenable. Hence by Theorem 2.5, $\alpha(Y) < 2k - 1$ and $\beta(Y) < 2k$, as required.

REFERENCES

[1] ALONSO, J., T. BRADY, D. COOPER, V. FERLINI, M. LUSTIG, M. MIHALIK, M. SHAPIRO and H. SHORT. Notes on hyperbolic groups. In: *Group Theory from a Geometrical Viewpoint*, Proceedings of the workshop held in Trieste, É. Ghys, A. Haefliger and A. Verjovsky (editors). World Scientific Publishing Co., 1991.

[2] ANCONA, A. *Théorie du potentiel sur les graphes et les variétés*, École d'été de Probabilités de Saint-Flour XVIII–1988, 1–112. Lecture Notes in Mathematics *1427*. Springer, Berlin, 1990.

[3] ARZHANTSEVA, G. On quasiconvex subgroups of word hyperbolic groups. *Geom. Dedicata 87* (2001), 191–208.

[4] BAUMSLAG, G., S. M. GERSTEN, M. SHAPIRO and H. SHORT. Automatic groups and amalgams. *J. Pure Appl. Algebra 76* (1991), 229–316.

[5] BARTHOLDI, L. Counting paths in graphs. *L'Enseignement Math. (2) 45* (1999), 83–131.

[6] BENJAMINI, I., R. LYONS and O. SCHRAMM. Percolation perturbations in potential theory and random walks. In: *Random Walks and Discrete Potential Theory (Cortona, 1997)*, 56–84. Sympos. Math. XXXIX, Cambridge Univ. Press, Cambridge, 1999.

[7] BENJAMINI, I. and O. SCHRAMM. Every graph with a positive Cheeger constant contains a tree with a positive Cheeger constant. *Geom. Funct. Anal. 7* (1997), 403–419.

[8] BESTVINA, M. and M. FEIGHN. A combination theorem for negatively curved groups. *J. Differential Geom. 35* (1992), 85–101.

[9] BLOCK, J. and S. WEINBERGER. Aperiodic tilings, positive scalar curvature and amenability of spaces. *J. Amer. Math. Soc. 5* (1992), 907–918.

[10] BOROVIK, A., A. G. MYASNIKOV and V. REMESLENNIKOV. Multiplicative measures on free groups. To appear in *Internat. J. Algebra Comput.*

[11] BOWDITCH, B. Cut points and canonical splittings of hyperbolic groups. *Acta Math. 180* (1998), 145–186.

[12] BOWERS, P. Negatively curved graph and planar metrics with applications to type. *Michigan Math. J. 45* (1998), 31–53.

[13] BRINKMANN, P. Hyperbolic automorphisms of free groups. *Geom. Funct. Anal. 10* (2000), 1071–1089.

[14] CANNON, J. W. The theory of negatively curved spaces and groups. In: *Ergodic Theory, Symbolic Dynamics, and Hyperbolic Spaces (Trieste, 1989)*, 315–369. Oxford Sci. Publ., Oxford Univ. Press, New York, 1991.

[15] CAO, J. Cheeger isoperimetric constants of Gromov-hyperbolic spaces with quasi-poles. *Commun. Contemp. Math. 2* (2000), 511–533.

[16] CECCHERINI-SILBERSTEIN, T., R. GRIGORCHUCK and P. DE LA HARPE. Amenability and paradoxical decompositions for pseudogroups and discrete metric spaces (in Russian). *Tr. Mat. Inst. Steklova 224* (1999). Algebra. Topol. Differ. Uravn. i ikh Prilozh., 68–111; translation in *Proc. Steklov Inst. Math. 224* (1999), 57–97.

[17] CHUNG, F. R. K. Laplacians of graphs and Cheeger's inequalities. In: *Combinatorics, Paul Erdős is Eighty, Vol. 2 (Keszthely, 1993)*, 157–172. Bolyai Soc. Math. Stud. 2. János Bolyai Math. Soc., Budapest, 1996.

[18] CHUNG, F. R. K. and K. ODEN. Weighted graph Laplacians and isoperimetric inequalities. *Pacific J. Math. 192* (2000), 257–273.

[19] COHEN, J. Cogrowth and amenability of discrete groups. *J. Funct. Anal. 48* (1982), 301–309.

[20] COORNAERT, M., T. DELZANT, and A. PAPADOPOULOS. *Géométrie et théorie des groupes. Les groupes hyperboliques de Gromov.* Lecture Notes in Mathematics, *1441*. Springer-Verlag, Berlin, 1990.

[21] DODZIUK, J. Difference equations, isoperimetric inequality and transience of certain random walks. *Trans. Amer. Math. Soc. 284* (1984), 787–794.

[22] DUNWOODY, M. and M. SAGEEV. JSJ-splittings for finitely presented groups over slender groups. *Invent. Math. 135* (1999), 25–44.

[23] DUNWOODY, M. and E. SWENSON. The algebraic torus theorem. *Invent. Math. 140* (2000), 605–637.

[24] ELEK, G. Amenability, l_p-homologies and translation invariant functionals. *J. Austral. Math. Soc. Ser. A 65* (1998), 111–119.

[25] EPSTEIN, D., J. CANNON, D. HOLT, S. LEVY, M. PATERSON and W. THURSTON. *Word Processing in Groups.* Jones and Bartlett, Boston, 1992.

[26] EPSTEIN, D. and D. HOLT. Efficient computation in word-hyperbolic groups. In: *Computational and Geometric Aspects of Modern Algebra (Edinburgh, 1998)*, 66–77. LMS Lecture Note Ser. *275*. Cambridge Univ. Press, Cambridge, 2000.

[27] FOORD, R. Automaticity and growth in certain classes of groups and monoids. PhD Thesis, Warwick University, 2000.

[28] FUJIWARA, K. and P. PAPASOGLU. JSJ decompositions and complexes of groups. Preprint, 1996.

[29] GERASIMOV, V.N. Semi-splittings of groups and actions on cubings. In: *Algebra, Geometry, Analysis and Mathematical Physics (Novosibirsk, 1996)*, 91–109, 190. Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 1997.

[30] GERL, P. Amenable groups and amenable graphs. In: *Harmonic Analysis (Luxembourg, 1987)*, 181–190. Lecture Notes in Mathematics *1359*. Springer, Berlin, 1988.

[31] GERSTEN, S.M. and H. SHORT. Rational subgroups of biautomatic groups. *Ann. of Math. (2) 134* (1991), 125–158.

[32] GHYS, E. and P. DE LA HARPE (eds). *Sur les groupes hyperboliques d'après Mikhael Gromov.* Progress in Mathematics series *83*. Birkhäuser, 1990.

[33] GITIK, R. On the combination theorem for negatively curved groups. *Internat. J. Algebra Comput. 6* (1996), 751–760.

[34] —— On quasiconvex subgroups of negatively curved groups. *J. Pure Appl. Algebra 119* (1997), 155–169.

[35] —— On the profinite topology on negatively curved groups. *J. Algebra 219* (1999), 80–86.

[36] —— Doubles of groups and hyperbolic LERF 3-manifolds. *Ann. of Math. (2) 150* (1999), 775–806.

[37] —— Tameness and geodesic cores of subgroups. *J. Austral. Math. Soc. Ser. A 69* (2000), 153–16.

[38] GITIK, R., M. MITRA, E. RIPS and M. SAGEEV. Widths of subgroups. *Trans. Amer. Math. Soc. 350* (1998), 321–329.

[39] GRIGORCHUK, R.I. Symmetrical random walks on discrete groups. In: *Multicomponent Random Systems*, 285–325. Adv. Probab. Related Topics *6*. Dekker, New York, 1998.

[40] GROMOV, M. Hyperbolic groups. In: *Essays in Group Theory*, (S.M. Gersten, ed.), 75–263. Math. Sci. Res. Inst. Publ. *8*, 1987.

[41] —— Asymptotic invariants of infinite groups. In: *Geometric Group Theory*, Vol. 2 (Sussex, 1991), 1–295. LMS Lecture Note Ser. *182*. Cambridge Univ. Press, Cambridge, 1993.

[42] HOLT, D. Automatic groups, subgroups and cosets. In: *The Epstein Birthday Schrift*, 249–260. Geom. Topol. Monogr. *1*. Geom. Topol., Coventry, 1998.

[43] KAIMANOVICH, V. Equivalence relations with amenable leaves need not be amenable. In: *Topology, Ergodic Theory, Real Algebraic Geometry*, 151–166. Amer. Math. Soc. Transl. Ser. 2 *202*. Amer. Math. Soc., Providence (R.I.), 2001.

[44] KAIMANOVICH, V. and W. WOESS. The Dirichlet problem at infinity for random walks on graphs with a strong isoperimetric inequality. *Probab. Theory Related Fields 91* (1992), 445–466.

[45] KAPOVICH, I. Detecting quasiconvexity: algorithmic aspects. In: *Geometric and Computational Perspectives on Infinite Groups (Minneapolis, MN and New Brunswick, NJ, 1994)*, 91–99. DIMACS Ser. Discrete Math. Theoret. Comput. Sci. *25*. Amer. Math. Soc., Providence (R.I.), 1996.

[46] —— Quasiconvexity and amalgams. *Internat. J. Algebra Comput. 7* (1997), 771–811.

[47] —— A non-quasiconvexity embedding theorem for word-hyperbolic groups. *Math. Proc. Cambridge Philos. Soc. 127* (1999), 461–486.

[48] —— The combination theorem and quasiconvexity. *Internat. J. Algebra Comput. 11* (2001), 185–216.

[49] —— The geometry of relative Cayley graphs for subgroups of hyperbolic groups. Preprint, 2002.

[50] KAPOVICH, I., A. MYASNIKOV, P. SCHUPP and V. SHPILRAIN. Generic-case complexity, decision problems in group theory and random walks. Preprint, 2002.

[51] KAPOVICH, I. and H. SHORT. Greenberg's theorem for quasiconvex subgroups of word hyperbolic groups. *Canad. J. Math. 48* (1996), 1224–1244.

[52] LUBOTZKY, A. *Discrete Groups, Expanding Graphs and Invariant Measures.* (With an appendix by Jonathan D. Rogawski.) Progress in Mathematics *125*. Birkhäuser Verlag, Basel, 1994.

[53] MIHALIK, M. Group extensions and tame pairs. *Trans. Amer. Math. Soc. 351* (1999), 1095–1107.

[54] MIHALIK, M. and W. TOWLE. Quasiconvex subgroups of negatively curved groups. *J. Pure Appl. Algebra 95* (1994), 297–301.

[55] MITRA, M. Cannon-Thurston maps for trees of hyperbolic metric spaces. *J. Differential Geom. 48* (1998), 135–164.

[56] NEUMANN, B. H. Groups covered by finitely many cosets. *Publ. Math. Debrecen 3* (1954), 227–242.

[57] REEVES, L. Rational subgroups of cubed 3-manifold groups. *Michigan Math. J. 42* (1995), 109–126.

[58] RIPS, E. Subgroups of small cancellation groups. *Bull. London Math. Soc. 14* (1982), 45–47.

[59] RIPS, E. and Z. SELA. Cyclic splittings of finitely presented groups and the canonical JSJ decomposition. *Ann. of Math. (2) 146* (1997), 53–109.

[60] SAGEEV, M. Ends of group pairs and non-positively curved cube complexes. *Proc. London Math. Soc. (3) 71* (1995), 585–617.

[61] —— Codimension-1 subgroups and splittings of groups. *J. Algebra 189* (1997), 377–389.

[62] SCHONMANN, R. Multiplicity of phase transitions and mean-field criticality on highly non-amenable graphs. *Comm. Math. Phys. 219* (2001), 271–322.

[63] SCOTT, G. P. and G. A. SWARUP. An algebraic annulus theorem. *Pacific J. Math. 196* (2000), 461–506.

[64] SCOTT, G. P. and G. A. SWARUP. Canonical splittings of groups and 3-manifolds. *Trans. Amer. Math. Soc. 353* (2001), 4973–5001.

[65] SELA, Z. Structure and rigidity in (Gromov) hyperbolic groups and discrete groups in rank 1 Lie groups. II. *Geom. Funct. Anal. 7* (1997), 561–593.

[66] SHALOM, Y. Random ergodic theorems, invariant means and unitary representation. In: *Lie Groups and Ergodic Theory (Mumbai, 1996)*, 273–314. Tata Inst. Fund. Res. Stud. Math. *14*. Tata Inst. Fund. Res., Bombay, 1998.

[67] —— Expander graphs and amenable quotients. In: *Emerging Applications of Number Theory (Minneapolis, 1996)*, 571–581. IMA Vol. Math. Appl. *109*. Springer, New York, 1999.

[68] SHORT, H. Quasiconvexity and a theorem of Howson's. In: *Group Theory from a Geometrical Viewpoint (Trieste, 1990)*, 168–176. World Sci. Publishing, River Edge (N.J.), 1991.

[69] SWARUP, G. A. Geometric finiteness and rationality. *J. Pure Appl. Algebra 86* (1993), 327–333.

[70] —— Proof of a weak hyperbolization theorem. *Quarterly J. Math. 51* (2000), 529–533.

[71] WOESS, W. Random walks on infinite graphs and groups – a survey on selected topics. *Bull. London Math. Soc. 26* (1994), 1–60.

[72] —— *Random Walks on Infinite Graphs and Groups.* Cambridge Tracts in Mathematics *138*. Cambridge University Press, Cambridge, 2000.

[73] ZUK, A. On property (T) for discrete groups. Preprint, 2001.

(Reçu le 22 avril 2002)

Ilya Kapovich

Department of Mathematics
University of Illinois at Urbana-Champaign
1409 West Green Street, Urbana, IL 61801
U. S. A.
e-mail: kapovich@math.uiuc.edu

vide-leer-empty