

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TRIPLE RATIO ON THE UNITARY STIEFEL MANIFOLD
Autor: Clerc, Jean-Louis
Kapitel: 1. Geometric setting
DOI: <https://doi.org/10.5169/seals-66067>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

is the unit ball in the matrix space $\text{Mat}(p \times q, \mathbf{C})$, S is the unitary Stiefel manifold $S_{p,q}$ and $G = \text{PSU}(p, q)$. The invariant we construct for triples is of matrix-valued nature (it is a conjugacy class) and we give two versions of it (see Theorems 4.3 and 4.4). The basic strategy is to approach the Shilov boundary from inside. The (matrix-valued) *automorphy kernel* for the domain D is used to build a kernel for triples of points inside D which transforms nicely under the action of G . It remains to look carefully at the boundary behaviour of the kernel when the points approach the Shilov boundary S . This is only possible for triples satisfying a generic condition called *transversality* (see Proposition 2.1 for a definition). The *Cayley transform* plays an important role in the proofs. Finally the problem is reduced to a *linear* problem, which is related to the description of some orbits for the action $(g, X) \mapsto gXg^*$ of GL_q on $\text{Mat}(q \times q, \mathbf{C})$ (see Theorem 3.9).

For general references on bounded symmetric domains and their geometric properties, see [S], and Part III in [Fal]. For explicit calculations related to our example, see [P] and [H].

1. GEOMETRIC SETTING

Let p, q be two integers with $1 \leq q \leq p$, and let

$$(1) \quad D = \{z \in \text{Mat}(p \times q, \mathbf{C}) \mid \mathbf{1}_q - z^*z \gg 0\}.$$

Let $G = \text{SU}(p, q) \subset \text{GL}(p + q, \mathbf{C})$. An element $g \in \text{GL}(p + q, \mathbf{C})$ will often be written as

$$g = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where

$$a \in \text{Mat}(p \times p, \mathbf{C}), \quad b \in \text{Mat}(p \times q, \mathbf{C}), \quad c \in \text{Mat}(q \times p, \mathbf{C}), \quad d \in \text{Mat}(q \times q, \mathbf{C}).$$

In this notation, the conditions for g to belong to $\text{U}(p, q, \mathbf{C})$ can be written as

$$(2) \quad \begin{aligned} a^*a - c^*c &= \mathbf{1}_p \\ b^*a - d^*c &= 0 \\ d^*d - b^*b &= \mathbf{1}_q. \end{aligned}$$

Define an action of the group $\text{GL}(p + q, \mathbf{C})$ on $\text{Mat}(p \times q, \mathbf{C})$ by

$$(3) \quad g(z) = (az + b)(cz + d)^{-1}.$$

The action is not everywhere defined, but it is certainly defined if $g \in G$ and $z \in D$. It defines an action of G on D , and G (or rather $\mathbf{PSU}(p, q)$) is the neutral component of the group of all biholomorphic transformations of D .

The stabilizer of the base point $0 \in D$ is the maximal compact subgroup $K = S(\mathrm{U}(p) \times \mathrm{U}(q))$. Its complexification is the complex group $K^{\mathbb{C}} = S(\mathrm{GL}(p, \mathbb{C}) \times \mathrm{GL}(q, \mathbb{C}))$. We also define the following subgroups

$$P^+ = \left\{ \begin{pmatrix} \mathbf{1}_p & z \\ 0 & \mathbf{1}_q \end{pmatrix}, z \in \mathrm{Mat}(p \times q, \mathbb{C}) \right\}$$

$$P^- = \left\{ \begin{pmatrix} \mathbf{1}_p & 0 \\ w & \mathbf{1}_q \end{pmatrix}, w \in \mathrm{Mat}(q \times p, \mathbb{C}) \right\}.$$

The corresponding *Harish Chandra decomposition* is the following identity

$$(4) \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \mathbf{1}_p & bd^{-1} \\ 0 & \mathbf{1}_q \end{pmatrix} \begin{pmatrix} a - bd^{-1}c & 0 \\ 0 & d \end{pmatrix} \begin{pmatrix} \mathbf{1}_p & 0 \\ d^{-1}c & \mathbf{1}_q \end{pmatrix}$$

valid for $g \in \mathrm{GL}(p+q, \mathbb{C})$ if d is invertible.

The *automorphy kernel* $k(z, w)$ is defined for $z, w \in \mathrm{Mat}(p \times q, \mathbb{C})$ wherever it makes sense by the formula

$$(5) \quad k(z, w) = (\mathbf{1}_q - w^* z)^{-1}.$$

In particular it is always well defined for $z, w \in D$ and has values in $\mathrm{GL}(q, \mathbb{C})$. It has the following law of transformation for $g \in G$

$$(6) \quad k(g(z), g(w)) = j(g, z) k(z, w) j(g, w)^*,$$

where

$$(7) \quad j(g, z) = cz + d.$$

The *Shilov boundary* of D is the unitary Stiefel manifold S defined by

$$(8) \quad S = \left\{ \sigma \in \mathrm{Mat}(p \times q, \mathbb{C}) \mid \sigma^* \sigma = \mathbf{1}_q \right\}.$$

The action of G extends to S , and it is clearly transitive on S . In fact the action of K is already transitive.

To go further, we need to make a specific choice of a base point in S . For this we first systematically write elements in $\mathrm{Mat}(p \times q, \mathbb{C})$ as

$$z = \begin{pmatrix} z_q \\ z' \end{pmatrix}$$

where $z_q \in \mathrm{Mat}(q \times q, \mathbb{C})$ and $z' \in \mathrm{Mat}((p-q) \times q, \mathbb{C})$. With this convention, let $ie = \begin{pmatrix} i\mathbf{1}_q \\ 0 \end{pmatrix}$ be the base point in S . Associated to this choice is the Cayley transform c , given by

$$z = \begin{pmatrix} z_q \\ z' \end{pmatrix} \longmapsto c(z) = \begin{pmatrix} w_q \\ w' \end{pmatrix}$$

with

$$(9) \quad \begin{aligned} w_q &= (z_q + i\mathbf{1}_q)(iz_q + \mathbf{1}_q)^{-1} \\ w' &= -z'(iz_q + \mathbf{1}_q)^{-1}. \end{aligned}$$

The inverse of the Cayley transform is the map which to $\begin{pmatrix} w_q \\ w' \end{pmatrix}$ associates the matrix $\begin{pmatrix} z_q \\ z' \end{pmatrix}$ given by

$$(10) \quad \begin{aligned} z_q &= (iw_q - \mathbf{1}_q)^{-1}(i\mathbf{1}_q - w_q) \\ z' &= 2w'(iw_q - \mathbf{1}_q)^{-1}. \end{aligned}$$

The Cayley transform is a rational map, well defined on D . The image of D is the *Siegel domain of type II* defined by

$$(11) \quad {}^cD = \left\{ \begin{pmatrix} w_q \\ w' \end{pmatrix}, \frac{1}{2i}(w_q - w_q^*) - w'^*w' \gg 0 \right\}$$

and the image of the Shilov boundary (more exactly the part of the Shilov boundary where the Cayley transform is defined) is

$$(12) \quad {}^cS = \left\{ \begin{pmatrix} w_q \\ w' \end{pmatrix}, \frac{1}{2i}(w_q - w_q^*) = w'^*w' \right\}.$$

To the data

$$\begin{aligned} w_0 &\in \text{Mat}((p-q) \times q, \mathbf{C}) \\ h &\in \text{GL}(q, \mathbf{C}), u \in \text{U}(p-q, \mathbf{C}), \text{ such that } \det h = (\det u)^{-1} \\ s &\in \text{Herm}(q, \mathbf{C}) \end{aligned}$$

we associate the transform

$$(13) \quad \begin{aligned} w_q &\longmapsto h^*w_qh + s + 2iw_0^*uw'h + iw_0^*w_0 \\ w' &\longmapsto uw'h + w_0. \end{aligned}$$

Any such transform maps cD in a one-to-one fashion into itself. These transforms form a group and it is exactly the group of affine holomorphic transforms of the domain cD .

Let B be the stabilizer of the point ie in G . The conjugate group under the Cayley transform is ${}^cB = c \circ B \circ c^{-1}$ and it turns out to be exactly the group of affine transforms of cD we just described. Observe that the group cB is transitive on cD and on cS .