

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TRIPLE RATIO ON THE UNITARY STIEFEL MANIFOLD
Autor: Clerc, Jean-Louis
Kapitel: 2. Action of G on $S \times S$ and $S \times S \times S$
DOI: <https://doi.org/10.5169/seals-66067>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. ACTION OF G ON $S \times S$ AND $S \times S \times S$

We now study the action of G on pairs of points of S . The main notion to be introduced is *transversality*, a notion that could be defined for any bounded symmetric domain. We give several equivalent definitions for our case.

PROPOSITION 2.1. *Let σ and ξ be two elements of S . Then the following are equivalent:*

- (i) $\det(\mathbf{1}_q - \xi^* \sigma) \neq 0$;
- (ii) $\xi - \sigma$ injective;
- (iii) $\det(\mathbf{1}_p - \xi \sigma^*) \neq 0$.

If one of these equivalent conditions is satisfied, then σ and ξ are said to be transverse.

Proof. Assume (i). As $\mathbf{1}_q = \xi^* \xi$, this condition amounts to $\det(\xi^*(\xi - \sigma)) \neq 0$, which in particular shows that $\xi - \sigma$ is injective. Conversely, assume $\xi - \sigma$ is injective and let $v \in \mathbf{C}^q$ be such that $v = \xi^* \sigma v$. Now

$$\|v\| = \|\xi^* \sigma v\| \leq \|\sigma v\| \leq \|v\|,$$

and hence $\|\xi^* \sigma v\| = \|\sigma v\|$, which is possible only if $\sigma v \in \text{Im } \xi$. So there exists $w \in \mathbf{C}^q$, such that $\sigma v = \xi w$. But taking the image of both sides by ξ^* yields $v = w$, and hence $\sigma v = \xi v$, so that $v = 0$. So $\mathbf{1}_q - \xi^* \sigma$ is injective and hence (ii) \Rightarrow (i). Under the same assumption (ii), let us prove that $\xi \sigma^*$ cannot have 1 as an eigenvalue. Suppose $v \in \mathbf{C}^p$ is such that $\xi \sigma^* v = v$. As ξ is a partial isometry, this forces $\|\sigma^* v\| = \|v\|$, and hence v belongs to the image of the map σ , so there exists $w \in \mathbf{C}^q$ such that $v = \sigma w$. But then we also have $v = \xi \sigma^* \sigma w = \xi w$ and hence $(\sigma - \xi)w = 0$ which forces $w = 0$. Hence (iii) follows from (ii). Finally assume (iii). Then as σ is injective, $(\mathbf{1}_p - \xi \sigma^*) \circ \sigma = \sigma - \xi$ is also injective. Hence (iii) \Rightarrow (ii). \square

We will use the notation $\sigma \top \xi$ to denote transversality. It is a symmetric condition. It is invariant under the action of G , as can easily be concluded from (6). For $\sigma \in S$, let

$$S_{\top}^{\sigma} = \{\xi \mid \sigma \top \xi\}.$$

Observe that the set S_{\top}^{ie} is exactly the subset in S where the Cayley transform is defined.

Let

$$(14) \quad S_{\top}^2 = \{(\sigma, \xi) \in S \times S \mid \sigma \top \xi\}.$$

As base point in S_{\top}^2 we choose $(ie, -ie)$. Observe that $c(-ie) = 0$.

THEOREM 2.2. *The group G acts transitively on S_{\top}^2 .*

Proof. Let $(\sigma, \xi) \in S_{\top}^2$ and let us show that there exists an element of G which maps (σ, ξ) to $(ie, -ie)$. As G is transitive on S , we may assume that $\sigma = ie$. Then the transversality condition shows that ξ belongs to the domain of the Cayley transform. The element $c(\xi)$ belongs to ${}^c S$, and we have already noticed that ${}^c B$ is transitive on ${}^c S$. Hence $c(\xi)$ can be mapped to $0 = c(-ie)$. Taking the image under the inverse Cayley transform gives the result. \square

Denote by L the stabilizer of the base point $(ie, -ie)$ in B . Under a Cayley transform, the group ${}^c L = c \circ L \circ c^{-1}$ is the stabilizer in ${}^c B$ of the element 0 . Hence it is the subgroup of linear transformations given by

$$\begin{aligned} w_q &\longmapsto h^* w_q h \\ w' &\longmapsto uw h \end{aligned}$$

where $h \in \mathrm{GL}(q, \mathbf{C})$, $u \in \mathrm{U}(p-q)$ and $\det h = (\det u)^{-1}$.

LEMMA 2.3. *Let $\begin{pmatrix} w_q \\ w' \end{pmatrix}, \begin{pmatrix} v_q \\ v' \end{pmatrix} \in {}^c S$. Then they belong to the same orbit under the action of ${}^c L$ if and only if w_q and v_q belong to the same orbit under the action of $\mathrm{GL}(q, \mathbf{C})$.*

Proof. One implication being trivial, we only have to prove the other one. So assume there exists $h \in \mathrm{GL}(q, \mathbf{C})$ such that $v_q = h^* w_q h$. Let μ be a complex number such that $\mu^{p-q} = \det h$ and let $u = \mu^{-1} \mathbf{1}_{p-q}$. Clearly $(\det u)^{-1} = \det h$. Using the action of (h, u) we may assume that $v_q = w_q$. Let $s_q = \frac{1}{2i}(w_q - w_q^*)$. This is an Hermitian matrix and as w_q and v_q belong to ${}^c S$, we get

$$s_q = w'^* w' = v'^* v'.$$

Looking to the columns of w' (or v'), we may think of w' as a family of q vectors in \mathbf{C}^{p-q} . Then the matrix s_q is the Gram matrix of these vectors. But two sets of vectors in \mathbf{C}^{p-q} are conjugate under the action of the unitary group $\mathrm{U}(p-q)$ if and only if they have the same Gram matrix. Hence there exists $u \in \mathrm{U}(p-q)$ such that $v' = uw'$. Let λ be a complex number such that $\lambda^q = \det u$. Then using the action of $(\lambda^{-1} \mathbf{1}_q, u)$, we get the result. \square

Let us denote by H_q the real vector space of $q \times q$ Hermitian matrices, and let Ω_q be the subset of all positive-definite matrices. For any integer r such that $0 \leq r \leq q$ let $\Omega_q^{(r)}$ be the set of all positive semi-definite $q \times q$ Hermitian matrices of rank less than r . For $r < q$, the set $\Omega_q^{(r)}$ is contained in the boundary of Ω_q , whereas for $r = q$, $\Omega_q^{(q)} = \overline{\Omega}_q$.

Let

$$T_q^{(r)} = \{x + iy \mid x \in H_q, y \in \Omega_q^{(r)}\}.$$

The group $\mathrm{GL}(q, \mathbf{C})$ acts on $T_q^{(r)}$ by the action $(h, w) \mapsto hwh^*$.

Finally let

$$\tilde{T}_q^{(r)} = \{z \in T_q^{(r)} \mid z \text{ invertible}\}.$$

Clearly the action of $\mathrm{GL}(q, \mathbf{C})$ preserves $\tilde{T}_q^{(r)}$.

Let $\begin{pmatrix} w_q \\ w' \end{pmatrix}$ be in ${}^c S$. Then $w_q = x_q + iw'^*w'$, with $x_q \in H_q$. Let

$$r = \inf(q, p - q).$$

The rank of the matrix w'^*w' is at most r . Hence w_q belongs to $T_q^{(r)}$. Conversely, it is easily seen that any positive semi-definite Hermitian matrix of rank at most r can be written as w'^*w' for some $w' \in \mathrm{Mat}((p-q) \times q, \mathbf{C})$.

Let

$$(15) \quad S_{\top}^3 = \{(\sigma_1, \sigma_2, \sigma_3) \in S \times S \times S \mid \sigma_1 \top \sigma_2, \sigma_2 \top \sigma_3, \sigma_3 \top \sigma_1\}.$$

THEOREM 2.4. *The G -orbits in S_{\top}^3 are in one-to-one correspondance with the orbits of $\mathrm{GL}(q, \mathbf{C})$ in $\tilde{T}_q^{(r)}$.*

Proof. From Theorem 2.2 we already know that any orbit contains an element of the form $(ie, -ie, \sigma)$ with $\sigma \in S$. Now use the Cayley transform. The element $w = c(\sigma)$ is in ${}^c S$, and the transversality condition is equivalent to the condition $\det(w_q) \neq 0$. In other words, $w_q \in \tilde{T}_q^{(r)}$. The result now follows from Lemma 2.3. \square