Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 48 (2002)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: TRIPLE RATIO ON THE UNITARY STIEFEL MANIFOLD

Autor: Clerc, Jean-Louis

Kapitel: 3. Orbits for the \$GL_q\$-action on \$\tilde{T}_q\$

DOI: https://doi.org/10.5169/seals-66067

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. Orbits for the GL_q -action on \widetilde{T}_q

Any $z \in \text{Mat}(q \times q, \mathbb{C})$ can be written in a unique way as z = x + iy, with $x, y \in H_q$. We will be concerned with the set \widetilde{T}_q defined by

(16)
$$\widetilde{T}_q = \{ z \in \operatorname{Mat}(q \times q, \mathbb{C}) \mid z = x + iy, \ x \in H_q, \ y \in \overline{\Omega}_q, \ \det z \neq 0 \}.$$

Its interior is the classical tube domain over the cone Ω_q , namely

$$T_q = \{ z \in \operatorname{Mat}(q \times q, \mathbb{C}) \mid z = x + iy, \ y \in \Omega_q \}.$$

Let $G = GL(q, \mathbb{C})$ act on $Mat(q \times q, \mathbb{C})$ by

$$(17) (g,z) \longmapsto gzg^*.$$

The spaces $H_q, \Omega_q, \overline{\Omega}_q$ are stable under this action, and hence \widetilde{T}_q and T_q are invariant subsets under this action. We investigate the orbits and describe a full set of invariants for this action.

There is a natural invariant associated to a $GL(q, \mathbb{C})$ -orbit. To any $z \in \widetilde{T}_q$, we associate its angular matrix defined by

(18)
$$a = a(z) = z^{*^{-1}}z.$$

Then the matrix associated to gzg^* is $g^{*^{-1}}ag^*$, so that the angular matrix a(z) belongs to the same conjugacy class when z runs through a $GL(q, \mathbb{C})$ -orbit. As we shall see (Theorem 3.3 and Theorem 3.13), this invariant is close to characterizing the orbits.

Let us first prove some elementary properties of the angular matrix.

PROPOSITION 3.1. Let $z = x + iy \in \widetilde{T}_q$, and let $a = z^{*^{-1}}z$ be its angular matrix. Then

- (i) $Sp(a) \subset U_1 = \{ \mu \in \mathbb{C}, |\mu| = 1 \};$
- (ii) if $1 \in Sp(a)$, then y is degenerate and

$$\{v \in \mathbf{C}^q \mid av = v\} = \{v \in \mathbf{C}^q \mid yv = 0\}.$$

Proof. Let μ be an eigenvalue of a, and let $v \neq 0$ be an eigenvector for the eigenvalue μ . Then $zv = \mu z^*v$, and hence

$$(zv, v) = \mu(z^*v, v) = \mu(v, zv) = \mu \overline{(zv, v)}.$$

If $(zv, v) \neq 0$, then $|\mu| = 1$. So we now assume (zv, v) = 0. This amounts to (xv, v) + i(yv, v) = 0, so that in particular (yv, v) = 0. Now recall that y is positive semi-definite. So the condition (yv, v) = 0 implies that yv = 0. From this it follows that $zv = xv = z^*v$, and as z is assumed to be invertible, this implies $\mu = 1$. This shows (i) and part of (ii). Conversely, the condition yv = 0 implies trivially av = v.

In particular, we may consider the polynomial $d(\mu) = \det(z - \mu z^*)$. The roots of d are the eigenvalues of the angular matrix. The set of these roots, counted with their multiplicities, will be called the *angular spectrum* of z.

We first consider the case of T_q . So let $z = x + iy \in T_q$. Then as y is positive-definite, we may define its square root $y^{1/2}$ as the unique positive-definite Hermitian matrix whose square is y. Then we may write

$$x + iy = y^{\frac{1}{2}}(y^{-\frac{1}{2}}xy^{-\frac{1}{2}} + i1_q)y^{\frac{1}{2}}.$$

This shows that any $GL(q, \mathbf{C})$ -orbit contains some element of the form $x+i1_q$, where $x \in H_q$. But by the classical diagonalization theorem for Hermitian forms, there exists an orthonormal basis in which the Hermitian form associated to x is diagonal. In other words, there exists a unitary matrix u and real numbers $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_q$ such that

$$uxu^* = \Lambda = egin{pmatrix} \lambda_1 & & & & \ & \lambda_2 & & & \ & & \ddots & & \ & & & \lambda_q \end{pmatrix}.$$

Moreover, if Λ and Λ' are two such diagonal matrices, then $\Lambda+i\mathbf{1}_q$ and $\Lambda'+i\mathbf{1}_q$ are not conjugate under the action of $\mathrm{GL}(q,\mathbf{C})$ unless $\Lambda=\Lambda'$. Hence we have shown the following result, which of course is the well-known fact that there is a simultaneous diagonalization for two Hermitian forms if one of them is positive-definite.

THEOREM 3.2. The set of matrices of the form

(19)
$$\Lambda = \begin{pmatrix} \lambda_1 + i & & \\ & \lambda_2 + i & \\ & & \ddots & \\ & & & \lambda_q + i \end{pmatrix}$$

with $\lambda_1 \geq \lambda_2 \cdots \geq \lambda_q$ is a full set of representatives of the $GL(q, \mathbb{C})$ -orbits in T_q .

The angular matrix associated to Λ is

(20)
$$\begin{pmatrix} \frac{\lambda_1+i}{\lambda_1-i} & & & \\ & \frac{\lambda_2+i}{\lambda_2-i} & & & \\ & & \ddots & & \\ & & & \frac{\lambda_q+i}{\lambda_q-i} \end{pmatrix}.$$

The latter is a semi-simple matrix with spectral values

$$\mu_j = \frac{\lambda_j + i}{\lambda_j - i}$$

for $1 \le j \le q$. Observe that these spectral values are complex numbers of modulus 1, but always different from 1. From the u_j we may recover the λ_j by the formula

$$\lambda_j = i \, \frac{1 + \mu_j}{1 - \mu_j} \, .$$

From these observations we get the following result.

THEOREM 3.3. Two elements z and z' of T_q belong to the same $GL(q, \mathbb{C})$ -orbit if and only if their angular matrices are conjugate. The angular spectrum is a full set of invariants for the action of $GL(q, \mathbb{C})$ on T_q .

The situation for \widetilde{T}_q is more complicated. In fact we may consider the extreme case where y=0. Then x corresponds to a non-degenerate Hermitian form, and the orbit picture is given by the signature. So we need to consider matrices of the form

$$\Upsilon = \Upsilon_{n_+,n_-} = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & -1 & \\ & & & \ddots & \\ & & & & -1 \end{pmatrix}$$

with n_+ diagonal entries equal to +1 and n_- diagonal entries equal to -1, n_+ and n_- being arbitrary nonnegative integers such that $n_+ + n_- = q$. The corresponding angular matrix is the identity matrix $\mathbf{1}_q$.

Another source of difficulty comes from the fact that it is not always possible to find a basis in which both Hermitian forms associated to x and y are diagonal. For instance if q = 2, consider the matrix

$$z = \begin{pmatrix} i & \frac{i}{2} \\ -\frac{i}{2} & 0 \end{pmatrix} = \begin{pmatrix} 0 & \frac{i}{2} \\ -\frac{i}{2} & 0 \end{pmatrix} + i \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}.$$

Notice that its angular matrix is

$$a = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix},$$

which is not semisimple.

From these examples we see that neither the angular spectrum of z nor the conjugacy class of the angular matrix characterizes the orbit of z.

Let n_1, n_2, n_3, n_4 be four nonnegative integers such that $n_1+2n_2+n_3+n_4=q$, and let $\lambda_1, \lambda_2, \ldots, \lambda_{n_1}$ be n_1 real numbers satisfying the condition

$$\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_{n_1}$$
.

To such data we associate the matrix $\Lambda = \Lambda(\lambda_1, \lambda_2, \dots, \lambda_{n_1}, n_2, n_3, n_4)$ given by

where there are n_2 diagonal 2×2 submatrices of the form $\begin{pmatrix} i & 1 \\ 1 & 0 \end{pmatrix}$, n_3 diagonal terms equal to 1 and n_4 diagonal terms equal to -1.

THEOREM 3.4. Any $GL(q, \mathbb{C})$ orbit in \widetilde{T}_q contains one and only one matrix of the form $\Lambda(\lambda_1, \lambda_2, \dots, \lambda_{n_1}, n_2, n_3, n_4)$.

Before beginning the proof, let us prove a couple of lemmas. Lemmas 3.6 and 3.7 are related to the classical Gauss's algorithm for diagonalizing an Hermitian form. Let r, s, n be three nonnegative integers such that r + s = n.

LEMMA 3.5. The stabilizer in $GL(n, \mathbb{C})$ of the matrix $y_r = \begin{pmatrix} \mathbf{1}_r & \\ & \mathbf{0}_s \end{pmatrix}$ is the subgroup

$$(22) G_r = \left\{ \begin{pmatrix} u & v \\ 0 & h \end{pmatrix} \right\}$$

where $u \in U(r)$, $v \in Mat(r, s)$, $h \in GL(s, \mathbb{C})$.

Proof. Easy computation.

Now we study the action of G_r in H_n . If $x \in H_n$, let us write

$$x = \begin{pmatrix} \alpha & b \\ b^* & \gamma \end{pmatrix}$$

where $\alpha \in H_r, b \in \operatorname{Mat}(r \times s, \mathbb{C})$ and $\gamma \in H_s$. If $g = \begin{pmatrix} u & v \\ 0 & h \end{pmatrix} \in G_r$, then $gxg^* = \begin{pmatrix} \alpha' & b' \\ b'^* & \gamma' \end{pmatrix}$, with

$$\alpha' = u\alpha u^* + ubv^* + vb^*u^* + v\gamma v^*$$
$$b' = ubh^* + v\gamma h^*$$
$$\gamma' = h\gamma h^*.$$

LEMMA 3.6. Let $x = \begin{pmatrix} \alpha & b \\ b^* & \gamma \end{pmatrix} \in H_n$, with $\alpha \in H_r$, $b \in \operatorname{Mat}(r \times s, \mathbb{C})$ and $\gamma \in H_s$. Assume $\det \gamma \neq 0$. Then the orbit of x under G_r contains a matrix of the form $\begin{pmatrix} \alpha' & 0 \\ 0 & \gamma \end{pmatrix}$ with $\alpha' \in H_r$.

Proof. This is a consequence of the previous formula with $u = \mathbf{1}_r$, $v = -b\gamma^{-1}$ and $h = \mathbf{1}_s$.

LEMMA 3.7. Let $x = \begin{pmatrix} \alpha & b \\ b^* & 0 \end{pmatrix} \in H_n$, with rank b = s (so in particular $r \geq s$). Then the orbit of x under G_r contains an element of the form

$$\begin{pmatrix} \beta & 0 & 0 \\ 0 & 0 & \mathbf{1}_s \\ 0 & \mathbf{1}_s & 0 \end{pmatrix}$$

with $\beta \in H_{r-s}$.

Proof. Consider the subgroup $\left\{ \begin{pmatrix} u & 0 \\ 0 & h \end{pmatrix}, u \in U(r), h \in GL_s(\mathbf{C}) \right\}$. It acts on the component b by $b' = ubh^*$. As $\operatorname{rank}(b) = s$, we may think of b as a set of s independent vectors in \mathbf{C}^r . By the Gram-Schmidt process, it is possible to find $h \in \operatorname{GL}_s(\mathbf{C})$ such that bh^* is a s-orthonormal frame in \mathbf{C}^r . But now two such frames are conjugate by the (left) action of U(r). Hence there exists $u \in U(r)$ such that

$$ubh^* = \begin{pmatrix} 0 \\ \mathbf{1}_s \end{pmatrix}.$$

The matrix x we started with is conjugate under G_r to a matrix of the form

$$\begin{pmatrix} \alpha' & c & 0 \\ c^* & \beta & \mathbf{1}_s \\ 0 & \mathbf{1}_s & 0 \end{pmatrix}$$

where $\alpha' \in H_{r-s}$, $\beta \in H_s$ and $c \in \text{Mat}((r-s) \times s, \mathbb{C})$. Now we use the action of the element

$$g = \begin{pmatrix} \mathbf{1}_{r-s} & 0 & -c \\ 0 & \mathbf{1}_s & -\frac{\beta}{2} \\ 0 & 0 & \mathbf{1}_s \end{pmatrix} \in G_r$$

to get the result.

We are now ready to start the proof of Theorem 3.4.

STEP 1. Let $z = x + iy \in \widetilde{T}_q$. As y is positive semidefinite, there exists an element $g \in GL(q, \mathbb{C})$ such that

$$gyg^* = \begin{pmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & 0 & & \\ & & & \ddots & \\ & & & 0 \end{pmatrix},$$

with r diagonal entries equal to 1, and s diagonal entries equal to 0, r and s being nonnegative integers satisfying r+s=q. In other terms, any $\mathrm{GL}(q,\mathbf{C})$ -orbit in \widetilde{T}_q contains an element of the form

$$\begin{pmatrix} \alpha + i \mathbf{1}_r & b \\ b^* & \gamma \end{pmatrix}$$

with $\alpha \in H_r, \gamma \in H_s, b \in \operatorname{Mat}(r \times s, \mathbb{C})$.

STEP 2. Now assume x is of the form

$$x = \begin{pmatrix} \alpha + i\mathbf{1}_r & b \\ b^* & \gamma \end{pmatrix}.$$

Consider γ . It is an Hermitian matrix of size s, and under the action of $GL(s, \mathbb{C})$ it can be transformed to

$$\begin{pmatrix} \mathbf{0}_{n_2} & 0 & 0 \\ 0 & \mathbf{1}_{n_3} & 0 \\ 0 & 0 & -\mathbf{1}_{n_4} \end{pmatrix}$$

where $n_2 + n_3 + n_4 = s$. Hence x is conjugate under the action of G_r to an element of the form

$$\begin{pmatrix} \alpha & b' & c' \\ b'^* & 0 & 0 \\ c'^* & 0 & \Upsilon \end{pmatrix}$$

where $\alpha \in H_r$, $b' \in \text{Mat}(r \times n_2, \mathbb{C})$, $c' \in \text{Mat}(r \times (n_3 + n_4), \mathbb{C})$ and

$$\Upsilon = \left(egin{array}{cc} \mathbf{1}_{n_3} & 0 \ 0 & -\mathbf{1}_{n_4} \end{array}
ight).$$

Using Lemma 3.6, we see that x is conjugate under the action of G_s to an element of the form

$$\begin{pmatrix} \alpha'' & b'' & 0 \\ b''^* & 0 & 0 \\ 0 & 0 & \Upsilon \end{pmatrix},$$

with $\alpha'' \in H_r$, $b'' \in \operatorname{Mat}(r \times n_2, \mathbb{C})$.

STEP 3. Assume now that

$$x = \begin{pmatrix} \alpha & b & 0 \\ b^* & 0 & 0 \\ 0 & 0 & \Upsilon \end{pmatrix}$$

with $\alpha \in H_r$ and $b \in \text{Mat}(r \times n_2, \mathbb{C})$. Recall that

$$x + iy = \begin{pmatrix} \alpha + i\mathbf{1}_r & b & 0 \\ b^* & 0 & 0 \\ 0 & 0 & \Upsilon \end{pmatrix}$$

is assumed to be invertible. This shows that $rank(b) = n_2$. So we may apply Lemma 3.7 to see that x is conjugate under G_r to an element of the form

$$egin{pmatrix} eta & 0 & 0 & 0 \ 0 & 0 & \mathbf{1}_{n_2} & 0 \ 0 & \mathbf{1}_{n_2} & 0 & 0 \ 0 & 0 & 0 & \Upsilon \end{pmatrix}$$

with $\beta \in H_{r-n_2}$.

STEP 4. Set $n_1 = r - n_2$. The last step is just to put the element $\beta \in H_{n_1}$ in diagonal form under the action of $U(n_1)$. Up to minor rearrangements of the matrix, this shows that any $GL(q, \mathbb{C})$ -orbit in \widetilde{T}_q contains an element of the form $\Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4)$.

STEP 5. It remains to show that two Λ 's are not conjugate under $GL(q, \mathbb{C})$. The angular matrix associated to $\Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4)$ is

$$\begin{pmatrix} \frac{\lambda_{1}+i}{\lambda_{1}-i} & & & & & & & \\ & \ddots & & & & & & \\ & \frac{\lambda_{n_{1}}+i}{\lambda_{n_{1}}-i} & & & & & & \\ & & 1 & 0 & & & & \\ & & 2i & 1 & & & & \\ & & & & \ddots & & & \\ & & & & 1 & 0 & & \\ & & & & 2i & 1 & & \\ & & & & & \ddots & & \\ & & & & & & 1 \end{pmatrix}$$

where there are n_2 2 × 2 submatrices $\binom{1}{2i}$, and $n_3 + n_4$ diagonal elements equal to 1. From the Jordan normal form theorem, we deduce that if $\Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4)$ and $\Lambda(\lambda'_1, \ldots, \lambda'_{n_1}, n'_2, n'_3, n'_4)$ are in a same $GL(q, \mathbb{C})$ -orbit, then $n_1 = n'_1$, $\lambda_j = \lambda'_j$ for all $j, 1 \leq j \leq n_1$, $n_2 = n'_2$ and $n_3 + n_4 = n'_3 + n'_4$. Now the matrix $\Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4) = L + iM$ and $\Lambda' = L' + iM'$, with $L, L', M, M' \in H_n$. As Λ and Λ' are supposed to be in the same $GL(q, \mathbb{C})$ -orbit, L and L' are also in the same $GL(q, \mathbb{C})$ -orbit, and so they must have the same signature. This forces $n_3 = n'_3$ and $n_4 = n'_4$, and hence $\Lambda = \Lambda'$.

We can now give the solution to the orbit problem we addressed at the end of Section 2. Recall that for any integer r such that $0 \le r \le q$ we defined

$$\widetilde{T}_q^{(r)} = \{ z = x + iy \mid y \in \overline{\Omega}_q, \text{ rank}(y) \le r, z \text{ invertible} \}.$$

LEMMA 3.8. Let n_1, n_2, n_3, n_4 be four integers such that

$$n_1 + 2n_2 + n_3 + n_4 = q,$$

and let $\lambda_1, \ldots, \lambda_{n_1}$ be n_1 real numbers. Then the standard matrix $\Lambda = \Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4)$ belongs to $\widetilde{T}_q^{(r)}$ if and only if $n_1 + n_2 \leq r$.

In fact the rank of $\frac{1}{2i}(\Lambda - \Lambda^*)$ is $n_1 + n_2$.

THEOREM 3.9. Any $GL(q, \mathbf{C})$ -orbit in $\widetilde{T}_q^{(r)}$ contains a unique standard matrix $\Lambda((\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4))$ with $n_1 + n_2 \leq r$.

We now want an analog of Theorem 3.3. As we have already noticed, the conjugacy class of the angular matrix does not determine the orbit of the matrix. We need a finer invariant, which we will construct now.

LEMMA 3.10. The space \widetilde{T}_q is connected and simply connected.

Proof. As T_q is connected and $T_q \subset \widetilde{T}_q \subset \overline{T}_q$, the space \widetilde{T}_q is connected. Take $i\mathbf{1}_q$ as base point in \widetilde{T}_q , and observe that for any $z \in \widetilde{T}_q$ and any s > 0, $z + is\mathbf{1}_q$ is in T_q . So if $(\gamma(t), t \in [0,1])$ is a path in \widetilde{T}_q starting and ending at $i\mathbf{1}_q$ then we can deform it by homotopy to $\gamma_s(t) = \gamma(t) + is(s-1)\mathbf{1}_q$, which for s > 0 is a path inside T_q . But T_q as a tube-type domain is simply connected. \square

The function $z\mapsto \det(z)$ is a continuous function from \widetilde{T}_q into ${\bf C}^*$. From Lemma 3.10, there exists a unique continuous determination of the argument of $\det(z)$ denoted by $\arg\det:\widetilde{T}_q\longrightarrow {\bf R}$ such that $\arg\det i{\bf 1}_q=q\frac{\pi}{2}$. If $Y\in\Omega_q$, then $\arg\det iy=q\frac{\pi}{2}$. If $z\in\widetilde{T}_q$ and $g\in {\rm GL}(q,{\bf C})$, then $\det gzg^*=|\det g|^2\det z$, and $gi{\bf 1}_qg^*=igg^*\in i\Omega_q$, so that

$$\arg \det gzg^* = \arg \det z$$
.

This provides a new invariant for the action of $GL(q, \mathbb{C})$ on \widetilde{T}_q .

LEMMA 3.11. Let
$$\Lambda = \Lambda(\lambda_1, \ldots, \lambda_{n_1}, n_2, n_3, n_4)$$
. Then

(23)
$$\arg \det \Lambda = \arg(\lambda_1 + i) + \cdots + \arg(\lambda_{n_1} + i) + n_2 \pi + \dot{n_4} \pi$$

where arg is used for the principal determination of the argument of a non-zero complex number.

Proof. We need to describe a continuous path from $i\mathbf{1}_q$ to Λ inside T_q . For clarity of exposition, we describe successively the path for each diagonal block (either a one-dimensional or a two-dimensional submatrix) of Λ , and compute the contribution of each block to the function arg det.

For a block of the form $\lambda + i$, with $\lambda \in \mathbf{R}$ we use the path $t \mapsto t\lambda + i$, $0 \le t \le 1$, and so the contribution of this block is $\arg(\lambda + i)$.

For a block of the form $\begin{pmatrix} i & 1 \\ 1 & 0 \end{pmatrix}$, we use the path

$$t \mapsto \begin{pmatrix} i & t \\ t & i(1-t^2) \end{pmatrix}, \ 0 \le t \le 1.$$

The corresponding determinant of this 2×2 -block is constant along the path and equal to -1. Hence the contribution of this block is $2\frac{\pi}{2} = \pi$.

For a block of the form 1, we use the path $t \mapsto e^{i\frac{\pi}{2}(1-t)}$, $0 \le t \le 1$, and we see that the corresponding contribution is 0.

For a block of the form -1, we use the path $t \mapsto e^{i\frac{\pi}{2}(1+t)}$, $0 \le t \le 1$, and we see that the corresponding contribution is π .

Putting together the contribution of the blocks, we get the result. \Box

COROLLARY 3.12. Let Λ and Λ' be two standard matrices. Assume that their angular matrices coincide and that $\operatorname{arg} \det \Lambda = \operatorname{arg} \det \Lambda'$. Then $\Lambda = \Lambda'$.

Proof. In fact we noticed that the equality of angular matrices implies the equality of the parameters except for $n_3 = n'_3$ and $n_4 = n'_4$. But from (23), we see that the equality of the determination of the arguments of the determinants implies $n_4 = n'_4$ (and hence $n_3 = n'_3$).

Now we can state the conclusion of this section, which is a consequence of Theorem 3.4 and Corollary 3.12.

THEOREM 3.13. Let $z, z' \in \widetilde{T}_q$, and assume that the angular matrices of z and z' are conjugate, and that $\arg \det z = \arg \det z'$. Then z and z' belong to the same orbit under the action of $\mathrm{GL}(q, \mathbf{C})$.

REMARK. Let
$$z \in \widetilde{T}_q$$
. Let $a = z^{*^{-1}}z$. Then

$$\det a = \frac{\det z}{\det z} = \left| \det z \right|^{-2} (\det z)^{2}.$$

So 2 arg det z is a determination of $arg(\det a)$. If z and z' are two matrices in \widetilde{T}_q with the same angular matrix, then $arg \det z$ and $arg \det z'$ differ by an integral multiple of π . So the new invariant needed to characterize the orbits under $GL(q, \mathbb{C})$ has to be regarded as a \mathbb{Z} -valued function. In this sense, it is a generalization of the signature.