

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** TRIPLE RATIO ON THE UNITARY STIEFEL MANIFOLD  
**Autor:** Clerc, Jean-Louis

**Bibliographie**  
**DOI:** <https://doi.org/10.5169/seals-66067>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 04.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

is the complex number

$$T(\sigma_1, \sigma_2, \sigma_3) = (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1}.$$

The group  $GL(q, \mathbf{C}) \simeq \mathbf{C}^*$  acts on the upper halfplane by  $(\lambda, z) \mapsto |\lambda|^2 z$  and so the orbits are described by the argument of the complex number  $z$ . So the characteristic invariant in this case is just

$$\arg \left( (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1} \right).$$

It is equivalent to the invariant  $\theta$  considered in [KR]. This invariant, almost in our terms, was known to E. Cartan (see [Ca]).

REMARK 2. Let us consider the case where  $p = q$ . Then the Stiefel manifold is  $U(q)$ , and the content of Proposition 4.2 is that for  $(\sigma_1, \sigma_2, \sigma_3) \in S_{\top}^3$

$$T(\sigma_1, \sigma_2, \sigma_3) = (1 - \sigma_2^* \sigma_1)^{-1} (1 - \sigma_2^* \sigma_3) (1 - \sigma_1^* \sigma_3)^{-1}$$

is an invertible skew-Hermitian matrix. The orbits of  $GL(q, \mathbf{C})$  in its action on nondegenerate Hermitian forms are characterized by the signature. So the characteristic invariant as described in Theorem 4.3 in this case reduces to  $\text{sgn } iT(\sigma_1, \sigma_2, \sigma_3)$ . As concerns Theorem 4.4, notice that the invariant  $S$  is trivial (equal to  $-\mathbf{1}_q$ ), so one is only concerned with the invariant  $\arg \det T$ . The bounded domain  $D$  is of tube type and the description of the invariant through the function  $\arg \det$  coincides with the approach of this problem in [CØ], where the invariant was introduced under the name of *generalized Maslov index*.

## REFERENCES

- [Ca] CARTAN, E. Sur le groupe de la géométrie hypersphérique. *Comment. Math. Helv.* 4 (1932), 158–171.
- [CØ] CLERC, J.-L. and B. ØRSTED. The Maslov index revisited. *Transform. Groups* 6 (2001), 303–320.
- [FK] FARAUT, J. and A. KORÁNYI. *Analysis on Symmetric Cones*. Oxford Mathematical Monographs, Clarendon Press, Oxford, 1994.
- [Fal] FARAUT, J. et al. *Analysis and Geometry on Complex Homogeneous Domains*. Progress in Mathematics 185. Birkhäuser Verlag, Boston, 2000.
- [H] HUA, L.-K. Geometries of matrices, I. Generalizations of von Staudt's theorem. *Trans. Amer. Math. Soc.* 57 (1945), 441–481.
- [KR] KORÁNYI, A. and H. M. REIMANN. The complex cross ratio on the Heisenberg group. *L'Enseign. Math.* (2) 33 (1987), 291–300.

- [P] PIATETSKIĬ-SHAPIRO, L.I. *Geometry of Classical Domains and Theory of Automorphic Forms*. Gordon and Breach, New York, 1969.
- [S] SATAKE, I. *Algebraic Structures of Symmetric Domains*. Kanô Memorial Lectures (4). Iwanami Shoten and Princeton University Press, 1980.

(Reçu le 29 mars 2001)

Jean-Louis Clerc

Institut Élie Cartan  
Université Henri Poincaré  
B.P. 239  
F-54506 Vandœuvre-lès-Nancy Cedex  
France  
*e-mail*: [clerc@iecn.u-nancy.fr](mailto:clerc@iecn.u-nancy.fr)

**Vide-leer-empty**