

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE HILBERT METRIC AND GROMOV HYPERBOLICITY
Autor: Karlsson, Anders / NOSKOV, Guennadi A.
Kapitel: Introduction
DOI: <https://doi.org/10.5169/seals-66068>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE HILBERT METRIC AND GROMOV HYPERBOLICITY

by Anders KARLSSON¹⁾ and Guennadi A. NOSKOV²⁾

ABSTRACT. We give some sufficient conditions for Hilbert's metric on convex domains D to be Gromov hyperbolic. The conditions involve an intersecting chords property, which we in turn relate to the Menger curvature of triples of boundary points and, in the case the boundary is smooth, to differential geometric curvature of ∂D . In particular, the intersecting chords property and hence Gromov hyperbolicity is established for bounded, convex C^2 -domains in \mathbf{R}^n with non-zero curvature.

We also give some necessary conditions for hyperbolicity: the boundary must be of class C^1 and may not contain a line segment. Furthermore we prove a statement about the asymptotic geometry of the Hilbert metric on arbitrary convex (i.e. not necessarily strictly convex) bounded domains, with an application to maps which do not increase Hilbert distance.

INTRODUCTION

Let D be a bounded convex domain in \mathbf{R}^n and let h be the Hilbert metric, which is defined as follows. For any distinct points $x, y \in D$, let x' and y' be the intersections of the line through x and y with ∂D closest to x and y respectively. Then

$$h(x, y) = \log \frac{yx' \cdot xy'}{xx' \cdot yy'}$$

where zw denotes the Euclidean distance $\|z - w\|$ between two points. The expression $\frac{yx' \cdot xy'}{xx' \cdot yy'}$ is called the *cross-ratio* of four collinear points and is invariant under projective transformations. For the basic properties of the distance h we refer to [Bu55] or [dIH93].

¹⁾ Supported by SFB 343 of the Universität Bielefeld.

²⁾ Supported by SFB 343 of the Universität Bielefeld and GIF-grant G-454-213.06/95.

We will give here some sufficient conditions for the metric space (D, h) to be hyperbolic in the sense of Gromov. Namely, we show that a certain intersecting chords property implies Gromov hyperbolicity (Theorem 2.1). This intersecting chords property holds when the (Menger) curvature of any three points of the domain's boundary is uniformly bounded from both above and below in a certain way (Corollary 1.2). Domains with C^2 boundary of everywhere nonzero curvature satisfy this condition as will be proved in Section 3. Beardon showed in [Be97] (see also [Be99]) that a weaker intersecting chords property holds for any bounded strictly convex domain and he used this to establish some weak hyperbolicity results for the Hilbert metric. In Section 5 we prove a generalization of his results to any bounded convex domain.

It would also be interesting to understand what conditions on ∂D are necessary in order for (D, h) to be Gromov hyperbolic. For example, in Section 4 we give an argument showing that ∂D must be of class C^1 and that it may not contain a line segment.

Some parts of the results in this paper might already be known: Y. Benoist has told us that a convex domain with C^2 boundary is Gromov hyperbolic if the curvature of the boundary is everywhere nonzero. Benoist has also found examples of Gromov hyperbolic Hilbert geometries whose boundaries are C^1 but not C^2 . In [Be99] it is mentioned that C. Bell has proved an intersecting chords theorem in an unpublished work. However, we have not found the present arguments or our main results in the literature. Furthermore, we have not found the simple and attractive Proposition 1.1 and Corollary 3.5 stated or discussed anywhere, although these facts are most likely known. They should belong to ancient Greek geometry and classical differential geometry respectively.

Since the Hilbert distance can be defined in analogy with Kobayashi's pseudo-distance on complex spaces [Ko84], we would like to mention that Balogh and Bonk proved in [BB00] that the Kobayashi metric on any bounded strictly pseudoconvex domain with C^2 boundary is Gromov hyperbolic.

Note that metric spaces of this type are CAT(0) only in exceptional cases (see [BH99] for the definition). Indeed, Kelly and Straus proved in [KS58] that if (D, h) is nonpositively curved in the sense of Busemann then D is an ellipsoid and hence (D, h) is the n -dimensional hyperbolic space. Compare this to the situation for Banach spaces: a Banach space is CAT(0) if and only if it is a Hilbert space. Another category of results is of the following type: if D has a large (infinite, cocompact, etc.) automorphism group and C^2 -smooth boundary, then it is an ellipsoid; see the work of Socié-Méthou [SM00].

The Hilbert metric has found several applications, see [Bi57], [Li95] and [Me95] just to mention a few instances. Typically the idea is to apply the contraction mapping principle to maps which do not increase Hilbert distances (e.g. affine maps).

This work was mainly done during our stay at Bielefeld University. We are grateful to this university for its hospitality and for providing such excellent working conditions. Especially we wish to thank Professor H. Abels for inviting us there.

1. INTERSECTING CHORDS IN CONVEX DOMAINS

From elementary school we know that if c_1, c_2 are two intersecting chords in a circle, then $l_1 l'_1 = l_2 l'_2$ where l_1, l'_1 and l_2, l'_2 denote the respective lengths of the segments into which the two chords are divided. (This follows immediately from the similarity of the associated triangles, see Fig. 1.)




FIGURE 1

Intersecting chords in a circle

A generalization of this fact to any bounded strictly convex domain was given by Beardon in [Be97] by an elegant argument using the Hilbert metric. He proved that if D is such a domain then for each positive δ there is a positive number $M = M(D, \delta)$ such that for any intersecting chords c_1, c_2 , each of length at least δ , one has

$$(1.1) \quad M^{-1} \leq \frac{l_1 l'_1}{l_2 l'_2} \leq M,$$

where l_1, l'_1 , and l_2, l'_2 denote the respective lengths of the segments into which the two chords are divided.