Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE HILBERT METRIC AND GROMOV HYPERBOLICITY
Autor: Karlsson, Anders / NOSKOV, Guennadi A.
Kapitel: 1.1 Intersecting line segments and Menger curvature
DOI: https://doi.org/10.5169/seals-66068

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 12.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

We say that a domain satisfies the *intersecting chords property* (ICP) if (1.1) holds for *any* two intersecting chords c_1 and c_2. It is easy to see that ICP may fail for a general strictly convex domain (at a curvature zero point or a 'corner').

We show in this section that ICP holds for domains that satisfy a certain (non-differentiable) curvature condition. Domains with C^2 boundary of nonvanishing curvature are proved to satisfy this condition in Section 3.

1.1 Intersecting line segments and Menger curvature

This subsection clarifies the relation between the curvature of any triple of endpoints and the ratio considered above that two intersecting line segments define.

Three distinct points A, B and C in the plane, not all on a line, lie on a unique circle. Recall that the radius of this circle is

$$R(A,B,C) = \frac{c}{2\sin\gamma}, \tag{1.2}$$

where c is the length of a side of the triangle ABC and γ is the opposite angle. The reciprocal of R is called the *(Menger) curvature* of these three points and is denoted by $K(A,B,C)$.

Now consider two intersecting line segments as in Fig. 2.



FIGURE 2
Intersecting line segments

PROPOSITION 1.1. *In the above notation, the following equality holds:*

$$\frac{a_1a_2}{b_1b_2} = \frac{K(A_1,B_1,B_2)K(A_2,B_1,B_2)}{K(B_1,A_1,A_2)K(B_2,A_1,A_2)}.$$

Proof. Let α_i be the angle between the line segments A_iB_j and B_1B_2, and let β_i be the angle between B_iA_j and A_1A_2, for $\{i,j\} = \{1,2\}$. By the sine law we have

$$\frac{a_1a_2}{b_1b_2} = \frac{\sin\alpha_1 \sin\alpha_2}{\sin\beta_2 \sin\beta_1} = \frac{2\sin\alpha_1}{|A_1B_1|}\frac{|A_2B_2|}{2\sin\beta_2}\frac{2\sin\alpha_2}{|A_2B_2|}\frac{|A_1B_1|}{2\sin\beta_1}$$
$$= \frac{K(A_1,B_1,B_2)K(A_2,B_1,B_2)}{K(B_1,A_1,A_2)K(B_2,A_1,A_2)}. \qquad \square$$

COROLLARY 1.2. *Let D be a bounded convex domain in $\mathbf{R}^n$. Assume that there is a constant $C > 0$ such that*

$$\frac{K(x,y,z)}{K(x',y',z')} \le C$$

for any two triples of distinct points in ∂D all lying in the same 2-dimensional plane. Then D satisfies the intersecting chords property.

Proof. Any two intersecting chords define a plane and by Proposition 1.1 we have

$$\frac{a_1a_2}{b_1b_2} = \frac{K_{\alpha 1}K_{\alpha 2}}{K_{\beta 1}K_{\beta 2}} \le C^2. \qquad \square$$

REMARK 1.3. In view of this subsection it is clear that ICP implies restrictions on the curvature of the boundary, e.g. there cannot be any points of zero curvature. We were however not able to establish the converse of Corollary 1.2.

1.2 CHORDS LARGER THAN δ

The following proposition provides a different approach to the result in [Be97] mentioned above.

PROPOSITION 1.4. *Let D be a bounded convex domain in $\mathbf{R}^n$. Let δ be such that the length of any line segment contained in ∂D is bounded from above by some $\delta' < \delta$. Then there is a constant $C = C(D,\delta) > 0$ such that*

$$C(D,\delta) \le K(x,y,z) \le \frac{2}{\delta}, \tag{1.3}$$

whenever $x,y,z \in \partial D$ and $xy \ge \delta$.

Proof. The angle $\alpha(x,y,v) := \angle_y(xy,v)$ is continuous in $x,y \in \mathbf{R}^n$ and $v \in UT_y(\partial D)$, the unit tangent cone at y. The tangent cone at a boundary point y is the union of all hyperplanes containing y but which are disjoint from D. If $[x,y]$ does not lie in ∂D, then $0 < \alpha(x,y,v) < \pi$. The set

$$S = \{(x,y,v) \in \partial D \times \partial D \times UT_y(\partial D) \,:\, xy \ge \delta\}$$