

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE HILBERT METRIC AND GROMOV HYPERBOLICITY
Autor: Karlsson, Anders / NOSKOV, Guennadi A.
Kapitel: 1.2 Chords larger than
DOI: <https://doi.org/10.5169/seals-66068>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$$\begin{aligned} \frac{a_1 a_2}{b_1 b_2} &= \frac{\sin \alpha_1 \sin \alpha_2}{\sin \beta_2 \sin \beta_1} = \frac{2 \sin \alpha_1 |A_2 B_2|}{|A_1 B_1|} \frac{2 \sin \alpha_2 |A_1 B_1|}{2 \sin \beta_2 |A_2 B_2|} \frac{2 \sin \beta_1}{2 \sin \beta_1} \\ &= \frac{K(A_1, B_1, B_2) K(A_2, B_1, B_2)}{K(B_1, A_1, A_2) K(B_2, A_1, A_2)}. \quad \square \end{aligned}$$

COROLLARY 1.2. *Let D be a bounded convex domain in \mathbf{R}^n . Assume that there is a constant $C > 0$ such that*

$$\frac{K(x, y, z)}{K(x', y', z')} \leq C$$

for any two triples of distinct points in ∂D all lying in the same 2-dimensional plane. Then D satisfies the intersecting chords property.

Proof. Any two intersecting chords define a plane and by Proposition 1.1 we have

$$\frac{a_1 a_2}{b_1 b_2} = \frac{K_{\alpha_1} K_{\alpha_2}}{K_{\beta_1} K_{\beta_2}} \leq C^2. \quad \square$$

REMARK 1.3. In view of this subsection it is clear that ICP implies restrictions on the curvature of the boundary, e.g. there cannot be any points of zero curvature. We were however not able to establish the converse of Corollary 1.2.

1.2 CHORDS LARGER THAN δ

The following proposition provides a different approach to the result in [Be97] mentioned above.

PROPOSITION 1.4. *Let D be a bounded convex domain in \mathbf{R}^n . Let δ be such that the length of any line segment contained in ∂D is bounded from above by some $\delta' < \delta$. Then there is a constant $C = C(D, \delta) > 0$ such that*

$$(1.3) \quad C(D, \delta) \leq K(x, y, z) \leq \frac{2}{\delta},$$

whenever $x, y, z \in \partial D$ and $xy \geq \delta$.

Proof. The angle $\alpha(x, y, v) := \angle_y(xy, v)$ is continuous in $x, y \in \mathbf{R}^n$ and $v \in UT_y(\partial D)$, the unit tangent cone at y . The tangent cone at a boundary point y is the union of all hyperplanes containing y but which are disjoint from D . If $[x, y]$ does not lie in ∂D , then $0 < \alpha(x, y, v) < \pi$. The set

$$S = \{(x, y, v) \in \partial D \times \partial D \times UT_y(\partial D) : xy \geq \delta\}$$

is compact. Hence there is a constant $\alpha_0 > 0$ such that

$$(1.4) \quad \alpha_0 \leq \alpha(x, y, v) \leq \pi - \alpha_0$$

for every $(x, y, v) \in S$. By the definition of the tangent cone and compactness there is an $\varepsilon > 0$ such that for any $y, z \in \partial D$, $0 < yz < \varepsilon$ there is an element $v \in UT_y(\partial D)$ for which

$$(1.5) \quad 0 \leq \angle_y(yz, v) \leq \alpha_0/2.$$

The estimates (1.4) and (1.5) imply the existence of $C > 0$ and the other inequality in (1.3) is trivial. \square

As an immediate consequence of Propositions 1.1 and 1.4 we have:

COROLLARY 1.5 (cf. [Be99]). *Let D be a bounded convex domain such that any line segment in ∂D has length less than $\delta' < \delta$. Then the intersecting chords property holds for any two chords each of length greater than δ .*

2. HYPERBOLICITY OF HILBERT'S METRIC

Let (Y, d) be a metric space. Given two points $z, w \in Y$, let

$$(z | w)_y = \frac{1}{2}(d(z, y) + d(w, y) - d(z, w))$$

be their *Gromov product* relative to y . We think of y as a fixed base point. The metric space Y is *Gromov hyperbolic* (or δ -*hyperbolic*) if there is a constant $\delta \geq 0$ such that the inequality

$$(x | z)_y \geq \min\{(x | w)_y, (w | z)_y\} - \delta$$

holds for any four points x, y, z, w in Y . As is known, it is enough to show such an inequality for a fixed y (the δ changes by a factor of 2); see [BH99] for a proof of this and we also refer to this book for a general exposition of this important notion of hyperbolicity. By expanding the terms the above inequality is equivalent to

$$(2.1) \quad d(x, z) + d(y, w) \leq \max\{d(x, y) + d(z, w), d(y, z) + d(x, w)\} + 2\delta.$$