

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE HILBERT METRIC AND GROMOV HYPERBOLICITY  
**Autor:** Karlsson, Anders / NOSKOV, Guennadi A.  
**Kapitel:** 2. Hyperbolicity of Hilbert's metric  
**DOI:** <https://doi.org/10.5169/seals-66068>

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 06.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

is compact. Hence there is a constant  $\alpha_0 > 0$  such that

$$(1.4) \quad \alpha_0 \leq \alpha(x, y, v) \leq \pi - \alpha_0$$

for every  $(x, y, v) \in S$ . By the definition of the tangent cone and compactness there is an  $\varepsilon > 0$  such that for any  $y, z \in \partial D$ ,  $0 < yz < \varepsilon$  there is an element  $v \in UT_y(\partial D)$  for which

$$(1.5) \quad 0 \leq \angle_y(yz, v) \leq \alpha_0/2.$$

The estimates (1.4) and (1.5) imply the existence of  $C > 0$  and the other inequality in (1.3) is trivial.  $\square$

As an immediate consequence of Propositions 1.1 and 1.4 we have:

**COROLLARY 1.5** (cf. [Be99]). *Let  $D$  be a bounded convex domain such that any line segment in  $\partial D$  has length less than  $\delta' < \delta$ . Then the intersecting chords property holds for any two chords each of length greater than  $\delta$ .*

## 2. HYPERBOLICITY OF HILBERT'S METRIC

Let  $(Y, d)$  be a metric space. Given two points  $z, w \in Y$ , let

$$(z \mid w)_y = \frac{1}{2}(d(z, y) + d(w, y) - d(z, w))$$

be their *Gromov product* relative to  $y$ . We think of  $y$  as a fixed base point. The metric space  $Y$  is *Gromov hyperbolic* (or  $\delta$ -hyperbolic) if there is a constant  $\delta \geq 0$  such that the inequality

$$(x \mid z)_y \geq \min\{(x \mid w)_y, (w \mid z)_y\} - \delta$$

holds for any four points  $x, y, z, w$  in  $Y$ . As is known, it is enough to show such an inequality for a fixed  $y$  (the  $\delta$  changes by a factor of 2); see [BH99] for a proof of this and we also refer to this book for a general exposition of this important notion of hyperbolicity. By expanding the terms the above inequality is equivalent to

$$(2.1) \quad d(x, z) + d(y, w) \leq \max\{d(x, y) + d(z, w), d(y, z) + d(x, w)\} + 2\delta.$$

**THEOREM 2.1.** *Let  $D$  be a bounded convex domain in  $\mathbf{R}^n$  satisfying the intersection chords property. Then the metric space  $(D, h)$  is Gromov hyperbolic.*

*Proof.* Suppose that the intersecting chords property holds with a constant  $M$ . Let  $y$  be a fixed reference point and consider any other three points  $x, z, w$  in  $D$ . Set  $A(u, v) = h(u, v) + h(w, y) - h(u, w) - h(v, y)$  for any two points  $u, v$ . By (2.1) we need to show that there is a constant  $\delta$  independent of  $x, z, w$  such that

$$(2.2) \quad \min\{A(x, z), A(z, x)\} \leq 2\delta.$$




FIGURE 3

Four points

Using the definition of  $h$  and the notation in Fig. 3, we have (by rearranging the terms of the product)

$$\begin{aligned} A(x, z) &= \log \left( \frac{xz'' \cdot zx''}{xx'' \cdot zz''} \frac{wy'' \cdot yw''}{ww'' \cdot yy''} \frac{xx' \cdot ww'}{xw' \cdot wx'} \frac{zz' \cdot yy'}{zy' \cdot yz'} \right) \\ &= \log \left( \frac{xx' \cdot xz''}{xx'' \cdot xw'} \frac{yy' \cdot yw''}{yy'' \cdot yz'} \frac{zz' \cdot zx''}{zz'' \cdot zy'} \frac{ww' \cdot wy''}{ww'' \cdot wx'} \right). \end{aligned}$$

Hence, by using  $\frac{xx'}{xx''} \leq M \frac{xz''}{xw'}$  and similar inequalities for the other fractions,

$$A(x, z) \leq M' + 2 \log \left( \frac{xz''}{xw'} \frac{yw''}{yz'} \frac{zx''}{zy'} \frac{wy''}{wx'} \right).$$

Now,  $y$  is fixed and  $zy', wy''$  are bounded from above and below respectively, so that

$$A(x, z) \leq M'' + 2 \log \left( \frac{xz'' \cdot zx''}{xw' \cdot wx'} \right).$$

So (2.2) is equivalent to the boundedness of

$$\min \left\{ \frac{xz'' \cdot zx''}{xw' \cdot wx'}, \frac{zx'' \cdot xz''}{zw''' \cdot wz'''} \right\}$$

from above. By symmetry we may assume without loss of generality that  $zz'' \leq xx''$ . Now we have two cases:

CASE 1:  $xw \geq xx''$  or  $zw \geq xx''$ .

If  $xw \geq zw$  (so in particular  $xw \geq xx''$ ), then

$$\begin{aligned} \frac{xz'' \cdot zx''}{xw' \cdot wx'} &\leq \frac{(xz + zz'')(zx + xx'')}{(xw)^2} \\ &\leq \frac{(xw + wz + zz'')(zw + wx + xx'')}{(xw)^2} \leq \frac{(3xw)^2}{(xw)^2} \leq 9. \end{aligned}$$

When  $zw \geq xw$ , we estimate the other fraction instead (obtained by interchanging  $x$  and  $z$ ) in the same way.

CASE 2:  $xw \leq xx''$  and  $zw \leq xx''$ .

Considering chords at  $x$  we have

$$\frac{xz'' \cdot zx''}{xw' \cdot wx'} \leq M \frac{xx' \cdot zx''}{xx'' \cdot wx'} \leq M \frac{xx'}{wx'} \frac{(xw + wz + xx'')}{xx''} \leq 3M$$

since  $xx'' \cdot xz'' \leq M(xx' \cdot xw')$ .  $\square$

REMARK 2.2. Since the  $n$ -dimensional ball  $B^n$  obviously satisfies the assumption in Corollary 1.2 with  $C = 1$ , Theorem 2.1 contains the standard fact that  $(B^n, h)$ , which is Klein's model of the  $n$ -dimensional hyperbolic space, is Gromov hyperbolic.

REMARK 2.3. The above proof does not appeal to compactness and therefore goes through in infinite dimensions provided that  $y$  lies at positive distance from the boundary. In particular, it proves that the unit ball in a Hilbert space with the Hilbert metric, which is the infinite dimensional hyperbolic space, is Gromov hyperbolic. Note however that ICP is not affinely invariant in infinite dimensions. (Kaimanovich brought this remark to our attention.)