

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 48 (2002)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** THE HILBERT METRIC AND GROMOV HYPERBOLICITY  
**Autor:** Karlsson, Anders / NOSKOV, Guennadi A.  
**Kapitel:** 5. NON-STRICTLY CONVEX DOMAINS  
**DOI:** <https://doi.org/10.5169/seals-66068>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 07.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

neighbourhood  $U_\varepsilon$  of 0 such that  $f(x) < \varepsilon|x|$  in this neighbourhood. But in view of  $f_{x_i}(0) = 0$ ,  $i = 1, \dots, n$ , we have  $f(0, \dots, 0, x_i, 0, \dots, 0) < \varepsilon|x_i|$  for sufficiently small  $x_i$  and hence by convexity  $f(x) < \varepsilon|x|$  for sufficiently small  $|x|$ .  $\square$

REMARK 4.3. The following was announced in [B00]: *If a strictly convex domain  $D$  is divisible, that is, if it admits a proper cocompact group of isometries  $\Gamma$ , then  $D$  is Gromov hyperbolic if and only if  $\partial D$  is  $C^1$ .* Our Theorem 4.2 shows that in the implication (Gromov hyperbolicity + divisibility  $\Rightarrow C^1$ ) the condition of divisibility is superfluous.

## 5. NON-STRICTLY CONVEX DOMAINS

This section owes much of its existence to [Be97] and [Be99]. Using a different argument, we prove certain extensions to arbitrary convex bounded domains of some of the results obtained in those papers.

LEMMA 5.1. *Let  $D$  be a bounded convex domain in  $\mathbf{R}^n$ . Let  $\{x_n\}, \{y_n\}$  be two sequences of points in  $D$ . Assume that  $x_n \rightarrow \bar{x} \in \partial D$ ,  $y_n \rightarrow \bar{y} \in \bar{D}$  and  $[\bar{x}, \bar{y}] \not\subseteq \partial D$ . Let  $x'_n$  and  $y'_n$  denote the endpoints of the chord through  $x_n$  and  $y_n$  as usual. Then  $x'_n$  converges to  $\bar{x}$  and  $y'_n$  converges to the endpoint  $\bar{y}'$  of the chord defined by  $\bar{x}$  and  $\bar{y}$  different from  $\bar{x}$ .*

*Proof.* Compare with Lemma 5.3. in [Be97]. Every limit point of chord endpoints must belong to the line through  $\bar{x}$  and  $\bar{y}$ . In addition, in the case of  $x'_n$  for example, any limit point must lie on the halfline from  $\bar{x}$  not containing  $\bar{y}$ . At the same time each limit point must belong to the boundary of  $D$ , and the statement follows since the line through  $\bar{x}$  and  $\bar{y}$  intersects  $\partial D$  only in  $\bar{x}$  and  $\bar{y}'$ .  $\square$

THEOREM 5.2. *Let  $D$  be a bounded convex domain. Let  $\{x_n\}$  and  $\{z_n\}$  be two sequences of points in  $D$ . Assume that  $x_n \rightarrow \bar{x} \in \partial D$ ,  $z_n \rightarrow \bar{z} \in \partial D$  and  $[\bar{x}, \bar{z}] \not\subseteq \partial D$ . Then there is a constant  $K = K(\bar{x}, \bar{z})$  such that for the Gromov product  $(x_n | z_n)_y$  in Hilbert distances relative to some fixed point  $y$  in  $D$  we have*

$$\limsup_{n \rightarrow \infty} (x_n | z_n)_y \leq K.$$

*Proof.* By Lemma 5.1, the endpoints of the chords through  $x_n$  and  $z_n$  converge to  $\bar{x}$  and  $\bar{z}$ .




FIGURE 5  
Partial hyperbolicity

Since  $[\bar{x}, \bar{z}]$  is not contained in the boundary of  $D$ , there are small compact neighbourhoods  $\bar{U}_x$  and  $\bar{U}_z$  of  $\bar{x}$  and  $\bar{z}$  respectively, in  $\partial D$ , such that every chord with endpoints in  $\bar{U}_x$  and  $\bar{U}_z$  is contained in  $D$ . In particular the Euclidean midpoint of every such chord lies inside  $D$  and by compactness there is an upper bound  $K$  on  $h(y, w)$ , where  $w$  is the midpoint of such a chord.

Consider three points  $x, y, z$  and a point  $w$  on a (minimizing) geodesic segment  $[x, z]$  in a (geodesic) metric space  $(Y, d)$ . Then

$$\begin{aligned} (x \mid z)_y &= \frac{1}{2}(d(x, y) + d(z, y) - d(x, z)) \\ &= \frac{1}{2}(d(x, y) + d(z, y) - d(x, w) - d(w, z)) \\ &\leq \frac{1}{2}(d(y, w) + d(y, w)) = d(y, w) \end{aligned}$$

by the triangle inequality. It follows from this estimate and the above considerations that eventually

$$(x_n \mid z_n)_y \leq K. \quad \square$$

REMARK 5.3. The content of Theorem 5.2 is that  $(D, h)$  satisfies a weak notion of hyperbolicity. This property should be compared with Gromov hyperbolicity, especially with the fact that for Gromov hyperbolic spaces, two sequences converge to the same point of the boundary if and only if their Gromov product tends to infinity. Theorem 5.2 can be applied as in [Ka01, Theorem 8] to the study of random walks on the automorphism group of  $D$ , and it is also likely to be useful for analyzing commuting nonexpanding maps or isometries of  $(D, h)$ .

REMARK 5.4. We suggest that a similar statement might hold for the classical Teichmüller spaces and perhaps also for more general Kobayashi hyperbolic complex spaces. Hilbert geodesic rays from a point  $y$  that terminate on a line segment contained in the boundary may correspond to the Teichmüller geodesic rays defined by Jenkins-Strebel differentials that H. Masur considered when demonstrating the failure of  $\text{CAT}(0)$  for the Teichmüller space of Riemann surfaces of genus  $g \geq 2$ . The complement of the union of all line segments in the boundary  $\partial D$  may correspond to the uniquely ergodic foliation points on the Thurston boundary of Teichmüller space.

Using the arguments in [Ka01], see Proposition 5.1 of that paper, we obtain the following result as an application of Theorem 5.2:

THEOREM 5.5. *Let  $D$  be a bounded convex domain and  $\varphi: D \rightarrow D$  be a map which does not increase Hilbert distances. Then either the orbit  $\{\varphi^n(y)\}_{n=1}^\infty$  is bounded or there is a limit point  $\bar{y}$  of the orbit such that for any other limit point  $\bar{x}$  of the orbit it holds that  $[\bar{x}, \bar{y}] \subset \partial D$ .*

This theorem, which extends a theorem in [Be97], provides a general geometric explanation for a part of the main theorem in [Me01].

## REFERENCES

- [BB00] BALOGH, Z.M. and M. BONK. Gromov hyperbolicity and the Kobayashi metric on strictly pseudoconvex domains. *Comment. Math. Helv.* 75 (2000), 504–533.
- [Be97] BEARDON, A.F. The dynamics of contractions. *Ergodic Theory Dynam. Systems* 17(6) (1997), 1257–1266.
- [Be99] ——— The Klein, Hilbert and Poincaré metrics of a domain. *J. Comput. Appl. Math.*, 105(1–2) (1999), 155–162. Continued fractions and geometric function theory (CONFUN), Trondheim, 1997.
- [B00] BENOIST, Y. Convexes divisibles. *C.R. Acad. Sci. Paris Sér. I Math.* 322 (2001), no. 5, 387–390.
- [BG88] BERGER, M. and B. GOSTIAUX. *Differential Geometry: Manifolds, Curves, and Surfaces*. Translated from the French by Silvio Levy. Springer-Verlag, New York, 1988.
- [Bi57] BIRKHOFF, G. Extensions of Jentzsch's theorem. *Trans. Amer. Math. Soc.* 85 (1957), 219–227.
- [BH99] BRIDSON, M. and A. HAEFLIGER. *Metric Spaces of Non-Positive Curvature*. Grundlehren der Math. Wissenschaften, 319. Springer-Verlag (Berlin), 1999.