Zeitschrift:	L'Enseignement Mathématique				
Herausgeber:	Commission Internationale de l'Enseignement Mathématique				
Band:	48 (2002)				
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE				
Artikel:	SEMISTABLE K3-SURFACES WITH ICOSAHEDRAL SYMMETRY				
Autor:	Stevens, Jan				
Kapitel:	4. HODGE ALGEBRAS				
DOI:	https://doi.org/10.5169/seals-66069				

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

### Download PDF: 06.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3.15. A nice construction for studying the homology of the general fibre is given by [A'C]. Let  $\sigma_i: \mathcal{Z}_i \to \mathcal{X}$  be the oriented real blow-up of  $X_i \subset \mathcal{X}$ . This is a manifold with boundary, whose boundary  $\partial \mathcal{Z}_i = \sigma_i^{-1}(X_i)$  is isomorphic to the boundary of a tubular neighbourhood of  $X_i$  in  $\mathcal{X}$ . The fibred product  $\sigma: \mathcal{Z} \to \mathcal{X}$  of the  $\sigma_i$  is a manifold with corners. Its boundary  $\mathcal{N} := \partial \mathcal{Z}$  comes with a map to X. It also fibres over  $S^1$ : the composed map  $\mathcal{Z} \to \mathcal{X} \to S \ni 0$  extends to a map from  $\mathcal{Z}$  to the real oriented blow-up of S in 0 (polar coordinates !). A fibre of  $\mathcal{N} \to S^1$  is then a topological model of the general fibre.

This model is not sufficient to describe the monodromy. One has first to replace X by the geometric realisation of the simplicial object  $X^{[\cdot]}$ : one replaces each double point by an interval, and each triple point by a 2-simplex. A final fibred product then gives the new model. For details see [A'C, §2].

## 4. HODGE ALGEBRAS

4.1. STANLEY-REISNER RINGS. Let  $\Delta$  be a simplicial complex with set of vertices  $V = \{v_1, \ldots, v_n\}$ . A monomial on V is an element of  $\mathbb{N}^V$ . Each subset of V determines a monomial on V by its characteristic function. The support of a monomial  $M: V \to \mathbb{N}$  is the set supp $M = \{v \in V \mid M(v) \neq 0\}$ . The set  $\Sigma_{\Delta}$  of monomials whose support is not a face is an ideal, generated by the monomials corresponding to minimal non-simplices.

Given a ring R and an injection  $\phi: V \to R$  we can associate to each monomial M on V the element  $\phi(M) = \prod_{v \in V} \phi(v)^{M(v)} \in R$ . We will usually identify V and  $\phi(V)$  and write  $M \in R$  for  $\phi(M)$ . This applies in particular to the polynomial ring K[V] over a field K. The ideal  $\Sigma_{\Delta}$  gives rise to the Stanley-Reisner ideal  $I_{\Delta} \subset K[V]$ . The Stanley-Reisner ring is  $A_{\Delta} = K[V]/I_{\Delta}$ .

Deformations of Stanley-Reisner rings are studied in [A-C].

4.2. EXAMPLE. Let  $\Delta$  be an octahedron. We map the set of vertices to  $\mathbb{C}[x_1, \ldots, x_6]$  such that opposite vertices correspond to variables with index sum 7.

The Stanley-Reisner ring is minimally generated by the three monomials  $x_i x_{7-i}$ . The spaces smoothes to a K3-surface, the complete intersection of three general quadrics. A general 1-parameter deformation is not semi-stable, because the total space has singularities at the six quadruple points of the special fibre.



FIGURE 4.1 Octahedron

4.3. REMARK. To get an octahedron as dual graph we need the incidence relations of a cube. The toric variety associated to a cube is  $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1$ . The general anticanonical divisor is a smooth K3, whereas the complement of the torus is the unions of six quadrics (each of the form  $\mathbf{P}^1 \times \mathbf{P}^1$ ) with dual graph the octahedron. A small resolution of a general pencil yields a semistable degeneration.

4.4. DEFINITION [D-E-P]. Let H be a finite partially ordered set and  $\Sigma$  an ideal of monomials on H. Let K be a commutative ring and R a commutative K-algebra. Suppose an injection  $\phi: H \to R$  is given. Then R is a *Hodge algebra* (or algebra with straightening law) governed by  $\Sigma$  if H-1 R is a free K-module admitting the set of monomials not in  $\Sigma$  as basis,

H-2 for each generator M of  $\Sigma$  in the unique expression

(\*) 
$$M = \sum_{N \notin \Sigma} k_{M,N} N, \qquad k_{M,N} \in K,$$

guaranteed by H-1, for each  $x \in H$  dividing M and each  $N \notin \Sigma$  with  $k_{M,N} \neq 0$  there is a  $y_{M,N}$  dividing N and satisfying  $y_{M,N} < x$ .

The relations (\*) are called *straightening relations* for R.

4.5. If R is graded and the elements of  $\phi(H)$  are homogeneous the straightening relations give a presentation for R [D-E-P, p. 15].

We note that R is a deformation of the discrete Hodge algebra governed by  $\Sigma$ , whose ideal is generated by the monomials M.

4.6. EXAMPLE. The equations of the tetrahedron of degree 12 of 2.2 are straightening relations. We take  $\Sigma$  as Stanley-Reisner ideal  $\Sigma_{\Delta}$ , where  $\Delta$  is the stellation of the tetrahedron: in each top-dimensional face we take an additional vertex, which is joined to all vertices on the face. The partial order on the set of vertices is obtained by declaring the new vertices to be smaller. The discrete Hodge algebra then has equations  $x_i x_j$ ,  $x_i y_j$  and  $y_j y_k y_l$ .

# 5. THE DODECAHEDRON

5.1. To get an icosahedron as dual graph we need the incidence relations of a dodecahedron. Each side should be a rational surface and the intersection with the other surfaces should have a pentagon as dual graph. A pentagon occurs as hyperplane section of a Del Pezzo surface of degree 5. So we can realise our dodecahedron by gluing together 12 Del Pezzo surfaces.

We first describe the Del Pezzo surfaces. Each of those is an extension of its pentagonal hyperplane section. Its coordinate ring can be obtained as Stanley-Reisner ring of a pentagon as 1-dimensional simplicial complex. Introducing variables  $y_i$ , we get the equations  $y_{i-1}y_{i+1}$ . With an extra variable x the Del Pezzo surface has equations

$$y_{i-1}y_{i+1} - xy_i - x^2$$
.

These are the Pfaffians of the matrix

( 0	<i>y</i> <sub>1</sub>	X	-x	$-y_5$	
$-y_{1}$	0	<i>y</i> <sub>2</sub>	x	-x	
-x	$-y_{2}$	0	<i>y</i> <sub>3</sub>	x	
x	-x	$-y_{3}$	0	У4	
<b>y</b> 5	X	-x	$-y_{4}$	0 /	

We can check that this is indeed a smooth Del Pezzo of degree 5 by giving an explicit birational map from  $\mathbf{P}^2$ , which blows up four points, see Figure 5.1.

To the variable x corresponds a new vertex at the centre of the pentagon. By joining it to all other vertices we obtain a 2-dimensional simplicial complex, and the homogeneous coordinate ring of the Del Pezzo surface is a graded Hodge algebra governed by the Stanley-Reisner ideal of the complex: to satisfy H-2 we take x to be less than all  $y_i$ .