

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 48 (2002)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE FANNING METHOD FOR CONSTRUCTING EVEN UNIMODULAR LATTICES. I
Autor: ROEGNER, Katherine
Kapitel: Introduction
DOI: <https://doi.org/10.5169/seals-66072>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

THE FANNING METHOD FOR CONSTRUCTING EVEN UNIMODULAR LATTICES. I

by Katherine ROEGNER

ABSTRACT. This paper provides a formal study of isofans and discusses their use in the theory of even unimodular lattices. Examples are given that illustrate how isofans simplify the construction of certain types of even unimodular lattices. A classification of isofans concludes the paper.

INTRODUCTION

The history of even unimodular lattices dates back to the 19th century when H. J. S. Smith [Sm] showed the existence of what is known today as the E_8 lattice. The even unimodular lattices have been classified for dimensions 8 [M], 16 [W2], and 24 [N]. The next dimension of interest is 32 due to the fact that even unimodular lattices only occur in dimensions divisible by 8; see e.g. [Sch]. In dimension 32, there are millions of nonisometric even unimodular lattices. Although no classification in this dimension is available, there has been considerable progress. Conway and Pless [CP] determined the doubly-even self-dual binary codes, the results of which can be transformed into a classification statement for even unimodular lattices with complete root systems of a particular type. Within their work, they noted that it is possible to build some codes using known codes by making appropriate substitutions. Kervaire [Ke] classified the remaining cases of complete even unimodular lattices in dimension 32 using a lengthy elimination procedure and a lot of machine testing. Venkov [V] has shown that, except for 15 cases, the even unimodular lattices in dimension 32 can be generated by the roots and vectors with scalar square 4. In that article, Venkov introduced an important operation on lattices, which he called “fanning”. It turns out that Venkov’s fanning method is comparable to Conway and Pless’ substitution method.

The purpose of this article is to provide an indepth study of the fanning method. To do so, Venkov's fanning method is generalized to the isofan, a special isomorphism between rational bilinear form modules associated to root lattices. Some examples are given illustrating the construction of new complete even unimodular lattices from already known ones using isofans. In particular, an easy construction for a lattice that Conway and Pless found using "several processes including divination" is given. A classification of isofans concludes the paper.

The author is indebted to Helmut Koch for the hints and suggestions he has provided. Special thanks are due to Boris Venkov for the many helpful discussions concerning even unimodular lattices and to the referee for suggesting many improvements to the original version of this paper.

1. LATTICES

Let \mathbf{R}^n be n -dimensional euclidean space equipped with the standard scalar product

$$x \cdot y = \sum_{i=1}^n x_i y_i \text{ for all } x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbf{R}^n .$$

A free \mathbf{Z} -module $\Lambda \subset \mathbf{R}^n$ of rank $k := \dim_{\mathbf{R}} \mathbf{R} \otimes_{\mathbf{Z}} \Lambda$ is called a *lattice of rank k* . A *basis* of a rank k lattice Λ is a subset $\{\lambda_1, \dots, \lambda_k\} \subset \Lambda$ that generates Λ over \mathbf{Z} .

Let $\Lambda \subset \mathbf{R}^n$ be a lattice. Λ is said to be *integral* if $\lambda_i \cdot \lambda_j \in \mathbf{Z}$ for all $\lambda_i, \lambda_j \in \Lambda$. It is an *even* lattice if, in addition to being integral, $\lambda^2 := \lambda \cdot \lambda \in 2\mathbf{Z}$ for all $\lambda \in \Lambda$. Let $\Lambda^\# = \{x \in \mathbf{R}^n \mid x \cdot \lambda \in \mathbf{Z} \text{ for all } \lambda \in \Lambda\}$ denote the *dual lattice*. Clearly, Λ is integral if and only if $\Lambda \subseteq \Lambda^\#$. Λ is called *unimodular* if in fact $\Lambda = \Lambda^\#$. Thus, an even unimodular lattice is a self-dual lattice such that $\lambda^2 \in 2\mathbf{Z}$ for all $\lambda \in \Lambda$.

Let $\Lambda_1, \dots, \Lambda_m$ be nontrivial sublattices of the integral lattice Λ whose direct sum is equal to Λ . If $x \cdot y = 0$ for all $x \in \Lambda_i, y \in \Lambda_j, i \neq j$, then Λ is called the orthogonal direct sum of the sublattices $\Lambda_1, \dots, \Lambda_m$ and denoted by $\Lambda = \Lambda_1 \oplus \dots \oplus \Lambda_m$. Λ is called *decomposable* if there exists such an orthogonal direct sum with $m > 1$, otherwise Λ is said to be *indecomposable*.

The *root system* of an even lattice Λ is the set

$$\Lambda_{\text{rt}} := \{\lambda \in \Lambda \mid \lambda^2 = 2\},$$

the elements of which are called *roots*. Λ is called a *root lattice* if Λ is