

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE ENTROPY OF HOLOMORPHIC MAPS
Autor: GROMOV, Mikhaïl
Kapitel: §1. Notation and definitions
DOI: <https://doi.org/10.5169/seals-66687>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

me to write this paper. I also appreciate the hospitality of IHES, which made possible my involvement in this story. I am especially thankful to Dennis Sullivan, who took pains to read the paper and to clean it up of multiple errors.

STRUCTURE OF THE PAPER. We start with a geometric outlook on topological entropy and reduce our inequality (0.1) to standard facts about minimal varieties. We discuss next the real algebraic analogue of (0.1) and a generalization to maps. We conclude with an estimate of the entropy involving the mean curvature.

§ 1. NOTATION AND DEFINITIONS

For a space X we denote by X^k the product $X \times X \times \dots \times X$ (k factors). A *graph* Γ over X is by definition an arbitrary set $\Gamma \subset X^2$. When X is finite this is the usual definition of an oriented graph (with loops). The graph of a map $X \rightarrow X$ gives another example.

For a graph Γ we denote by $\Gamma_k \subset X^k$ the set of strings $(x_1, \dots, x_i, \dots, x_k)$, $x_i \in X$, where each pair $(x_{i-1}, x_i) \in X^2$ is contained in Γ .

When X is endowed with a metric, we call ϵ -*cubes* products in X^k of balls from X of radius ϵ . For a set $Y \subset X^k$ we denote by $\text{Cap}_\epsilon Y$ the minimal number of ϵ -cubes needed to cover Y .

ENTROPY

Set $h_\epsilon(\Gamma) = \limsup_{k \rightarrow \infty} \frac{1}{k} \log \text{Cap}_\epsilon \Gamma_k$, and $h(\Gamma) = \lim_{\epsilon \rightarrow 0} h_\epsilon(\Gamma)$, for $\Gamma \subset X^2$.

When f is an endomorphism $X \rightarrow X$, we define its *entropy* $h(f)$ as the entropy of its graph Γ_f . If the space X is compact, the definition does not depend on the choice of the metric [2]. Observe that the entropy of a general graph Γ is equal to the entropy of the shift in $\Gamma_\infty \subset X^\infty$: Γ_∞ is the space of doubly infinite strings $(x_i)_{i=\dots,-1,0,1,\dots}$ with the product topology, and the shift maps (x_i) to (x_{i+1}) . For finite X , we come to the usual definition of the Markov shift.

VOLUME

From now on, X is a Riemannian manifold and $n = \dim \Gamma$, $\Gamma \subset X^2$. We denote by $\text{Vol } \Gamma_k$ the n -dimensional volume of $\Gamma_k \subset X^k$, i.e. the

n -dimensional Hausdorff measure with respect to the Riemann product metric in X^k . Set

$$\text{lov } \Gamma = \limsup_{k \rightarrow \infty} \frac{1}{k} \log \text{Vol } \Gamma_k.$$

For an f we set $\text{lov } f = \text{lov } \Gamma_f$. This is a smooth invariant of f (it does not depend on the choice of the Riemann metric).

Our invariant “lov” is sometimes more accessible than entropy and for a holomorphic f we are going to prove that

$$(1.0) \quad h(f) \leq \text{lov } f.$$

DENSITY

Denote by $\text{Dens}_\epsilon(\Gamma_k, \gamma)$, for $\gamma \in \Gamma_k \subset X^k$, the n -dimensional measure of the intersection of Γ_k with the ball (in the Riemannian product metric) of radius ϵ centered at γ . Set $\text{Dens}_\epsilon(\Gamma_k) = \inf_{\gamma \in \Gamma_k} \text{Dens}_\epsilon(\Gamma_k, \gamma)$, and then $\text{lodn}_\epsilon \Gamma = \liminf_{k \rightarrow \infty} \frac{1}{k} \log \text{Dens}_\epsilon \Gamma_k$, and finally

$$\text{lodn } \Gamma = \lim_{\epsilon \rightarrow 0} \text{lodn}_\epsilon \Gamma.$$

Observe that $\text{Vol} \geq \text{Cap}_{2\epsilon} \text{Dens}_\epsilon$ and hence that

$$(1.1) \quad h(V) \leq \text{lov } \Gamma - \text{lodn } \Gamma.$$

§2. ESTIMATES OF DENSITY

Consider a Riemannian manifold W (it will be X^k in the sequel) and an n -dimensional subvariety $V \subset W$. We suppose that the boundary of V (if there is such) does not intersect the ball $B_\epsilon \subset W$ of radius $\epsilon > 0$ centered at a point $v_0 \in V$. We suppose also that the injectivity radius of W at v_0 is at least ϵ , i.e. the exponential map $T_{v_0}(W) \rightarrow W$ is injective in the ϵ -ball in $T_{v_0}(W)$.

DENSITY OF A MINIMAL VARIETY

If the sectional curvature in B_ϵ is not greater than K and V is minimal, then

$$(2.0) \quad \text{Vol}(V \cap B_\epsilon) \geq C > 0,$$

where the constant C depends on n , K , and ϵ , but does not depend on $\dim W$.