

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES
Autor: Gautero, François
Kurzfassung
DOI: <https://doi.org/10.5169/seals-66690>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES

by François GAUTERO

ABSTRACT. This paper deals with the geometry of metric ‘two-dimensional’ spaces, equipped with semi-flows admitting transverse foliations by forests. Our main theorem relates the Gromov-hyperbolicity of such spaces, for instance mapping-telescopes of \mathbf{R} -trees, with the dynamical behaviour of the semi-flow. As a corollary, we give a new proof of the following theorem [3]: *Let α be a hyperbolic injective endomorphism of the rank n free group F_n . If the image of α is a malnormal subgroup of F_n , then $G_\alpha = F_n \rtimes_\alpha \mathbf{Z}$ is a hyperbolic group.*

INTRODUCTION

The subject of 3-dimensional topology changed completely in the seventies with Thurston’s geometric methods. His geometrization conjecture involves eight classes of manifolds, among which the hyperbolic manifolds play the most important role. In this context, a hyperbolic manifold is a compact manifold which admits (or whose interior admits in the case of non-empty boundary) a metric of constant curvature -1 . According to another conjecture of Thurston, any closed hyperbolic 3-manifold should have a finite cover which is a mapping-torus. This gives a particular interest to these mapping-tori manifolds. Recall that a mapping-torus is a manifold which fibers over the circle. Namely this is a 3-manifold constructed from a homeomorphism h of a compact surface Σ as

$$M = (\Sigma \times [0, 1]) / ((x, 1) \sim (h(x), 0)).$$

For these manifolds, the hyperbolization conjecture has been proved, see for instance [25]: the manifold M constructed from Σ and h as above is hyperbolic if and only if Σ has negative Euler characteristic and h is a pseudo-Anosov homeomorphism (see [12]).