

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES
Autor: Gautero, François
Kapitel: 2. Mapping-telescopes and forest-stacks
DOI: <https://doi.org/10.5169/seals-66690>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REMARK 1.1. The above computations fail, and the space is no longer Gromov-hyperbolic, if one replaces $d_y = \lambda^{|y|}d_0$ by $d_y = P(|y|)d_0$, where $P(\cdot)$ is a polynomial function of y . Indeed, in this case, the length of the horizontal interval between the two considered orbits, evaluated at the height where the minimum of the length-function $f(t)$ is attained, depends, even in the optimal case, on the horizontal length of the interval connecting one point to the orbit of the other. Whereas in the exponential case it equals $\frac{2}{\ln \lambda}$ unless it belongs to the horizontal axis.

2. MAPPING-TELESCOPES AND FOREST-STACKS

Let X be a topological space. Call X a *topological tree* if there exists a unique arc between any two points in X . A *topological forest* is a union of disjoint topological trees. By ‘arc’ we mean the image of an injective path. A path in X is a continuous map from a bounded interval of the real line into X . A *forest-map* is a continuous map of a topological forest into itself.

DEFINITION 2.1. Let $\psi: X \rightarrow X$ be a forest-map. The *mapping-telescope* K_ψ of (ψ, X) is the topological space resulting from $K_X = \bigsqcup_{n \in \mathbf{Z}} X \times [n, n + 1]$ by the identification of each point $(x, n + 1) \in X \times [n, n + 1]$ with the point $(\psi(x), n + 1) \in X \times [n + 1, n + 2]$.

Let us examine somewhat more closely the topology of these mapping-telescopes.

For any integer $n \in \mathbf{Z}$, for any $(x, r) \in X \times [n, n + 1]$, for any real number $t \geq 0$, we define $\tilde{\sigma}_t((x, r))$ as the point $(\psi^{E[t-(n+1-r)]+1}(x), r + t)$ in $X \times [E[r + t], E[r + t] + 1]$, where $E[r]$ denotes the integer part of r . The map $\tilde{\sigma}_t$ is defined on K_X (the disjoint union of the $X \times [n, n + 1]$) for every $t \geq 0$. Moreover $\tilde{\sigma}_{t+t'} = \tilde{\sigma}_t \circ \tilde{\sigma}_{t'}$.

If $a = (x, n + 1) \in X \times [n + 1, n + 2]$, then $\tilde{\sigma}_t(a) = (\psi^{E[t]}(x), n + 1 + t) \in [n + 1 + E[t], E[t] + n + 2]$. Whereas if $a = (x, n + 1) \in X \times [n, n + 1]$ then $\tilde{\sigma}_t(a) = (\psi^{E[t]+1}(x), n + 1 + t) \in X \times [n + 1 + E[t], E[t] + n + 2]$, which is equal to $\tilde{\sigma}_t(b)$ with $b = (\psi(x), n + 1) \in X \times [n + 1, n + 2]$. Therefore $(\tilde{\sigma}_t)_{t \in \mathbf{R}^+}$ descends to the mapping-telescope K_ψ , where it defines a one parameter family $(\sigma_t)_{t \in \mathbf{R}^+}$ of continuous maps of K_ψ . This family depends continuously on the parameter $t \in \mathbf{R}^+$. It satisfies furthermore $\sigma_0 = \text{Id}_{K_\psi}$ and $\sigma_{t+t'} = \sigma_t \circ \sigma_{t'}$. Such a family is called a *semi-flow* on K_ψ .

Let $f: K_\psi \rightarrow \mathbf{R}$ be defined by $f(a) = r$ if $a \in X \times \{r\}$. Then f is a continuous surjective map. The preimage of any real number r is $X \times \{r\}$, a topological forest. Furthermore, for any $t \geq 0$, $f \circ \sigma_t = \tau_t \circ f$, where $\tau_t: \mathbf{R} \rightarrow \mathbf{R}$ is defined by $\tau_t(r) = r + t$.

We extracted above the two properties shared by mapping-telescopes which are really important for our work. We now define a class of spaces which satisfy these two properties, and in particular generalize the mapping-telescopes.

DEFINITION 2.2. Let X be a topological space. Let $(\sigma_t)_{t \in \mathbf{R}^+}$ be a semi-flow on X . Let $f: X \rightarrow \mathbf{R}$ be a surjective continuous map such that:

1. For any real number r , the *stratum* $f^{-1}(r)$ is a topological forest.
2. For any $t \geq 0$, $f \circ \sigma_t = \tau_t \circ f$, where $\tau_t(r) = r + t$ for any real number r .

Then X is a *forest-stack*, denoted by (X, f, σ_t) .

REMARK 2.3. All the strata of a mapping-telescope are homeomorphic. This is not required in the definition of a forest-stack.

As we just saw, a mapping-telescope is an example of a forest-stack. In Section 13, we show that a Cayley complex for the mapping-torus group of an injective free group endomorphism is a mapping-telescope of a forest-map, and thus a forest-stack. The reader can also find there, and in Section 12, an illustration of the horizontal and vertical metrics on forest-stacks, which we are now going to define.

3. METRICS

The aim of this section is to introduce a particular metric on forest-stacks, called the *telescopic metric*. We sometimes deal with metric spaces which are not necessarily connected, for instance forests. In this case, when considering the distance between two points, it will always be tacitly assumed that the two points lie in a same connected component of the space.

3.1 HORIZONTAL AND VERTICAL METRICS

Let us consider a forest-stack (\tilde{X}, f, σ_t) , see Definition 2.2. We want to define a natural metric on the orbits of the semi-flow.