Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES
Autor: Gautero, François
Kapitel: 6. About straight quasi geodesics
DOI: https://doi.org/10.5169/seals-66690

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

6. About straight quasi geodesics

DEFINITION 6.1. Let $(\widetilde{X}, f, \sigma_t, \mathcal{H})$ be a forest-stack. A (J, J')-*quasi geodesic*, $J \geq 1$, $J' \geq 0$, in $(\widetilde{X}, d_{(\widetilde{X},\mathcal{H})})$ is a telescopic path S of which each subpath S' satisfies the inequality

$$|S'|_{(\widetilde{X},\mathcal{H})} \leq J d_{(\widetilde{X},\mathcal{H})}(i(S'), t(S')) + J' .$$

LEMMA 6.2. *Let p be a straight (J, J')-quasi geodesic with*

$$|r_{max} - f(i(p))| \leq t_0 ,$$

where $r_{max} = max_{x \in p} f(x)$. There exists a constant $C_{6.2}(J, J') \geq M$, which increases with J and J', such that if $|[p]_{r_{max}}|_{r_{max}} \geq C_{6.2}(J, J')$ then $[p]_{r_{max}}$ is dilated both in the future and in the past after $C_{6.2}(J, J')t_0$.

Proof. By the bounded-dilatation property, $|p|_{(\widetilde{X},\mathcal{H})} \geq \lambda_+^{-t_0} |[p]_{r_{max}}|_{r_{max}} + t_0$. We choose n_* so that $\lambda_+^{-t_0} - J\lambda^{-n_* t_0} > 0$. For any n greater than n_*, the inequality

$$J(2t_0 + 2nt_0 + \lambda^{-nt_0} |[p]_{r_{max}}|_{r_{max}}) + J' < \lambda_+^{-t_0} |[p]_{r_{max}}|_{r_{max}} + t_0$$

is satisfied for $|[p]_{r_{max}}|_{r_{max}} > \frac{(2J-1)t_0 + 2nJt_0 + J'}{\lambda_+^{-t_0} - J\lambda^{-nt_0}}$. This is in contradiction with p being a (J, J')-quasi geodesic. If $|[p]_{r_{max}}|_{r_{max}} > \lambda_+^{n_* t_0} M$, then, by the bounded-dilatation property, the geodesic preimages of $[p]_{r_{max}}$ under $\sigma_{n_* t_0}$ have horizontal length at least M. Hence, if moreover $|[p]_{r_{max}}|_{r_{max}} > \frac{(2J-1)t_0 + 2n_* Jt_0 + J'}{\lambda_+^{-t_0} - J\lambda^{-n_* t_0}}$ then the hyperbolicity of the semi-flow implies that they are dilated in the past after t_0. The bounded-dilatation property implies that these geodesic preimages have horizontal length at least $\lambda_+^{-n_* t_0} |[p]_{r_{max}}|_{r_{max}}$. Choosing N_* such that $\lambda^{N_* t_0} \geq \lambda_+^{n_* t_0}$, we conclude that $[p]_{r_{max}}$ is dilated in the past after $(N_* + 1)t_0$. The same arguments allow us to find a lower bound on $|[p]_{r_{max}}|_{r_{max}}$ for $[p]_{r_{max}}$ to be dilated in the future after some fixed finite time. $\square$

DEFINITION 6.3. Let $(\widetilde{X}, f, \sigma_t)$ be a forest-stack. A *stair* in $\widetilde{X}$ is a telescopic path along which the function f is monotone.

LEMMA 6.4. *Let p be a straight (J, J')-quasi geodesic stair between two points a and b, $f(a) \leq f(b)$. There exists a constant $C_{6.4}(J, J') \geq M$, which increases with J and J', such that if the horizontal length of a horizontal geodesic I between a and $O^-(b)$ (resp. b and $O^+(a)$) is at least $C_{6.4}(J, J')$, then I is dilated in the past (resp. in the future) after t_0.*

Proof. Let X be such that $\lambda^{t_0}X > X + \lambda_+^{t_0}C_{6.2}(J,J')$. Assume that the horizontal length of some horizontal geodesic I between a and $O^-(b)$ is at least X. By Lemma 6.2, the choice of X implies that if I is dilated in the future after t_0, then the first point a_1 along p satisfying $f(a_1) = f(a) + t_0$ is at horizontal distance greater than X from $O^-(b)$. By induction, we thus obtain an infinite sequence of points $a_1, a_2, \ldots, a_n, \ldots$ in p such that $f(a_i) = f(a_{i-1}) + t_0$ and each a_i is at horizontal distance at least X from $O^-(b)$. This is absurd. The other case of Lemma 6.4 is treated similarly. □

DEFINITION 6.5. Let S_0, S_1 be two telescopic paths whose pulled-tight projections agree after some finite time. We say that S_0 and S_1 *are in fine position* if, for any two points x, y, $x \neq y$, satisfying $x \in S_i \cap O(y)$, $y \in S_{i+1}$, $i = 0, 1 \mod 2$, then $x \in O^+(y) \cup O^-(y)$.

Let us observe that a path is always in fine position with respect to any of its pulled-tight projections.

DEFINITION 6.6. A $+$*-hole* (resp. $-$*-hole*) is a telescopic path with both endpoints in a same stratum, which is in fine position with respect to the horizontal geodesic I between its endpoints, and which satisfies furthermore $\min_{x\in p} f(x) \geq f(I)$ (resp. $\max_{x\in p} f(x) \leq f(I)$).

LEMMA 6.7. *Let p be a straight (J,J')-quasi geodesic $+$-hole (resp. $-$-hole). There exists a constant $C_{6.7}(J,J') \geq M$, which increases with J and J', such that, if I is the horizontal geodesic between the endpoints of p and if $|I|_{f(I)} \geq C_{6.7}(J,J')$, then I is dilated in the past (resp. future) after $C_{6.7}(J,J')t_0$.*

Proof. We consider a decomposition $p_1 p_2 \ldots p_l$ of p such that

$$\max_{x\in p_i} |f(x) - f(i(p_i))| \leq t_0 ,$$

and a decomposition $I_1 \ldots I_l$ of I, where I_k joins the past orbits of the endpoints of p_k. We denote by I_D the union of the I_k's which are dilated in the past after $C_{6.2}(J,J')t_0$, and by I_C the union of the other intervals in I. By Lemma 6.2, the horizontal length of any interval in I_C is at most $C_{6.2}(J,J')$.

Let n be some positive integer. We consider a horizontal geodesic h with $I = [h]_{f(h)+nC_{6.2}(J,J')t_0}$ and assume that h is dilated in the future after t_0. Then,

$$\lambda^n |I_D|_{f(I)} + \lambda_+^{-n} |I_C|_{f(I)} \leq |h|_{f(h)} \leq \lambda^{-n}(|I_D|_{f(I)} + |I_C|_{f(I)}) .$$

Hence $|I_C|_{f(I)} \geq \frac{\lambda^n-\lambda_+^{-n}}{\lambda^n-\lambda^{-n}}|I_D|_{f(I)}$, so that $|I_C|_{f(I)} \geq \frac{X(n)}{1+X(n)}|I|_{f(I)}$ with $X(n) = \frac{\lambda^n-\lambda_+^{-n}}{\lambda^n-\lambda^{-n}}$. Now $\lim_{n\to+\infty}\frac{X(n)}{1+X(n)} = 1$, so that for some $n_* \geq 1$, for any $n \geq n_*$, $\frac{X(n)}{1+X(n)} \geq \frac{1}{2}$. Since the horizontal length of any interval I_k in I_C is at most $C_{6.2}(J,J')$, and the telescopic length of the associated $p_k \subset p$ is at least t_0, we obtain

$$|p|_{(\widetilde{X},\mathcal{H})} \geq \frac{t_0}{2C_{6.2}(J,J')}|I|_{f(I)}.$$

On the other hand, $|p|_{(\widetilde{X},\mathcal{H})} \leq 2Jnt_0 + \lambda^{-n}J|I|_{f(I)} + J'$ for any $n \geq n_*$. The last two inequalities give, for $n \geq n_*$, $2Jnt_0 + \lambda^{-n}J|I|_{f(I)} + J' \geq \frac{t_0}{2C_{6.2}(J,J')}|I|_{f(I)}$, equivalently $2Jnt_0 + J' \geq (\frac{t_0}{2C_{6.2}(J,J')} - \lambda^{-n}J)|I|_{f(I)}$. We choose $n_\circ \geq n_*$ such that $\frac{t_0}{2C_{6.2}(J,J')} - \lambda^{-n_\circ}J > 0$. We get

$$\frac{2Jn_\circ t_0 + J'}{\frac{t_0}{2C_{6.2}(J,J')} - \lambda^{-n_\circ}J} \geq |I|_{f(I)}.$$

Thus, for $|I|_{f(I)} > \frac{2Jn_\circ t_0+J'}{\frac{t_0}{2C_{6.2}(J,J')}-\lambda^{-n_\circ}J}$, h is not dilated in the future after t_0. If $|I|_{f(I)} > \lambda_+^{n_\circ}M$, then $|h|_{f(h)} \geq M$. Therefore h is dilated in the past after t_0. We choose N such that $\lambda^N\lambda_+^{-n_\circ} > \lambda$. Thus, if $|I|_{f(I)} \geq \max(\lambda_+^{n_\circ}M, \frac{2Jn_\circ t_0+J'}{\frac{t_0}{2C_{6.2}(J,J')}-\lambda^{-n_\circ}J})$ then I is dilated in the past after $(n_\circ C_{6.2}(J,J') + N)t_0$. The arguments and computations in the case where $\max_{x\in p} f(x) \leq f(I)$ are the same. □

7. Substitution of quasi geodesics

Lemma 7.1. *Let p be a (J,J')-quasi geodesic. Let q be obtained from p by replacing subpaths $p_i \subset p$ by (L,L')-quasi geodesics q_i satisfying the following properties:*

- *q_i has the same endpoints as p_i,*
- *q_i is L-close to p_i,*
- *$|q_i|_{(\widetilde{X},\mathcal{H})} \leq L|p_i|_{(\widetilde{X},\mathcal{H})}$.*

There exists a constant $C_{7.1}(L,L',J,J')$, which increases in each variable, such that q is a $(C_{7.1}(L,L',J,J'), C_{7.1}(L,L',J,J'))$-quasi geodesic which is L-close to p.

Proof. Since each q_i is L-close to a p_i, and with the same endpoints, q is L-close to p. Let us consider any two points x, y in q and let $q_{xy} \subset q$