Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 49 (2003)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES

Autor: Gautero, François

Kapitel: 10. Straight quasi geodesic bigons are thin

DOI: https://doi.org/10.5169/seals-66690

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

10. STRAIGHT QUASI GEODESIC BIGONS ARE THIN

PROPOSITION 10.1. There exists a constant Bi(J, J') such that any straight (J, J')-quasi geodesic bigon is Bi(J, J')-thin.

Proof. We denote by g, g' the two sides of a (J, J')-quasi geodesic bigon. We assume for a while that some horizontal geodesic connects the past orbits of the endpoints of the bigon. We choose such a horizontal geodesic h satisfying $f(h) \leq \min_{x \in g \cup g'} f(x) - C_{9.1}(J, J')$. Proposition 9.1 gives a $(C_{9.1}(J, J'), C_{9.1}(J, J'))$ -quasi geodesic bigon, with the same vertices, which is $C_{9.1}(J, J')$ -close to $g \cup g'$. We denote the sides of this bigon by \mathcal{G} and \mathcal{G}' .

Let us call a *diagonal* a horizontal geodesic which minimizes the horizontal distance between the future and past orbits of its endpoints. From the hyperbolicity of the semi-flow, any diagonal with horizontal length at least M is dilated both in the future and in the past after $2t_0$.

We choose a real number $L_0 \geq C_{5.4}(2,3,\lambda_+^{2t_0}C_{5.3}(2)) \geq M$ (the meaning of the constant $C_{5.4}(2,3,\lambda_+^{2t_0}C_{5.3}(2))$ will become clear later). Let $P \in \mathcal{G}$. We assume that there exist two points $P_1, P_2 \in h$, whose future orbits intersect \mathcal{G} , such that P is at telescopic distance $L_1 > C_{8.1}(L_0, C_{9.1}(J, J'), C_{9.1}(J, J'))$ from $O^+(P_i) \cup O^-(P_i)$, i = 1, 2.

We consider a diagonal D between $O^+(P_1) \cup O^-(P_1)$ and $O^+(P_2) \cup O^-(P_2)$. This diagonal is in fine position with respect to h. Since \mathcal{G} is in fine position with respect to h, and D connects the future or past orbits of points in h, and the future or past orbits of points in \mathcal{G} , then \mathcal{G} is in fine position with respect to D. Since the point P is at telescopic distance $L_1 > C_{8.1}(L_0, C_{9.1}(J, J'), C_{9.1}(J, J'))$ from $O^+(P_1) \cup O^-(P_1)$ and from $O^+(P_2) \cup O^-(P_2)$, Proposition 8.1 implies that $|D|_{f(D)} > L_0$.

Since \mathcal{G} is in fine position with respect to D, and connects the union of the future and past orbits of the endpoints of D, some horizontal geodesics connect $P \in \mathcal{G}$ to $O^+(P_1)$ and to $O^+(P_2)$. Either these horizontal geodesics are contained in the pulled-tight image of D, or some pulled-tight image of their concatenation contains D. Because of the bounded-cancellation and bounded-dilatation properties, the telescopic distance between a point and an orbit tends to infinity with the horizontal distance between this point and that orbit. Since the telescopic distance between P and $O^+(P_1) \cup O^-(P_1)$, and between P and $O^+(P_2) \cup O^-(P_2)$ is L_1 , this simple observation gives an upper bound X, depending on L_1 , for the horizontal length of each of these horizontal geodesics. Therefore some horizontal geodesic connecting $O^+(P_1) \cup O^-(P_1)$ to $O^+(P_2) \cup O^-(P_2)$ has horizontal length at most equal to

some constant 2X (depending on L_1). In particular, $|D|_{f(D)} \leq 2X$.

We observed that a diagonal D with $|D|_{f(D)} \ge M$ is dilated both in the future and in the past after $2t_0$. Here $|D|_{f(D)} > L_0 \ge M$. Since the concatenation of the above two horizontal geodesics, which lie in the future or in the past of D, has horizontal length at most 2X, a straightforward computation gives Y > 0, still depending on L_1 , such that $|f(P) - f(D)| \le Y$. Lemma 5.6 then implies that P is at telescopic distance smaller than $C_{5.6}(Y)$ from some point in D.

Since \mathcal{G}' and D are in fine position, if no point of \mathcal{G}' lies in the future or past orbit of an endpoint of D, this endpoint belongs to a cancellation. Thus we can write $D = D_1 D_2 D_3$, where

- D_1 (resp. D_3) is non trivial if and only if no point of \mathcal{G}' lies in the future or past orbit of the initial (resp. terminal) point of D.
- D_1 and D_3 , if non trivial, are contained in cancellations.
- \mathcal{G}' connects the future or past orbits of the endpoints of D_2 .

Let us assume that D_1 and D_3 are both trivial. Then, since $2X \ge |D|_{f(D)} \ge L_0$, Proposition 8.1 tells us that some subpath of \mathcal{G}' is $C_{8.1}(2X, C_{9.1}(J, J'), C_{9.1}(J, J'))$ -close to the orbit-segments which connect its endpoints to the endpoints of D. We observed that D is dilated both in the future and in the past after $2t_0$. We proved that $2X \ge |D|_{f(D)} \ge L_0$. An easy computation gives a time t_* after which the pulled-tight images and the geodesic preimages of D have horizontal length at least $3C_{8.1}(2X, C_{9.1}(J, J'), C_{9.1}(J, J'))$. Thus some point Q of the above subpath of \mathcal{G}' satisfies $|f(Q) - f(D)| \le t_*$. Lemma 5.6 gives $C_{5.6}(t_*)$ such that Q is $C_{5.6}(t_*)$ -close to D. Therefore $P \in \mathcal{G}$ and $Q \in \mathcal{G}'$ are $C_{5.6}(t_*) + C_{5.6}(Y) + X$ -close.

Consider now $D=D_1D_2D_3$ with D_1 or D_3 non trivial. Since $|D|_{f(D)} \ge C_{5.4}(2,3,\lambda_+^{2t_0}C_{5.3}(2))$, and D is dilated in the future after $2t_0$, Lemmas 5.3 and 5.4, together with the bounded-dilatation property, give $|D_2|_{f(D)} \ge \lambda_+^{-2t_0}\lambda_+^{2t_0}C_{5.3}(2) \ge M$. Also obviously $|D_2|_{f(D)} \le 2X$. As in the case where D_1 and D_3 are trivial, on replacing D by D_2 in the above arguments, Proposition 8.1 and Lemma 5.6 eventually give a constant $C_{5.6}(t_0)$ such that some point $Q \in \mathcal{G}'$ is $C_{5.6}(t_0)$ -close to D_2 . Thus $P \in \mathcal{G}$ and $Q \in \mathcal{G}'$ are $C_{5.6}(t_0) + C_{5.6}(Y) + X$ -close.

Consider now the case in which the points P_1 , P_2 do not exist. Then P is L_1 -close to some point P' in the orbit of an endpoint, say a, of the bigon. By arguing as above (putting paths in fine position and applying Proposition 8.1), we find a horizontal geodesic h', with one endpoint in the future or past orbit

of a, such that both paths \mathcal{G} and \mathcal{G}' have one point A-close to h', for some constant A. Since \mathcal{G} and \mathcal{G}' both end or begin at the point a, this implies that \mathcal{G}' admits a point B-close to each point of the orbit-segment between a and h'. In particular there exists $Q \in \mathcal{G}'$ which is $B + L_1$ -close to $P \in \mathcal{G}$.

It remains to consider the case where no horizontal geodesic connects the past orbits of the endpoints of the considered (J,J')-quasi geodesic bigon. Then, in the future orbit of the initial endpoint there exists a point z whose past orbit can be connected to the past orbit of the terminal endpoint, and this property is not satisfied by the point w with $f(z) - f(w) = t_0$, which is either in the future or past orbit of the initial endpoint. The strong hyperbolicity of the semi-flow and Proposition 8.1 then give a constant $C_{8.1}(M,J,J')$ such that initial subpaths of both sides of the bigon are $C_{8.1}(M,J,J') + t_0$ -close to the orbit-segment connecting the initial endpoint of the bigon to z. From what precedes, any (R,R')-quasi geodesic bigon between z and the terminal endpoint of the considered bigon is X(R,R')-thin, for some constant X(R,R'). This easily implies that the given bigon is $2(C_{8.1}(M,J,J') + t_0) + X(R,R' + C_{8.1}(M,J,J') + t_0)$ -thin. \square

11. GEODESIC TRIANGLES ARE THIN

The following lemma was suggested to the author by I. Kapovich, and allows us to simplify the conclusion. Let us recall that, in the context of quasi geodesic metric spaces, an (r', s')-chain bigon is a bigon whose sides are (r', s')-chains. Still with this terminology, an (r, s)-chain triangle is a triangle whose sides are (r, s)-chains.

LEMMA 11.1. Let X be an (r,s)-quasi geodesic metric space. If (r',s')-chain bigons are $\delta(r',s')$ -thin, $r' \geq r$, $s' \geq s$, then X is $2\delta(r,3s)$ -hyperbolic.

Proof. We consider an (r,s)-chain triangle with vertices a, b, c and sides [ab], [ac] and [bc]. We consider a point x in the (r,s)-chain [ab] which is closest to c. We claim that $[cx] \cup [xb]$ is an (r,3s)-chain, where [cx] and [xb] denote (r,s)-chains from c to x and from x to b. Indeed, for any points u,v in [xb] or [cx], one obviously has $rd_X(u,v) \ge |[uv]|_X$. Let us