

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	49 (2003)
<b>Heft:</b>	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	 HYPERBOLICITY OF MAPPING-TORUS GROUPS AND SPACES
<b>Autor:</b>	Gautero, François
<b>Kapitel:</b>	13.1 Relationships with mapping-telescopes
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-66690">https://doi.org/10.5169/seals-66690</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

**DEFINITION 13.1** ([19], [3]). An injective endomorphism  $\alpha$  of the rank  $n$  free group  $F_n$  is hyperbolic if there exist  $\lambda_\alpha > 1$  and  $j_\alpha > 0$  such that for any  $w \in F_n$ , either  $\lambda_\alpha |w| \leq |\alpha^{j_\alpha}(w)|$  or  $w$  admits a preimage  $\alpha^{-j_\alpha}(w)$  such that  $\lambda_\alpha |w| \leq |\alpha^{-j_\alpha}(w)|$ , where  $|.|$  denotes the usual word-metric.

We recall that a subgroup  $H$  in a group  $G$  is *malnormal* if  $w^{-1}Hw \cap H = \{1\}$  for any element  $w \notin H$  of  $G$ . We state our theorem about mapping-torus groups as follows:

**THEOREM 13.2.** *Let  $\alpha$  be an injective hyperbolic endomorphism of the rank  $n$  free group  $F_n$ . If the image of  $\alpha$  is a malnormal subgroup of  $F_n$  then the mapping-torus group  $G_\alpha = \langle x_1, \dots, x_n, t ; t^{-1}x_i t = \alpha(x_i), i = 1, \dots, n \rangle$  is a hyperbolic group.*

### 13.1 RELATIONSHIPS WITH MAPPING-TELESCOPES

We consider the rank  $n$  free group  $F_n = \langle x_1, \dots, x_n \rangle$ . Let  $\alpha$  be an injective endomorphism of  $F_n$ . Let  $G_\alpha = \langle x_1, \dots, x_n, t ; t^{-1}x_i t = \alpha(x_i), i = 1, \dots, n \rangle$  be the mapping-torus group of  $(\alpha, F_n)$ . We consider the Cayley graph  $\Gamma$  associated to the given system of generators. Let  $l$  be a loop in  $\Gamma$  whose associated word in the edges of  $\Gamma$  reads a relation  $t^{-1}x_i t \alpha(x_i)^{-1}$ . We attach a 2-cell by its boundary circle along any such loop  $l$ . The resulting topological space is a 2-complex. This is the Cayley complex of the mapping-torus group  $G_\alpha$  for the given presentation.

Let us check that the above Cayley complex is a mapping-telescope of a forest-map. We consider the rose  $\mathcal{R}_n$  with  $n$  petals. We label each edge by a generator  $x_i$  of  $F_n$ . We denote by  $\psi$  the simplicial map on  $\mathcal{R}_n$  such that  $\psi(x_i)$  is a locally injective path whose associated word in the edges of  $\mathcal{R}_n$  reads  $\alpha(x_i)$ . Let us denote by  $T$  the universal covering of  $\mathcal{R}_n$  ( $T$  is a tree) and by  $\pi: T \rightarrow \mathcal{R}_n$  the associated covering-map. We denote by  $\widehat{\psi}: T \rightarrow T$  a simplicial lift of  $\psi$  to  $T$ , that is  $\pi \circ \widehat{\psi} = \psi \circ \pi$ . We consider the mapping-torus of  $(\psi, \mathcal{R}_n)$ , i.e. the 2-complex  $\mathcal{R}_n \times [0, 1]/(x, 1) \sim (\psi(x), 0)$ . Then the universal covering of this mapping-torus is the mapping-telescope of  $\widetilde{\psi}: F \rightarrow F$ , where  $F$  and  $\widetilde{\psi}$  are defined as follows:

- We denote by  $I$  the set of integers from 1 to  $\text{Card}(F_n / \text{Im}(\alpha))$ . The different classes are written  $w_i \text{Im}(\alpha)$ ,  $i = 0, 1, \dots$ . We denote by  $\gamma: I \rightarrow \{w_0, w_1, \dots\}$  the bijection. Then the connected components of  $F$  are in bijection with  $\mathbf{N}^{\text{Card}(I)}$ . Each connected component is the image, by a

bijection  $\mu$ , of a sequence of  $\text{Card}(I)$  integers. Each connected component  $\mu(x_0, x_1, \dots)$  of  $F$  is homeomorphic to  $T$  via  $\beta_{(x_0, x_1, \dots)} : \mu(x_0, x_1, \dots) \rightarrow T$ .

- We define the restriction of  $\tilde{\psi}$  to any connected component  $\mu((x_0, x_1, \dots))$  as follows :

If  $\text{Card}(I) < +\infty$  then

$$\tilde{\psi}|_{\mu((x_0, x_1, \dots))} : \begin{cases} \mu((x_0, x_1, \dots)) & \rightarrow \quad \mu(E[\frac{x_0}{\text{Card}(I)}], x_1, \dots) \\ x & \rightarrow \quad (\gamma(j)\beta_{(x_0, x_1, \dots)}^{-1}\widehat{\psi}\beta_{(x_0, x_1, \dots)})(x) \end{cases}$$

where  $j < \text{Card}(I)$  satisfies  $E[\frac{x_0}{\text{Card}(I)}] = k \text{Card}(I) + j$ .

If  $\text{Card}(I) = +\infty$  then

$$\tilde{\psi}|_{\mu((x_0, x_1, \dots))} : \begin{cases} \mu((x_0, x_1, \dots)) & \rightarrow \quad \mu((x_1, x_2, \dots)) \\ x & \rightarrow \quad (\gamma(x_0)\beta_{(x_0, x_1, \dots)}^{-1}\widehat{\psi}\beta_{(x_0, x_1, \dots)})(x). \end{cases}$$

The mapping-torus of  $(\psi, \mathcal{R}_n)$  is a 2-complex whose 1-skeleton is the rose with  $n + 1$  petals in bijection with  $\{x_1, \dots, x_n, t\}$ . There is one 2-cell for each relation  $t^{-1}x_i t \alpha(x_i)^{-1}$ . Thus the universal covering described above is the Cayley complex for  $G_\alpha$  with the presentation  $G_\alpha = \langle x_1, \dots, x_n, t ; t^{-1}x_i t = \alpha(x_i), i = 1, \dots, n \rangle$ . We have thus proved

**LEMMA 13.3.** *Let  $\alpha$  be an injective endomorphism of  $F_n = \langle x_1, \dots, x_n \rangle$ . Let  $G_\alpha = \langle x_1, \dots, x_n, t ; t^{-1}x_i t = \alpha(x_i), i = 1, \dots, n \rangle$  be the mapping-torus group of  $\alpha$ . Let  $\mathcal{C}(G_\alpha)$  be the Cayley complex of  $G_\alpha$  for the given presentation. Then  $\mathcal{C}(G_\alpha)$  is the mapping-telescope of a forest-map.*

**REMARK 13.4.** If the endomorphism  $\alpha$  is an automorphism then the above Cayley complex is the mapping-telescope of a tree-map. The tree is the universal covering of the rose with  $n$  petals. If the endomorphism  $\alpha$  is not injective then some element  $w \in F_n$  satisfies  $w = 1$  in  $G_\alpha$ ; the above construction fails because of the corresponding loops in the Cayley graph.

Let  $\alpha$  be an injective free group endomorphism. Let  $G_\alpha$  be the mapping-torus group of  $\alpha$ . Let  $\mathcal{C}(G_\alpha)$  be the Cayley complex of  $G_\alpha$  for the usual presentation  $G_\alpha = \langle x_1, \dots, x_n, t ; t^{-1}x_i t = \alpha(x_i), i = 1, \dots, n \rangle$ . By Lemma 13.3,  $\mathcal{C}(G_\alpha)$  is a mapping-telescope of a forest-map. We now want to see what happens with respect to metrics and dynamics. The Cayley graph of a group is equipped with a metric which makes each edge isometric to the interval  $(0, 1)$ . More generally, given a graph  $\Gamma$ , we call *standard metric*, and denote

by  $d_\Gamma^s$ , such a metric on  $\Gamma$ . We will call *mapping-telescope standard metric* any mapping-telescope  $d_\Gamma^s$ -metric on  $\mathcal{C}(G_\alpha)$ .

**LEMMA 13.5.** *The mapping-torus group  $G_\alpha$  of an injective free group endomorphism acts cocompactly, properly discontinuously and isometrically on the Cayley complex  $\mathcal{C}(G_\alpha)$  equipped with any mapping-telescope standard metric.*

*Proof.* We consider the usual action by left translations of the group on its Cayley graph. This action is extended in a natural way to a free action on the Cayley complex  $\mathcal{C}(G_\alpha)$ . Let  $f$  denote the map giving the strata for the structure of forest-stack of  $\mathcal{C}(G_\alpha)$ , see Lemma 13.3. For a mapping-telescope metric, all the strata  $f^{-1}(r)$  and  $f^{-1}(r+1)$  are isometric. And for a mapping-telescope standard metric all the strata  $f^{-1}(n)$ ,  $n \in \mathbf{Z}$ , are equipped with the standard metric. This readily implies that the above action is isometric.  $\square$

### 13.2 FREE GROUP ENDOMORPHISMS AND FOREST-MAPS

The main point of Lemma 13.6 below is the so-called ‘bounded-cancellation lemma’ of [7] for free group automorphisms, and of [10] for the injective free group endomorphisms.

**LEMMA 13.6.** *Let  $\alpha$  be an injective free group endomorphism. Let  $F$  and  $\tilde{\psi}$  be the forest and the forest-map on  $F$  given by Lemma 13.3. Then  $\tilde{\psi}$  is a weakly bi-Lipschitz forest-map of  $F$  equipped with the standard metric  $d_F^s$ .*

*Proof.* If  $w$  is any element in  $F_n = \langle x_1, \dots, x_n \rangle$ , and  $|.|_{F_n}$  denotes the word-metric on  $F_n$ , then  $|\alpha(w)|_{F_n} \leq (\max_{i=1, \dots, n} |\alpha(x_i)|_{F_n})|w|_{F_n}$ . By definition of the standard metric, and setting  $\mu_0 = \max_{i=1, \dots, n} |\alpha(x_i)|_{F_n}$ , the map  $\tilde{\psi}$  satisfies  $d_F^s(\tilde{\psi}(x), \tilde{\psi}(y)) \leq \mu_0 d_F^s(x, y)$  for any pair of vertices  $x, y$ . If  $x, y$  are not vertices, then they are joined in their stratum by a horizontal geodesic which is the concatenation of a path between two vertices, with two proper subsets of edges. By construction and simpliciality of  $\tilde{\psi}$ , proper subsets of edges are dilated by a bounded factor when applying  $\tilde{\psi}$ , so that the conclusion follows for the upper bound.

If  $w$  is any element in  $F_n$  then

$$|\alpha^{-1}(w)|_{F_n} \leq (\max_{i=1, \dots, n} |\alpha^{-1}(x_i)|_{F_n})|w|_{F_n}.$$

Setting  $\mu_1 = \max_{i=1, \dots, n} |\alpha^{-1}(x_i)|_{F_n}$  we get  $|\alpha(w)|_{F_n} \geq \frac{1}{\mu_1}|w|_{F_n}$ . Therefore  $d_F^s(\tilde{\psi}(x), \tilde{\psi}(y)) \geq \frac{1}{\mu_1}d_F^s(x, y)$  for any pair of vertices  $x, y$ . The inequality