Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE BASIC GERBE OVER A COMPACT SIMPLE LIE GROUP
Autor: Meinrenken, Eckhard
Kapitel: 2.1 Chatterjee-Hitchin gerbes
DOI: https://doi.org/10.5169/seals-66691

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

The L_{ij}, together with these isomorphisms, define a gerbe over $\mathrm{SU}(d+1)$, representing the generator of $H^3(\mathrm{SU}(d+1), \mathbf{Z})$.

More generally, consider any compact, simply connected, simple Lie group G of rank d. Up to conjugacy, G contains exactly $d+1$ elements with semi-simple centralizer. (For $G = \mathrm{SU}(d+1)$, these are the central elements.) Let $\mathcal{C}_1, \ldots, \mathcal{C}_{d+1} \subset G$ be their conjugacy classes. We will define an invariant open cover $V_1, \ldots, V_{d+1}$ of G, with the property that each member of this cover admits an equivariant retraction onto the conjugacy class $\mathcal{C}_j \subset V_j$. It turns out that every semi-simple centralizer has a distinguished central extension by $\mathrm{U}(1)$. This central extension defines an equivariant bundle gerbe on $\mathcal{C}_j$, hence (by pull-back) an equivariant bundle gerbe over V_j. We will find that these gerbes over V_j glue together to produce a gerbe over G, using a gluing rule developed in this paper.

The organization of the paper is as follows. In Section 2 we review the theory of gerbes and pseudo-line bundles with connections, and discuss 'strong equivariance' under a group action. Section 4 describes gluing rules for bundle gerbes. Section 3 summarizes some facts about gerbes coming from central extensions. In Section 5 we give the construction of the basic gerbe over G outlined above, and in Section 6 we study the 'pre-quantization of conjugacy classes'.

ACKNOWLEDGEMENTS. I would like to thank Ping Xu for fruitful discussions at the Poisson 2002 meeting in Lisbon, and for a preliminary version of his preprint [2] with Behrend and Zhang, giving yet another construction of the basic gerbe over G. Their (infinite-dimensional) approach is based on the notion of Morita equivalence of (quasi-)symplectic groupoids. I thank the referees for detailed comments and suggestions.

2. GERBES WITH CONNECTIONS

In this section we review gerbes on manifolds, along the lines of Chatterjee-Hitchin and Murray.

2.1 CHATTERJEE-HITCHIN GERBES

Let M be a manifold. Any Hermitian line bundle over M can be described by an open cover U_a, and transition functions $\chi_{ab}\colon U_a \cap U_b \to \mathrm{U}(1)$ satisfying a cocycle condition $(\delta\chi)_{abc} = \chi_{bc}\chi_{ac}^{-1}\chi_{ab} = 1$ on triple intersections. The

cohomology class in $H^1(M, \underline{\mathrm{U}(1)}) = H^2(M, \mathbf{Z})$ defined by this cocycle is the Chern class of the line bundle. Chatterjee-Hitchin [10, 18, 17] suggested to realize classes in $H^3(M, \mathbf{Z})$ in a similar fashion, replacing U(1)-valued functions with Hermitian line bundles. They define a gerbe to be a collection of Hermitian transition line bundles $L_{ab} \to U_a \cap U_b$ and a trivialization, i.e. unit length section, t_{abc} of the line bundle $(\delta L)_{abc} = L_{bc} L_{ac}^{-1} L_{ab}$ over triple intersections. These trivializations have to satisfy a compatibility relation over quadruple intersections,

$$(\delta t)_{abcd} \equiv t_{bcd} t_{acd}^{-1} t_{abd} t_{abc}^{-1} = 1\,,$$

which makes sense since $(\delta t)_{abcd}$ is a section of the *canonically* trivial bundle. (Each factor L_{ab} cancels with a factor L_{ab}^{-1}.) After passing to a refinement of the cover, such that all L_{ab} become trivializable, and picking trivializations, t_{abc} is simply a Čech cocycle of degree 2, hence defines a class in $H^2(M, \underline{\mathrm{U}(1)}) = H^3(M, \mathbf{Z})$. The class is independent of the choices made in this construction, and is called the *Dixmier-Douady class* of the gerbe.

Note that in practice, it is often not desirable to pass to a refinement. For example, if M is a connected, oriented 3-manifold, the generator of $H^3(M, \mathbf{Z}) = \mathbf{Z}$ can be described in terms of the cover U_1, U_2, where U_1 is an open ball around a given point $p \in M$, and $U_2 = M \backslash \{p\}$, using the degree one line bundle over $U_1 \cap U_2 \cong S^2 \times (0, 1)$.

2.2 Bundle gerbes

Bundle gerbes were invented by Murray [24], generalizing the following construction of line bundles. Let $\pi\colon X \to M$ be a fiber bundle, or more generally a surjective submersion. (Different components of X may have different dimensions.) For each $k \geq 0$ let $X^{[k]}$ denote the k-fold fiber product of X with itself. There are $k+1$ projections $\partial^i\colon X^{[k+1]} \to X^{[k]}$, omitting the ith factor in the fiber product. Suppose we are given a smooth function $\chi\colon X^{[2]} \to \mathrm{U}(1)$, satisfying a cocycle condition $\delta\chi = 1$ where

$$\delta\chi := \partial_0^* \chi \partial_1^* \chi^{-1} \partial_2^* \chi\colon X^{[3]} \to \mathrm{U}(1)\,.$$

Then χ determines a Hermitian line bundle $L \to M$, with fibers at $m \in M$ the space of all linear maps $\phi\colon X_m = \pi^{-1}(m) \to \mathbf{C}$ such that $\phi(x) = \chi(x, x')\phi(x')$. Given local sections $\sigma_a\colon U_a \to X$ of X, the pull-backs of χ under the maps $(\sigma_a, \sigma_b)\colon U_a \cap U_b \to X^{[2]}$ give transition functions χ_{ab} for the line bundle.

Again, replacing U(1)-valued functions by line bundles in this construction, one obtains a model for gerbes: A bundle gerbe is given by a line bundle $L \to X^{[2]}$ and a trivializing section t of the line bundle $\delta L = \partial_0^* L \otimes \partial_1^* L^{-1} \otimes \partial_2^* L$