Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE BASIC GERBE OVER A COMPACT SIMPLE LIE GROUP
Autor: Meinrenken, Eckhard
Kapitel: 5.2 The basic 3-form on G
DOI: https://doi.org/10.5169/seals-66691

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

5.2 THE BASIC 3-FORM ON G

Let $\theta^L, \theta^R \in \Omega^1(G, \mathfrak{g})$ be the left- and right-invariant Maurer-Cartan forms on G, respectively. The 3-form $\eta \in \Omega^3(G)$ given by[3])

$$\eta = \frac{1}{12}\theta^L \cdot [\theta^L, \theta^L] = \frac{1}{12}\theta^R \cdot [\theta^R, \theta^R]$$

is closed, and has a closed equivariant extension $\eta_G \in \Omega^3_G(G)$ given by

$$\eta_G(\xi) := \eta - \tfrac{1}{2}(\theta^L + \theta^R) \cdot \xi\,.$$

Their cohomology classes represent generators of $H^3(G, \mathbf{Z}) = \mathbf{Z}$ and $H^3_G(G, \mathbf{Z}) = \mathbf{Z}$, respectively. The pull-back of η_G to any conjugacy class $\iota_{\mathcal{C}} : \mathcal{C} \hookrightarrow G$ is exact. In fact, let $\omega_{\mathcal{C}} \in \Omega^2(\mathcal{C})^G \subset \Omega^2_G(\mathcal{C})$ be the invariant 2-form given on generating vector fields $\xi_{\mathcal{C}}, \xi'_{\mathcal{C}}$ for $\xi, \xi' \in \mathfrak{g}$ by the formula

$$\omega_{\mathcal{C}}(\xi_{\mathcal{C}}(g), \xi'_{\mathcal{C}}(g)) = \tfrac{1}{2}\xi \cdot (\mathrm{Ad}_g - \mathrm{Ad}_{g^{-1}})\xi'\,.$$

Then [1, 16]

$$\mathrm{d}_G\, \omega_{\mathcal{C}} + \iota^*_{\mathcal{C}} \eta_G = 0\,.$$

We will now show that η_G is exact over each of the open subsets V_j. Let $\mathcal{C}_j = q^{-1}(\mu_j) \subset V_j$ be the conjugacy classes corresponding to the vertices.

LEMMA 5.1. *The linear retraction*

$$[0,1] \times \mathfrak{A}_j \to \mathfrak{A}_j, \quad (t, \mu_j + \zeta) \mapsto \mu_j + (1-t)\zeta$$

of $\mathfrak{A}_j$ onto the vertex μ_j lifts uniquely to a smooth G-equivariant retraction from V_j onto $\mathcal{C}_j$.

Proof. Recall that the slice S_j is an open neighborhood of $\exp(\mu_j)$ in G_j. Any G_j-equivariant retraction from S_j onto $\exp \mu_j$ extends uniquely to a G-equivariant retraction from $V_j = G \times_{G_j} S_j$ onto $\mathcal{C}_j$. Note that $S'_j = G_j \,.\, (\mathfrak{A}_j - \mu_j)$ is a star-shaped open neighborhood of 0 in $\mathfrak{g}_j$, and that $S'_j \to S_j$, $\zeta \mapsto \exp(\mu_j)\exp(\zeta)$ is a G_j-equivariant diffeomorphism. The linear retraction of S'_j onto the origin gives the desired retraction of S_j. Uniqueness is clear, since the retraction has to preserve $\exp(\mathfrak{A}_j) \subset V_j$, by equivariance.

---

[3]) For $\mathfrak{g}$-valued forms β_1, β_2, the bracket $[\beta_1, \beta_2]$ denotes the $\mathfrak{g}$-valued form obtained by applying the Lie bracket $[\cdot,\cdot]\colon \mathfrak{g} \otimes \mathfrak{g} \to \mathfrak{g}$ to the $\mathfrak{g} \otimes \mathfrak{g}$-valued form $\beta_1 \wedge \beta_2$.

Let

$$\mathbf{h}_j\colon \Omega^p(V_j) \to \Omega^p([0,1]\times V_j) \to \Omega^{p-1}(V_j)$$

be the de Rham homotopy operator for this retraction, given (up to a sign) by pull-back under the retraction, followed by integration over the fibers of $[0,1]\times V_j \to V_j$. It has the property

$$\mathrm{d}_G\,\mathbf{h}_j + \mathbf{h}_j\,\mathrm{d}_G = \mathrm{Id} - \pi_j^*\iota_j^* \tag{5.1}$$

where $\iota_j\colon \mathcal{C}_j \to V_j$ is the inclusion and $\pi_j\colon V_j = G\times_{G_j} S_j \to G/G_j = \mathcal{C}_j$ the projection. Let $(\varpi_j)_G = \mathbf{h}_j\eta_G - \pi_j^*\omega_{\mathcal{C}_j} \in \Omega^2_G(V_j)$, and write $(\varpi_j)_G = \varpi_j - \Psi_j$ where $\varpi_j \in \Omega^2(V_j)$ and $\Psi_j \in \Omega^0(V_j,\mathfrak{g})$.

PROPOSITION 5.2. *The equivariant 2-form $(\varpi_j)_G = \varpi_j - \Psi_j$ has the following properties.*

(a) $\mathrm{d}_G(\varpi_j)_G = \eta_G$.

(b) *The pull-back of $(\varpi_j)_G$ to a conjugacy class $\mathcal{C}\subset V_j$ is given by*

$$\iota_{\mathcal{C}}^*(\varpi_j)_G = \Psi_j^*(\omega_{\mathcal{O}})_G - \omega_{\mathcal{C}}\,,$$

where $(\omega_{\mathcal{O}})_G$ is the equivariant symplectic form on the adjoint orbit $\mathcal{O} = \Psi_j(\mathcal{C})$,

(c) *The pull-back of Ψ_j to the conjugacy class $\mathcal{C}_j$ vanishes. In fact, $\Psi_j(\exp\xi) = \xi - \mu_j$ for all $\xi\in\mathfrak{A}_j$.*

(d) *Over each intersection $V_{ij} = V_i\cap V_j$, the difference $\Psi_i - \Psi_j$ takes values in the adjoint orbit $\mathcal{O}_{ij}$ through $\mu_j - \mu_i \in \mathfrak{g}\cong\mathfrak{g}^*$. Furthermore,*

$$(\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G$$

where $p_{ij}\colon V_{ij}\to\mathcal{O}_{ij}$ is the map defined by $\Psi_i - \Psi_j$, and $(\omega_{\mathcal{O}_{ij}})_G$ is the equivariant symplectic form on the orbit.

Proof. (a) holds by construction. (b) follows from the observation that $\iota_{\mathcal{C}}^*(\varpi_j)_G + \omega_{\mathcal{C}}$ is an equivariantly closed 2-form on $\mathcal{C}_j$, with Ψ_j as its moment map. To prove (c) we note that since the retraction is equivariant, we have $\tilde{\mathbf{h}}_j\circ(\exp|_{\mathfrak{A}_j})^* = (\exp|_{\mathfrak{A}_j})^*\circ\mathbf{h}_j$ where $(\exp|_{\mathfrak{A}_j})^*$ is pull-back to $\mathfrak{A}_j\subset\mathfrak{t}$ and where $\tilde{\mathbf{h}}_j$ is the homotopy operator for the linear retraction of $\mathfrak{t}$ onto $\{\mu_j\}$. Let $\nu\colon\mathfrak{A}_j\to\mathfrak{t}$ be the coordinate function (inclusion). Then

$$\tilde{\mathbf{h}}_j\circ(\exp|_{\mathfrak{A}_j})^*\tfrac{1}{2}(\theta^L+\theta^R) = \tilde{\mathbf{h}}_j\circ\mathrm{d}\,\nu = \nu - \mu_j\,,$$

proving that $(\exp|_{\mathfrak{A}_j})^*\Psi_j = \nu - \mu_j$. This yields (c), by equivariance. For $\nu\in\mathfrak{A}_{ij}$ we have, using (c),

$$(\Psi_i - \Psi_j)(\exp \nu) = (\nu - \mu_i) - (\nu - \mu_j) = \mu_j - \mu_i .$$

By equivariance, it follows that $\Psi_i - \Psi_j$ takes values in the adjoint orbit through $\mu_j - \mu_i$. The difference $\varpi_i - \varpi_j$ vanishes on the maximal torus T, and is therefore determined by its contractions with generating vector fields. Since $\Psi_i - \Psi_j$ is a moment map for $\varpi_i - \varpi_j$, it follows that $\varpi_i - \varpi_j$ equals the pull-back of the symplectic form on $G.(\mu_j - \mu_i)$.

5.3 The special unitary group

For the special unitary group $G = \mathrm{SU}(d+1)$, the construction of the basic gerbe simplifies due to the fact that in this case all vertices μ_j of the alcove are contained in the weight lattice. In fact the gerbe is presented as a Chatterjee-Hitchin gerbe for the cover $\mathcal{V} = \{V_i,\ i = 0, \ldots, d\}$.

For each weight $\mu \in \Lambda^* \subset \mathfrak{t} \subset \mathfrak{g}$, let G_μ be its stabilizer for the adjoint action and let $\mathbf{C}_\mu$ the 1-dimensional G_μ-representation with infinitesimal character μ. Let the line bundle $L_\mu = G \times_{G_\mu} \mathbf{C}_\mu$ equipped with the unique left-invariant connection ∇. Then L_μ is a G-equivariant pre-quantum line bundle for the orbit $\mathcal{O} = G.\mu$. That is,

$$\frac{i}{2\pi} \operatorname{curv}_G(\nabla) = (\omega_{\mathcal{O}})_G := \omega_{\mathcal{O}} - \Phi_{\mathcal{O}}$$

where $\omega_{\mathcal{O}}$ is the symplectic form and $\Phi_{\mathcal{O}} \colon \mathcal{O} \hookrightarrow \mathfrak{g}^*$ is the moment map given as inclusion.

In particular, in the case of $\mathrm{SU}(d+1)$ all orbits $\mathcal{O}_{ij} = G.(\mu_j - \mu_i)$ carry G-equivariant pre-quantum line bundles. Recall the fibrations $p_{ij} \colon V_{ij} \to \mathcal{O}_{ij}$ defined by $\Psi_i - \Psi_j$, and let

$$L_{ij} = p_{ij}^*(L_{\mu_j - \mu_i}),$$

equipped with the pull-back connection. For any triple intersection $V_{ijk} = G \times_{G_{ijk}} S_{ijk}$, the tensor product $(\delta L)_{ijk} = L_{jk} L_{ik}^{-1} L_{ij}$ is the pull-back of the line bundle over G/G_{ijk}, defined by the zero weight

$$(\mu_k - \mu_j) - (\mu_k - \mu_i) + (\mu_j - \mu_i) = 0$$

of G_{ijk}. It is hence canonically trivial, with $(\delta\nabla)_{ijk}$ the trivial connection. The trivializing section $t_{ijk} = 1$ satisfies $\delta t = 1$ and $(\delta\nabla)t = 0$. Take $(B_j)_G = (\varpi_j)_G$. Then

$$(B_j)_G - (B_i)_G = (\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G = \frac{1}{2\pi i} \operatorname{curv}_G(\nabla^{L_{ij}}).$$

Thus $\mathcal{G} = (\mathcal{V}, L, t)$ is a equivariant gerbe with connection (∇, B). Since