Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 49 (2003)

Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE BASIC GERBE OVER A COMPACT SIMPLE LIE GROUP

Autor: Meinrenken, Eckhard

Kapitel: 5.3 The special unitary group

DOI: https://doi.org/10.5169/seals-66691

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$(\Psi_i - \Psi_j)(\exp \nu) = (\nu - \mu_i) - (\nu - \mu_j) = \mu_j - \mu_i$$
.

By equivariance, it follows that $\Psi_i - \Psi_j$ takes values in the adjoint orbit through $\mu_j - \mu_i$. The difference $\varpi_i - \varpi_j$ vanishes on the maximal torus T, and is therefore determined by its contractions with generating vector fields. Since $\Psi_i - \Psi_j$ is a moment map for $\varpi_i - \varpi_j$, it follows that $\varpi_i - \varpi_j$ equals the pull-back of the symplectic form on G. $(\mu_j - \mu_i)$.

5.3 THE SPECIAL UNITARY GROUP

For the special unitary group $G = \mathrm{SU}(d+1)$, the construction of the basic gerbe simplifies due to the fact that in this case all vertices μ_j of the alcove are contained in the weight lattice. In fact the gerbe is presented as a Chatterjee-Hitchin gerbe for the cover $\mathcal{V} = \{V_i, i = 0, \dots, d\}$.

For each weight $\mu \in \Lambda^* \subset \mathfrak{t} \subset \mathfrak{g}$, let G_{μ} be its stabilizer for the adjoint action and let \mathbf{C}_{μ} the 1-dimensional G_{μ} -representation with infinitesimal character μ . Let the line bundle $L_{\mu} = G \times_{G_{\mu}} \mathbf{C}_{\mu}$ equipped with the unique left-invariant connection ∇ . Then L_{μ} is a G-equivariant pre-quantum line bundle for the orbit $\mathcal{O} = G \cdot \mu$. That is,

$$\frac{i}{2\pi}\operatorname{curv}_G(\nabla) = (\omega_{\mathcal{O}})_G := \omega_{\mathcal{O}} - \Phi_{\mathcal{O}}$$

where $\omega_{\mathcal{O}}$ is the symplectic form and $\Phi_{\mathcal{O}} \colon \mathcal{O} \hookrightarrow \mathfrak{g}^*$ is the moment map given as inclusion.

In particular, in the case of SU(d+1) all orbits $\mathcal{O}_{ij} = G \cdot (\mu_j - \mu_i)$ carry G-equivariant pre-quantum line bundles. Recall the fibrations $p_{ij} \colon V_{ij} \to \mathcal{O}_{ij}$ defined by $\Psi_i - \Psi_j$, and let

$$L_{ij} = p_{ij}^* \left(L_{\mu_j - \mu_i} \right),$$

equipped with the pull-back connection. For any triple intersection $V_{ijk} = G \times_{G_{ijk}} S_{ijk}$, the tensor product $(\delta L)_{ijk} = L_{jk} L_{ik}^{-1} L_{ij}$ is the pull-back of the line bundle over G/G_{ijk} , defined by the zero weight

$$(\mu_k - \mu_i) - (\mu_k - \mu_i) + (\mu_i - \mu_i) = 0$$

of G_{ijk} . It is hence canonically trivial, with $(\delta \nabla)_{ijk}$ the trivial connection. The trivializing section $t_{ijk}=1$ satisfies $\delta t=1$ and $(\delta \nabla)t=0$. Take $(B_i)_G=(\varpi_i)_G$. Then

$$(B_j)_G - (B_i)_G = (\varpi_j)_G - (\varpi_i)_G = -p_{ij}^*(\omega_{\mathcal{O}_{ij}})_G = \frac{1}{2\pi i}\operatorname{curv}_G(\nabla^{L_{ij}}).$$

Thus $\mathcal{G} = (\mathcal{V}, L, t)$ is a equivariant gerbe with connection (∇, B) . Since

$$\mathrm{d}_G(B_j)_G=\mathrm{d}_G(\varpi_j)_G=\eta_G|_{V_j},$$

this is the basic gerbe for SU(d+1). The transition line bundles L_{ij} may be expressed in terms of eigenspace line bundles, leading to the description of the basic gerbe from the introduction.

REMARK 5.3. This description of the basic gerbe over the special unitary group was found independently by Gawędzki-Reis [13], who also discuss the much more difficult case of quotients of SU(d+1) by subgroups of the center.

A similar construction works for the group $C_d = \operatorname{Sp}(d)$, the only case besides $A_d = \operatorname{SU}(d+1)$ for which the vertices of the alcove are in the weight lattice. The following table lists, for all simply connected compact simple groups, the smallest integer $k_0 > 0$ such that $k_0 \mathfrak{A}$ is a weight lattice polytope⁴). The construction for $\operatorname{SU}(d+1)$ generalizes to describe the k_0 -th power of the basic gerbe in all cases.

5.4 The basic gerbe for general simple, simply connected G

The extra difficulty for the groups with $k_0 > 1$ comes from the fact that the pull-back maps $H_G^3(G, \mathbf{Z}) \to H_G^3(\mathcal{C}_j, \mathbf{Z}) \cong H_G^3(V_j, \mathbf{Z})$ may be a non-zero torsion class, in general. In this case the restriction of the basic gerbe to V_j will be non-trivial. Our strategy for the general case is to first construct equivariant bundle gerbes over V_j , and then glue the local data as explained in Section 4.

The centralizers G_g of elements $g \in G$ are always connected [11, Corollary (3.15)] but need not be simply-connected. The conjugacy classes $C_j = q^{-1}(\mu_j)$ corresponding to the vertices of the alcove are exactly the conjugacy classes of elements for which the centralizer is semi-simple. Since

$$H_G^3(\mathcal{C}_j, \mathbf{Z}) = H_G^3(G/G_j, \mathbf{Z}) = H_{G_j}^3(\operatorname{pt}, \mathbf{Z}),$$

we see that the torsion problem described above is related to a possibly non-trivial central extension of the centralizers G_j of $\exp(\mu_j)$ by the circle U(1).

⁴) This information is extracted from the tables in Bourbaki [5]. Letting w_1, \ldots, w_d be the fundamental weights, one determines k_0 as the least common multiple of the numbers $\alpha_{max} \cdot w_j$, using the basic inner product defined by $\alpha_{max} \cdot \alpha_{max} = 2$. The number k_0 is equal to the smallest Dynkin index of a representation $G \to SU(n)$, see [28, p. 128] where the same table appears in a different context.

PROPOSITION 5.4. Any vertex μ_j of the alcove \mathfrak{A} is in the dual of the co-root lattice for the corresponding centralizer G_j . It hence defines a homomorphism $\varrho_j \in \text{Hom}(\pi_1(G_j), U(1))$, or equivalently a central extension of G_j by U(1).

Proof. Let \widetilde{G}_j be the universal cover of G_j . A system of simple roots for \widetilde{G}_j is given by the list of all α_i $(i=0,\ldots,d)$ with $j\neq i$. The lattice Λ_j is spanned by the corresponding coroots $\check{\alpha}_i$. To show that μ_j is in the dual of the co-root lattice, we have to verify that $\langle \mu_j, \check{\alpha}_i \rangle \in \mathbb{Z}$ for $i \neq j$. For $i \neq 0$, j this is obvious since $\mu_j(\check{\alpha}_i) = 0$. For i = 0, we have $\|\check{\alpha}_0\|^2 = 2$, and therefore $\check{\alpha}_0 = \alpha_0$ and $\mu_j(\check{\alpha}_0) = \alpha_0(\mu_j) = -1$.

Recall that for $i \neq j$, G_{ij} is the centralizer of points $\exp \mu$ with $\mu = t\mu_j + (1-t)\mu_i$ for some 0 < t < 1. Let $\varrho_{ij} \in \operatorname{Hom}(\pi_1(G_{ij}), \operatorname{U}(1))$ be the quotient of $\pi_1(G_{ij}) \to \pi_1(G_j) \xrightarrow{\varrho_j} \operatorname{U}(1)$ by the homomorphism $\pi_1(G_{ij}) \to \pi_1(G_i) \xrightarrow{\varrho_i} \operatorname{U}(1)$.

LEMMA 5.5. The difference $\mu_j - \mu_i \in \mathfrak{g}_{ij}$ is fixed under G_{ij} , and $\varrho_{ij} \in \text{Hom}(\pi_1(G_{ij}), U(1))$ is its image under the exact sequence (3.2) for $K = G_{ij}$.

Proof. Since G_{ij} fixes the curve $g(t) = \exp(t\mu_j + (1-t)\mu_i) = \exp(\mu_i)\exp(t(\mu_j - \mu_i))$, it stabilizes the Lie algebra element $\mu_j - \mu_i$. The second claim is immediate from the definition.

We are now in position to explain our construction of the basic gerbe in the general case. For all $I \subset \{0, ..., d\}$ let $X_I \to V_I$ be the G-equivariant principal G_I -bundle,

$$X_I = G \times S_I \rightarrow V_I = G \times_{G_I} S_I$$
.

 X_I is the pull-back of the G_I -bundle $G \to G/G_I$, and in particular carries a G-invariant connection θ_I obtained by pull-back of the unique G-invariant connection on that bundle. For $I \supset J$ there are natural G-equivariant inclusions $f_I^J \colon X_I \to X_J$, and these are compatible as in Section 4. The homomorphisms $\varrho_j \colon \pi_1(G_j) \to \mathrm{U}(1)$ define flat, G-equivariant bundle gerbes $\mathcal{G}_j = (X_j, L_j, t_j)$ over V_j .

The quotient of the two gerbes on V_{ij} , obtained by pulling back \mathcal{G}_i , \mathcal{G}_j to X_{ij} , is just the gerbe defined by the homomorphism $\varrho_{ij} \colon \pi_1(G_{ij}) \to \mathrm{U}(1)$. By Lemma 5.5 and Proposition 3.2(b), it follows that this quotient gerbe has

a distinguished, equivariant pseudo-line bundle (E_{ij}, s_{ij}) (where E_{ij} is trivial), with connection $\nabla^{E_{ij}}$ induced from the connection θ_{ij} . From the definition of θ_{ij} , it follows that the equivariant error 2-form for this connection is the pull-back of the equivariant symplectic form on the coadjoint orbit through $\mu_i - \mu_i$.

We now modify the bundle gerbe connection by adding the equivariant 2-form $(\varpi_j)_G \in \Omega^2_G(V_j)$ to the gerbe connection. Proposition 5.2(d) shows that the equivariant error 2-form of $\nabla^{E_{ij}}$ with respect to the new gerbe connection vanishes. The other conditions from the gluing construction in §4 are trivially satisfied. Since the equivariant 3-curvature for the new gerbe connection on \mathcal{G}_j is $\mathrm{d}_G(\varpi_j)_G = \eta_G|_{V_j}$, we have constructed an equivariant bundle gerbe with connection, with equivariant curvature-form η_G .

REMARK 5.6. For $G = \mathrm{SU}(d+1)$ this construction reduces to the construction in terms of transition line bundles: All L_i , t_i , E_{ij} , u_{ijk} are trivial in this case, hence the entire information on the gerbe resides in the functions $s_{ij}: (X_{ij})^{[2]} \to \mathrm{U}(1)$ defined by the differences $\mu_j - \mu_i$. The condition $\delta s_{ij} = 1$ for these functions means that s_{ij} defines a line bundle L_{ij} over V_{ij} , as remarked at the beginning of Section 2.2. The condition $s_{ij}s_{jk}s_{ki} = 1$ over X_{ijk} is the compatibility condition over triple intersections.

6. Pre-quantization of conjugacy classes

It is a well-known fact from symplectic geometry that a coadjoint orbit $\mathcal{O}=G.\mu$ through $\mu\in\mathfrak{t}_+^*$ has integral symplectic form, i.e. admits a prequantum line bundle, if and only if μ is in the weight lattice Λ^* . The analogous question for conjugacy classes reads: For which $\mu\in\mathfrak{A}$ and $m\in\mathbf{N}$ does the pull-back of the mth power of the basic gerbe \mathcal{G}^m to the conjugacy class $\mathcal{C}=G.\exp(\mu)$ admit a pseudo-line bundle, with $m\omega_{\mathcal{C}}$ as its error 2-form? For any positive integer m>0 let

$$\Lambda_m^* = \Lambda^* \cap m\mathfrak{A}$$

be the set of level m weights. As is well-known [26], the set Λ_m^* parametrizes the positive energy representations of the loop group LG at level m.

THEOREM 6.1. The restriction of \mathcal{G}^m to a conjugacy class \mathcal{C} admits a pseudo-line bundle \mathcal{L} with connection, with error 2-form $m\omega_{\mathcal{C}}$, if and only if $\mathcal{C} = G \cdot \exp(\mu/m)$ with $\mu \in \Lambda_m^*$. Moreover \mathcal{L} has an equivariant extension in this case, with $m\omega_{\mathcal{C}}$ as its equivariant error 2-form.