

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE BASIC GERBE OVER A COMPACT SIMPLE LIE GROUP
Autor: Meinrenken, Eckhard
Kapitel: 6. Pre-quantization of conjugacy classes
DOI: <https://doi.org/10.5169/seals-66691>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

a distinguished, equivariant pseudo-line bundle (E_{ij}, s_{ij}) (where E_{ij} is trivial), with connection $\nabla^{E_{ij}}$ induced from the connection θ_{ij} . From the definition of θ_{ij} , it follows that the equivariant error 2-form for this connection is the pull-back of the equivariant symplectic form on the coadjoint orbit through $\mu_j - \mu_i$.

We now modify the bundle gerbe connection by adding the equivariant 2-form $(\varpi_j)_G \in \Omega_G^2(V_j)$ to the gerbe connection. Proposition 5.2(d) shows that the equivariant error 2-form of $\nabla^{E_{ij}}$ with respect to the new gerbe connection vanishes. The other conditions from the gluing construction in §4 are trivially satisfied. Since the equivariant 3-curvature for the new gerbe connection on \mathcal{G}_j is $d_G(\varpi_j)_G = \eta_G|_{V_j}$, we have constructed an equivariant bundle gerbe with connection, with equivariant curvature-form η_G .

REMARK 5.6. For $G = \mathrm{SU}(d+1)$ this construction reduces to the construction in terms of transition line bundles: All $L_i, t_i, E_{ij}, u_{ijk}$ are trivial in this case, hence the entire information on the gerbe resides in the functions $s_{ij} : (X_{ij})^{[2]} \rightarrow \mathrm{U}(1)$ defined by the differences $\mu_j - \mu_i$. The condition $\delta s_{ij} = 1$ for these functions means that s_{ij} defines a line bundle L_{ij} over V_{ij} , as remarked at the beginning of Section 2.2. The condition $s_{ij}s_{jk}s_{ki} = 1$ over X_{ijk} is the compatibility condition over triple intersections.

6. PRE-QUANTIZATION OF CONJUGACY CLASSES

It is a well-known fact from symplectic geometry that a coadjoint orbit $\mathcal{O} = G \cdot \mu$ through $\mu \in \mathfrak{t}_+^*$ has integral symplectic form, i.e. admits a pre-quantum line bundle, if and only if μ is in the weight lattice Λ^* . The analogous question for conjugacy classes reads: For which $\mu \in \mathfrak{A}$ and $m \in \mathbf{N}$ does the pull-back of the m th power of the basic gerbe \mathcal{G}^m to the conjugacy class $\mathcal{C} = G \cdot \exp(\mu)$ admit a pseudo-line bundle, with $m\omega_{\mathcal{C}}$ as its error 2-form? For any positive integer $m > 0$ let

$$\Lambda_m^* = \Lambda^* \cap m\mathfrak{A}$$

be the set of level m weights. As is well-known [26], the set Λ_m^* parametrizes the positive energy representations of the loop group LG at level m .

THEOREM 6.1. *The restriction of \mathcal{G}^m to a conjugacy class \mathcal{C} admits a pseudo-line bundle \mathcal{L} with connection, with error 2-form $m\omega_{\mathcal{C}}$, if and only if $\mathcal{C} = G \cdot \exp(\mu/m)$ with $\mu \in \Lambda_m^*$. Moreover \mathcal{L} has an equivariant extension in this case, with $m\omega_{\mathcal{C}}$ as its equivariant error 2-form.*

Proof. Given a conjugacy class $\mathcal{C} \subset G$, let $\mu \in m\mathfrak{A}$ be the unique point with $g := \exp(\mu/m) \in \mathcal{C}$, and let $K = G_g$ so that $\mathcal{C} = G/K$. Pick an index j with $\mathcal{C} \subset V_j$, and let

$$\nu = m\Psi_j(g) = \mu - m\mu_j.$$

Then

$$G_\mu \subset K \subset G_\nu.$$

Let $\mathcal{O}_\mu, \mathcal{O}_\nu \subset \mathfrak{g}$ denote the adjoint orbits of μ, ν , and $(\omega_\mu)_G, (\omega_\nu)_G$ their equivariant symplectic forms. The pull-back $\iota_{\mathcal{C}}^* \mathcal{G}^m$ is the gerbe over G/K defined as in Section 3 by the homomorphism $\varrho \in \text{Hom}(\pi_1(K), \text{U}(1))$, given as a composition

$$\pi_1(K) \rightarrow \pi_1(G_j) \rightarrow \text{U}(1),$$

where the first map is push-forward under the inclusion $K \hookrightarrow G_j$, and the second map is the homomorphism defined by the element $m\mu_j \in \mathfrak{t}$ for G_j .

Suppose now that $\mu \in \Lambda_m^*$. Then $m\mu_j$ equals $-\nu$ up to a weight lattice vector, which means that ϱ is the image of $-\nu \in (\mathfrak{k}^*)^K$ in the exact sequence (3.2). Hence, Proposition 3.2 says that we obtain an equivariant pseudo-line bundle for $\iota_{\mathcal{C}}^* \mathcal{G}^m$, with equivariant error 2-form

$$\Psi_j^*(\omega_\nu)_G - m\iota_{\mathcal{C}}^*(\varpi_j)_G = m\omega_{\mathcal{C}}.$$

Here we have used part (b) of Proposition 5.2.

Conversely, suppose that $\mathcal{G}^m|_{\mathcal{C}}$ admits a pseudo-line bundle with error 2-form $m\omega_{\mathcal{C}}$. Consider the pull-back of \mathcal{G} under the exponential map $\exp: \mathfrak{g} \rightarrow G$. The pull-back $\exp^* \eta \in \Omega^3(\mathfrak{g})$ is exact, and the homotopy operator for the linear retraction of \mathfrak{g} to the origin defines a 2-form $\varpi \in \Omega^2(\mathfrak{g})$ with $d\varpi = \exp^* \eta$. As in Proposition 5.2, one shows that for any adjoint orbit $\mathcal{O} \subset \mathfrak{g}$, with $\exp \mathcal{O} = \mathcal{C}$,

$$\iota_{\mathcal{O}}^* \varpi = \exp^* \omega_{\mathcal{C}} - \omega_{\mathcal{O}}$$

where $\omega_{\mathcal{O}}$ is the symplectic form on \mathcal{O} . In particular this applies to $\mathcal{O} = \mathcal{O}_{\mu/m}$. Choose a pseudo-line bundle for $\exp^* \mathcal{G}$ with error 2-form $-\varpi$. We then have two pseudo-line bundles for $\exp^* \mathcal{G}^m|_{\mathcal{O}}$ obtained by restricting the m th power of the pseudo-line bundle for $\exp^* \mathcal{G}$ or by pulling back the pseudo-line bundle for \mathcal{C} . Their quotient is a line bundle over \mathcal{O} , with curvature the difference of the error 2-forms:

$$m(\exp^* \omega_{\mathcal{C}} - \iota_{\mathcal{O}_{\mu}}^* \varpi) = m\omega_{\mathcal{O}}.$$

Thus $m(\mu/m) = \mu$ must be in the weight lattice.

REMARK 6.2. Z. Shahbazi has proved that if \mathcal{G} is a gerbe with connection over a manifold M , with curvature 3-form η , and $\Phi: N \rightarrow M$ is a map with $\Phi^*\eta + d\omega = 0$, then the pull-back gerbe $\Phi^*\mathcal{G}$ admits a pseudo-line bundle, with ω as its error 2-form, if and only if the pair (η, ω) defines an integral element of the relative de Rham cohomology $H^3(\Phi, \mathbf{R})$. This means that for any smooth 2-cycle $S \subset N$, and any smooth 3-chain $B \subset M$ with boundary $\Phi(S)$, one must have $\int_B \eta - \int_S \omega \in \mathbf{Z}$. The particular case where the target of Φ is a Lie group G is relevant for the pre-quantization of group-valued moment maps [1].

APPENDIX A. PROOF OF LEMMA 4.4

In this Appendix we prove Lemma 4.4, concerning the construction of a certain cover U_I of M from a given cover V_j . Write $M = \coprod_I A_I$ where

$$A_I = \bigcap_{i \in I} V_i \setminus \bigcup_{j \notin I} V_j.$$

Notice that $\bar{A}_I \subset \bigcup_{J \subset I} A_J$. By induction on the cardinality $k = |I|$ we will construct open sets $U_J \subset V_J$, having the following properties:

- (a) the closure \bar{U}_I does not meet \bar{U}_J for $|J| \leq |I|$ unless $J \subset I$,
- (b) each \bar{A}_I is contained in the union of U_J with $J \subset I$.

The induction starts at $k = 0$, taking $U_\emptyset = \emptyset$. Suppose we have constructed open sets U_I with $\bar{U}_I \subset V_I$ for $|I| < k$, such that the properties (a), (b) hold for all $|I| < k$. For $|I| = k$ consider the subsets

$$B_I := A_I \setminus \left(\bigcup_{J \subset I, |J| < k} U_J \right).$$

Note that (unlike A_I) the set B_I is closed. B_I does not meet \bar{A}_J unless $I \subset J$, and it also does not meet \bar{U}_J for $|J| < k$ unless $J \subset I$. That is, B_I is disjoint from

$$C_I := \bigcup_{J \not\subset I, |J| < k} \bar{U}_J \cup \bigcup_{K \not\supset I} \bar{A}_K.$$

Choose open sets U_I for $|I| = k$ with $B_I \subset U_I \subset \bar{U}_I \subset M \setminus C_I$, and such that the closures of the sets U_I for distinct I with $|I| = k$ are disjoint. The new collection of subsets will satisfy the properties (a), (b) for $|I| \leq k$. We next show that $V'_i = M \setminus \bigcup_{J \not\supset i} \bar{U}_J$ is a cover of M . Write $M = \coprod_I D_I$ with $D_I = \bar{U}_I \setminus \bigcup_{|J| < |I|} \bar{U}_J$. Then $D_I \cap \bar{U}_J = \emptyset$ unless $I \subset J$, so D_I is contained