

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 1.1 Définition of quasi-invariants
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

proofs are postponed until Lecture 3). In Lecture 2, we explain the origin of the ring of quasi-invariants in the theory of integrable systems, and introduce some tools from integrable systems, such as the Baker-Akhieser function. Finally, in Lecture 3, we develop the theory of the rational Cherednik algebra, the representation-theoretic techniques due to Opdam and Rouquier, and finish the proofs of the geometric statements from Chapter 1.

1. LECTURE 1

1.1 DEFINITION OF QUASI-INVARIANTS

In this lecture we will define the ring of quasi-invariants Q_m and discuss its main properties.

We will work over the field \mathbf{C} of complex numbers. Let W be a finite Coxeter group, i.e. a finite group generated by reflections. Let us denote by \mathfrak{h} its reflection representation. A typical example is the Weyl group of a semisimple Lie algebra acting on a Cartan subalgebra \mathfrak{h} . In the case the Lie algebra is $\mathfrak{sl}(n)$, we have that W is the symmetric group S_n on n letters and \mathfrak{h} is the space of diagonal traceless $n \times n$ matrices.

Let $\Sigma \subset W$ denote the set of reflections. Clearly, W acts on Σ by conjugation. Let $m: \Sigma \rightarrow \mathbf{Z}_+$ be a function on Σ taking non negative integer values, which is W -invariant. The number of orbits of W on Σ is generally very small. For example, if W is the Weyl group of a simple Lie algebra of ADE type, then W acts transitively on Σ , so m is a constant function.

For each reflection $s \in \Sigma$, choose $\alpha_s \in \mathfrak{h}^* - \{0\}$ so that, for $x \in \mathfrak{h}$, $\alpha_s(sx) = -\alpha_s(x)$ (this means that the hyperplane given by the equation $\alpha_s = 0$ is the reflection hyperplane for s).

DEFINITION 1.1 ([CV1, CV2]). A polynomial $q \in \mathbf{C}[\mathfrak{h}]$ is said to be *m -quasi-invariant* with respect to W if, for any $s \in \Sigma$, the polynomial $q(x) - q(sx)$ is divisible by $\alpha_s(x)^{2m_s+1}$.

We will denote by Q_m the space of m -quasi-invariant polynomials with respect to W .

Notice that every element of $\mathbf{C}[\mathfrak{h}]$ is a 0-quasi-invariant, and that every W -invariant is an m -quasi-invariant for any m . Indeed if $q \in \mathbf{C}[\mathfrak{h}]^W$, then we have $q(x) - q(sx) = 0$ for all $s \in \Sigma$, and 0 is divisible by all powers of $\alpha_s(x)$. Thus in a way, $\mathbf{C}[\mathfrak{h}]^W$ can be viewed as the set of ∞ -quasi-invariants.

EXAMPLE 1.2. The group $W = \mathbf{Z}/2$ acts on $\mathfrak{h} = \mathbf{C}$ by $s(v) = -v$. In this case m is a non negative integer and $\Sigma = \{s\}$. So this definition says that q is in Q_m iff $q(x) - q(-x)$ is divisible by x^{2m+1} . It is very easy to write a basis of Q_m . It is given by the polynomials $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$.

1.2 ELEMENTARY PROPERTIES OF Q_m

Some elementary properties of Q_m are collected in the following proposition.

PROPOSITION 1.3 (see [FV] and references therein).

- 1) $\mathbf{C}[\mathfrak{h}]^W \subset Q_m \subseteq \mathbf{C}[\mathfrak{h}]$, $Q_0 = \mathbf{C}[\mathfrak{h}]$, $Q_m \subset Q_{m'}$ if $m \geq m'$, $\bigcap_m Q_m = \mathbf{C}[\mathfrak{h}]^W$.
- 2) Q_m is a graded subalgebra of $\mathbf{C}[\mathfrak{h}]$.
- 3) The fraction field of Q_m is equal to $\mathbf{C}(\mathfrak{h})$.
- 4) Q_m is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and a finitely generated algebra. $\mathbf{C}[\mathfrak{h}]$ is a finite Q_m -module.

Proof. 1) is immediate and has already been mentioned in 1.1.

2) Clearly Q_m is closed under addition. Let $p, q \in Q_m$. Let $s \in \Sigma$. Then

$$p(x)q(x) - p(sx)q(sx) = (p(x) - p(sx))q(x) + p(sx)(q(x) - q(sx)).$$

Since both $p(x) - p(sx)$ and $q(x) - q(sx)$ are divisible by $\alpha_s^{2m_s+1}$, we deduce that $p(x)q(x) - p(sx)q(sx)$ is also divisible by $\alpha_s^{2m_s+1}$, proving the claim.

3) Consider the polynomial

$$\delta_{2m+1}(x) = \prod_{s \in \Sigma} \alpha_s(x)^{2m_s+1}.$$

This polynomial is uniquely defined up to scaling. One has $\delta_{2m+1}(sx) = -\delta_{2m+1}(x)$ for each $s \in \Sigma$, hence $\delta_{2m+1} \in Q_m$. Take $f(x) \in \mathbf{C}[\mathfrak{h}]$. We claim that $f(x)\delta_{2m+1}(x) \in Q_m$. As a matter of fact,

$$f(x)\delta_{2m+1}(x) - f(sx)\delta_{2m+1}(sx) = (f(x) + f(sx))\delta_{2m+1}(x),$$

and by its definition $\delta_{2m+1}(x)$ is divisible by $\alpha_s(x)^{2m_s+1}$ for all $s \in \Sigma$. This implies 3).

4) By Hilbert's theorem on the finiteness of invariants, we get that $\mathbf{C}[\mathfrak{h}]^W$ is a finitely generated algebra over \mathbf{C} and $\mathbf{C}[\mathfrak{h}]$ is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and hence a finite Q_m -module, proving the second part of 4).

Now $Q_m \subset \mathbf{C}[\mathfrak{h}]$ is a submodule of the finite module $\mathbf{C}[\mathfrak{h}]$ over the Noetherian ring $\mathbf{C}[\mathfrak{h}]^W$. Hence it is finite. This immediately implies that Q_m is a finitely generated algebra over \mathbf{C} . \square