

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	49 (2003)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
<b>Autor:</b>	Etingof, Pavel / Strickland, Elisabetta
<b>Kapitel:</b>	1.4 FURTHER PROPERTIES OF $X_m$
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-66677">https://doi.org/10.5169/seals-66677</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

$W_z$ -invariant, we deduce that  $p(x) - p(sx) = 0$ , so that in this case  $p(x) - p(sx)$  also is divisible by  $\alpha_s(x)^{2m_s+1}$ .

To conclude, notice that  $p(z) \neq 0$ . Indeed, for a reflection  $s$ ,  $\alpha_s$  vanishes exactly on the fixed points of  $s$ , so that  $\prod_{s \in \Sigma, sz \neq z} \alpha_s(z)^{2m_s+1} \neq 0$ . Also for all  $w \in W_z$ ,  $f(wz) = f(z) \neq 0$ . On the other hand, it is clear that  $p(y) = 0$ .  $\square$

EXAMPLE 1.5. Take  $W = \mathbf{Z}/2$ . As we have already seen,  $Q_m$  has a basis given by the monomials  $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$ . From this we deduce that setting  $z = x^2$  and  $y = x^{2m+1}$ ,  $Q_m = \mathbf{C}[y, z]/(y^2 - z^{2m+1}) = \mathbf{C}[K]$ , where  $K$  is the plane curve with a cusp at the origin, given by the equation  $y^2 = z^{2m+1}$ . The map  $\pi: \mathbf{C} \rightarrow K$  is given by  $\pi(t) = (t^{2m+1}, t^2)$ , which is clearly bijective.

#### 1.4 FURTHER PROPERTIES OF $X_m$

Let us get to some deeper properties of quasi-invariants. Let  $X$  be an irreducible affine variety over  $\mathbf{C}$  and  $A = \mathbf{C}[X]$ . Recall that, by the Noether Normalization Lemma, there exist  $f_1, \dots, f_n \in \mathbf{C}[X]$  which are algebraically independent over  $\mathbf{C}$  and such that  $\mathbf{C}[X]$  is a finite module over the polynomial ring  $\mathbf{C}[f_1, \dots, f_n]$ . This means that we have a finite morphism of  $X$  onto an affine space.

DEFINITION 1.6.  $A$  (and  $X$ ) is said to be *Cohen-Macaulay* if there exist  $f_1, \dots, f_n$  as above, with the property that  $\mathbf{C}[X]$  is a locally free module over  $\mathbf{C}[f_1, \dots, f_n]$ . (Notice that by the Quillen-Suslin theorem, this is equivalent to saying that  $A$  is a free module.)

REMARK. If  $A$  is Cohen-Macaulay, then for any  $f_1, \dots, f_n$  which are algebraically independent over  $\mathbf{C}$  and such that  $A$  is a finite module over the polynomial ring  $\mathbf{C}[f_1, \dots, f_n]$ , we have that  $A$  is a locally free  $\mathbf{C}[f_1, \dots, f_n]$ -module, see [Eis], Corollary 18.17.

THEOREM 1.7 ([EG2], [BEG], conjectured in [FV]).  $Q_m$  is Cohen-Macaulay.

Notice that, using Chevalley's result that  $\mathbf{C}[\mathfrak{h}]^W$  is a polynomial ring, it will suffice, in order to prove Theorem 1.7, to prove:

THEOREM 1.8 ([EG2, BEG], conjectured in [FV]).  $Q_m$  is a free  $\mathbf{C}[\mathfrak{h}]^W$ -module.

We show how one can prove this Theorem in 3.10. This proof follows [BEG] (the original proof of [EG2] is shorter but somewhat less conceptual). The main idea of the proof is to show that the  $\mathbf{C}[\mathfrak{h}]^W$ -module  $Q_m$  can be extended to a module over a bigger (noncommutative) algebra, namely the spherical subalgebra of the rational Cherednik algebra. Furthermore, this module belongs to an appropriate category of representations of this algebra, called category  $\mathcal{O}$ . On the other hand, it can be shown that any module over the spherical subalgebra that belongs to this category is free when restricted to the commutative algebra  $\mathbf{C}[\mathfrak{h}]^W$ .

### 1.5 THE POINCARÉ SERIES OF $Q_m$

Consider now the Poincaré series

$$h_{Q_m}(t) = \sum_{r \geq 0} \dim Q_m[r] t^r,$$

where  $Q_m[r]$  denotes the graded component of  $Q_m$  of degree  $r$ . For every irreducible representation  $\tau \in \widehat{W}$ , define

$$\chi_\tau(t) = \sum_{r \geq 0} \dim \text{Hom}_W(\tau, \mathbf{C}[\mathfrak{h}][r]) t^r.$$

Consider the element in the group ring  $\mathbf{Z}[W]$

$$\mu_m = \sum_{s \in \Sigma} m_s (1 - s).$$

The  $W$ -invariance of  $m$  implies that  $\mu_m$  lies in the center of  $\mathbf{Z}[W]$ . Hence it is clear that  $\mu_m$  acts as a scalar,  $\xi_m(\tau)$ , on  $\tau$ . Let  $d_\tau$  be the degree of  $\tau$ .

**LEMMA 1.9.** *The scalar  $\xi_m(\tau)$  is an integer.*

*Proof.*  $\mathbf{Z}[W]$  and hence also its center, is a finite  $\mathbf{Z}$ -module. This clearly implies that  $\xi_m(\tau)$  is an algebraic integer. Thus to prove that  $\xi_m(\tau)$  is an integer, it suffices to see that  $\xi_m(\tau)$  is a rational number. Let  $d_{\tau,s}$  be the dimension of the space of  $s$ -invariants in  $\tau$ . Taking traces we get

$$d_\tau \xi_m(\tau) = \sum_{s \in \Sigma} 2m_s (d_\tau - d_{\tau,s}),$$

which gives the rationality of  $\xi_m(\tau)$ .  $\square$