1.5 The Poincaré séries of Q_m

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.
Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.
Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*
ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

http://www.e-periodica.ch

We show how one can prove this Theorem in 3.10. This proof follows [BEG] (the original proof of [EG2] is shorter but somewhat less conceptual). The main idea of the proof is to show that the $\mathbf{C}[\mathfrak{h}]^W$-module Q_m can be extended to a module over a bigger (noncommutative) algebra, namely the spherical subalgebra of the rational Cherednik algebra. Furthermore, this module belongs to an appropriate category of representations of this algebra, called category $\mathcal{O}$. On the other hand, it can be shown that any module over the spherical subalgebra that belongs to this category is free when restricted to the commutative algebra $\mathbf{C}[\mathfrak{h}]^W$.

1.5 The Poincaré series of Q_m

Consider now the Poincaré series

$$h_{Q_m}(t) = \sum_{r \geq 0} \dim Q_m[r] t^r ,$$

where $Q_m[r]$ denotes the graded component of Q_m of degree r. For every irreducible representation $\tau \in \widehat{W}$, define

$$\chi_\tau(t) = \sum_{r \geq 0} \dim \mathrm{Hom}_W(\tau, \mathbf{C}[\mathfrak{h}][r]) t^r .$$

Consider the element in the group ring $\mathbf{Z}[W]$

$$\mu_m = \sum_{s \in \Sigma} m_s (1 - s) .$$

The W-invariance of m implies that μ_m lies in the center of $\mathbf{Z}[W]$. Hence it is clear that μ_m acts as a scalar, $\xi_m(\tau)$, on τ. Let d_τ be the degree of τ.

LEMMA 1.9. *The scalar $\xi_m(\tau)$ is an integer.*

Proof. $\mathbf{Z}[W]$ and hence also its center, is a finite $\mathbf{Z}$-module. This clearly implies that $\xi_m(\tau)$ is an algebraic integer. Thus to prove that $\xi_m(\tau)$ is an integer, it suffices to see that $\xi_m(\tau)$ is a rational number. Let $d_{\tau,s}$ be the dimension of the space of s-invariants in τ. Taking traces we get

$$d_\tau \xi_m(\tau) = \sum_{s \in \Sigma} 2 m_s (d_\tau - d_{\tau,s}) ,$$

which gives the rationality of $\xi_m(\tau)$. □

THEOREM 1.10. *One has*

$$h_{Q_m}(t) = \sum_{\tau \in \widehat{W}} d_\tau t^{\xi_m(\tau)} \chi_\tau(t) . \tag{1}$$

REMARK. This theorem was proved in [FeV] modulo Theorem 1.7 (conjectured in [FV]) using the so-called Matsuo-Cherednik correspondence (see [FeV] for details). Thus, Theorem 1.10 follows from [FeV] and [EG2]. Another proof of this theorem is given in [BEG]; this is the proof we will discuss below (in Lecture 3).

EXAMPLE 1.11. If $m = 0$, since $Q_0 = \mathbf{C}[\mathfrak{h}]$, the theorem says that

$$h_{Q_0}(t) = \frac{1}{(1-t)^n} = \sum_{\tau \in \widehat{W}} d_\tau \chi_\tau(t) .$$

Indeed, as a W-module one has

$$\mathbf{C}[\mathfrak{h}] = \oplus_\tau \tau \otimes \mathrm{Hom}_W(\tau, \mathbf{C}[\mathfrak{h}]) .$$

EXAMPLE 1.12. If $W = \mathbf{Z}/2$, then $\widehat{W} = \{+, -\}$, where $+$ (respectively $-$) denotes the trivial (respectively the sign) representation. One has

$$\mathbf{C}[x] = \mathbf{C}[x^2] \oplus \mathbf{C}[x^2]x ,$$

where $\mathbf{C}[x^2] = \mathbf{C}[x]^W$ and $\mathbf{C}[x^2]x$ is the isotypic component of the sign representation. Thus

$$\chi_+(t) = \frac{1}{1-t^2} , \quad \chi_-(t) = \frac{t}{1-t^2} ,$$

$\mu_m = m(1-s)$. Thus $\xi_m(+) = 0$, $\xi_m(-) = 2m$. We deduce that

$$h_{Q_m}(t) = \frac{1}{1-t^2} + \frac{t^{2m+1}}{1-t^2} ,$$

as we already know.

Recall now that as a graded W-module $\mathbf{C}[\mathfrak{h}]$ is isomorphic to $\mathbf{C}[\mathfrak{h}]^W \otimes H$, H being the space of harmonic polynomials. We deduce that the τ-isotypic component in $\mathbf{C}[\mathfrak{h}]$ is isomorphic to $\mathbf{C}[\mathfrak{h}]^W \otimes H_\tau$.

Set $K_\tau(t) = \sum_{r\geq 0} \dim \mathrm{Hom}_W(\tau, H[r])t^r$. This is a polynomial, called the Kostka polynomial relative to τ. We deduce that

$$\chi_\tau(t) = \frac{K_\tau(t)}{\prod_{i=1}^n (1-t^{d_i})}. \tag{2}$$

Also, if $\tau' = \tau \otimes \varepsilon$, ε being the sign representation, one has

$$K_{\tau'}(t) = K_\tau(t^{-1})t^{|\Sigma|}.$$

Set now

$$P_m(t) = \sum_{\tau\in\widehat{W}} d_\tau t^{\xi_m(\tau)} K_\tau(t).$$

We have

PROPOSITION 1.13 ([FeV]).

$$h_{Q_m}(t) = \frac{P_m(t)}{\prod_{i=1}^n (1-t^{d_i})}.$$

Furthermore $P_m(t) = t^{\xi_m(\varepsilon)+|\Sigma|}P_m(t^{-1})$.

Proof. Substituting the expression (2) for $\chi_\tau(t)$ in (1.10) and using the definition of $P_m(t)$, we get

$$h_{Q_m}(t) = \sum_{\tau\in\widehat{W}} d_\tau t^{\xi_m(\tau)} \frac{K_\tau(t)}{\prod_{i=1}^n (1-t^{d_i})} = \frac{P_m(t)}{\prod_{i=1}^n (1-t^{d_i})},$$

as desired.

Now notice that

$$\xi_m(\tau) + \xi_m(\tau') = \sum_{s\in\Sigma} 2m_s = \xi_m(\varepsilon).$$

Using this we get

$$\begin{aligned} t^{\xi_m(\varepsilon)+|\Sigma|}P_m(t^{-1}) &= \sum_{\tau\in\widehat{W}} d_\tau t^{\xi_m(\varepsilon)-\xi_m(\tau)}t^{|\Sigma|}K_\tau(t^{-1}) \\ &= \sum_{\tau'\in\widehat{W}} d_{\tau'} t^{\xi_m(\tau')}K_{\tau'}(t) = P_m(t), \end{aligned}$$

as desired. □

From this we deduce

THEOREM 1.14 ([EG2, BEG, FeV], conjectured in [FV]). *The ring Q_m of m-quasi-invariants is Gorenstein.*

Proof. By Stanley's theorem (see [Eis]), a positively graded Cohen-Macaulay domain A is Gorenstein iff its Poincaré series is a rational function $h(t)$ satisfying the equation $h(t^{-1}) = (-1)^n t^l h(t)$, where l is an integer and n is the dimension of the spectrum of A. Thus the result follows immediately from Proposition 1.13. □

1.6 THE RING OF DIFFERENTIAL OPERATORS ON X_m

Finally, let us introduce the ring $\mathcal{D}(X_m)$ of differential operators on X_m, that is the ring of differential operators with coefficients in $\mathbf{C}(\hbar)$ mapping Q_m to Q_m. It is clear that this definition coincides with Grothendieck's well-known definition ([Bj]).

THEOREM 1.15 ([BEG]). *$\mathcal{D}(X_m)$ is a simple algebra.*

REMARK 1.16. a) The ring of differential operators on a smooth affine algebraic variety is always simple (see [Bj], Chapter 3).

b) By a result of M. van den Bergh [VdB], for a non-smooth variety, the simplicity of the ring of differential operators implies the Cohen-Macaulay property of this variety.

2. LECTURE 2

We will now see how the ring Q_m appears in the theory of completely integrable systems.

2.1 HAMILTONIAN MECHANICS AND INTEGRABLE SYSTEMS

Recall the basic setup of Hamiltonian mechanics [Ar]. Consider a mechanical system with configuration space X (a smooth manifold). Then the phase space of this system is T^*X, the cotangent bundle on X. The space T^*X is naturally a symplectic manifold, and in particular we have an operation of Poisson bracket on functions on T^*X. A point of T^*X is a pair (x,p), where $x \in X$ is the position and $p \in T_x^*X$ is the momentum. Such pairs are