Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 49 (2003)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA

Autor: Etingof, Pavel / Strickland, Elisabetta

Kapitel: 2.1 Hamiltonian mechanics and integrable Systems

DOI: https://doi.org/10.5169/seals-66677

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

From this we deduce

THEOREM 1.14 ([EG2, BEG, FeV], conjectured in [FV]). *The ring Q_m of m-quasi-invariants is Gorenstein.*

Proof. By Stanley's theorem (see [Eis]), a positively graded Cohen-Macaulay domain A is Gorenstein iff its Poincaré series is a rational function $h(t)$ satisfying the equation $h(t^{-1}) = (-1)^n t^l h(t)$, where l is an integer and n is the dimension of the spectrum of A. Thus the result follows immediately from Proposition 1.13. □

1.6 THE RING OF DIFFERENTIAL OPERATORS ON X_m

Finally, let us introduce the ring $\mathcal{D}(X_m)$ of differential operators on X_m, that is the ring of differential operators with coefficients in $\mathbf{C}(\hbar)$ mapping Q_m to Q_m. It is clear that this definition coincides with Grothendieck's well-known definition ([Bj]).

THEOREM 1.15 ([BEG]). *$\mathcal{D}(X_m)$ is a simple algebra.*

REMARK 1.16. a) The ring of differential operators on a smooth affine algebraic variety is always simple (see [Bj], Chapter 3).

b) By a result of M. van den Bergh [VdB], for a non-smooth variety, the simplicity of the ring of differential operators implies the Cohen-Macaulay property of this variety.

2. LECTURE 2

We will now see how the ring Q_m appears in the theory of completely integrable systems.

2.1 HAMILTONIAN MECHANICS AND INTEGRABLE SYSTEMS

Recall the basic setup of Hamiltonian mechanics [Ar]. Consider a mechanical system with configuration space X (a smooth manifold). Then the phase space of this system is T^*X, the cotangent bundle on X. The space T^*X is naturally a symplectic manifold, and in particular we have an operation of Poisson bracket on functions on T^*X. A point of T^*X is a pair (x,p), where $x \in X$ is the position and $p \in T_x^*X$ is the momentum. Such pairs are

called states of the system. The dynamics of the system $x = x(t)$, $p = p(t)$ depends on the Hamiltonian, or energy function, $E(x,p)$ on T^*X. Given E and the initial state $x(0)$, $p(0)$, one can recover the dynamics $x = x(t)$, $p = p(t)$ from Hamilton's differential equations $\frac{df(x,p)}{dt} = \{f,E\}$. If X is locally identified with $\mathbf{R}^n$ by choosing coordinates $x_1,\dots,x_n$, then T^*X is locally identified with $\mathbf{R}^{2n}$ with coordinates $x_1,\dots,x_n,p_1,\dots,p_n$. In these coordinates, Hamilton's equations may be written in their standard form

$$\dot{x}_i = \frac{\partial E}{\partial p_i}, \qquad \dot{p}_i = -\frac{\partial E}{\partial x_i}.$$

A function $I(x,p)$ is called an integral of motion for our system if $\{I,E\} = 0$. Integrals of motion are useful, since for any such integral I the function $I(x(t),p(t))$ is constant, which allows one to reduce the number of variables by 2. Thus, if we are given n functionally independent integrals of motion $I_1,\dots,I_n$ with $\{I_l,I_k\} = 0$ for all $1 \le l,k \le n$, then all $2n$ variables x_i,p_i can be excluded, and the system can be completely solved by quadratures. Such a situation is called complete (or Liouville) integrability.

2.2 The classical Calogero-Moser system

Quasi-invariants are related to many-particle systems. Consider a system of n particles on the real line $\mathbf{R}$. A potential is an even function

$$U(x) = U(-x), \quad x \in \mathbf{R}.$$

Two particles at points a, b have energy of interaction $U(a-b)$. The total energy of our system of particles is

$$E = \sum_{i=1}^{n} \frac{p_i^2}{2} + \sum_{i<j} U(x_i - x_j).$$

Here, x_i are the coordinates of the particles, p_i their momenta. The dynamics of the particles $x_i = x_i(t)$, $p_i = p_i(t)$ is governed by the Hamilton equations with energy function E.

This is a system of nonlinear differential equations, which in general can be difficult to solve explicitly. However, for special potentials this system might be completely integrable. For instance, we will see that this is the case for the Calogero-Moser potential,

$$U(x) = \frac{\gamma}{x^2},$$

γ being a constant.