

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 2.5 Dunkl operators and symmetric quantum integrals
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

In general, m is not a homomorphism. However:

PROPOSITION 2.4. *Let $\mathcal{A}^W \subset \mathcal{A}$ denote the subalgebra of elements invariant under conjugation by W . Then the restriction of m to \mathcal{A}^W is an algebra homomorphism.*

Proof. If $A \in \mathcal{A}^W$, then clearly $m(A)$ is W -invariant. Now if we take $A, B \in \mathcal{A}^W$ and f a W -invariant function we have that $B(f)$ is also W -invariant. So

$$m(AB)(f) = (AB)(f) = A(B(f)) = A(m(B)(f)) = m(A)(m(B)(f)).$$

Thus $m(AB)$ and $m(A)m(B)$ coincide on W -invariant functions and hence coincide. \square

2.5 DUNKL OPERATORS AND SYMMETRIC QUANTUM INTEGRALS

In this subsection we will construct quantum integrals of the Calogero-Moser operator. This construction is due to Heckman [He] and is based on the Dunkl operators, introduced in [Du].

Fix a W -invariant function $c: \Sigma \rightarrow \mathbf{C}$ such that $\beta_s = c_s(c_s + 1)$ for each $s \in \Sigma$. Set $\delta_c := \prod_{s \in \Sigma} \alpha_s(x)^{c_s}$ and define

$$L = \delta_c(x)H\delta_c(x)^{-1}.$$

Then an easy computation shows that

$$L = \Delta - \sum_{s \in \Sigma} \frac{2c_s}{\alpha_s(x)} \partial_{\alpha_s},$$

where, for a vector $y \in \mathfrak{h}$, the symbol ∂_y denotes, as usual, the partial derivative in the y direction (notice that using the scalar product we are viewing α_s as a vector in \mathfrak{h} orthogonal to the hyperplane fixed by s).

From now on we will work with L instead of H and study the eigenvalue problem

$$(4) \quad L\psi = \lambda\psi.$$

It is clear that ψ is a solution of this equation if and only if $\delta_c(x)^{-1}\psi$ is a solution of (3).

Since for any $s \in \Sigma$ and $f \in \mathbf{C}[\mathfrak{h}]$ we have that $f(sx) - f(x)$ is divisible by $\alpha_s(x)$, the operator

$$\frac{1}{\alpha_s(x)}(s - 1) \in \mathcal{A}$$

maps $\mathbf{C}[\mathfrak{h}]$ to itself.

DEFINITION 2.5. Given $y \in \mathfrak{h}$, we define the *Dunkl operator* D_y on $\mathbf{C}[\mathfrak{h}]$ by

$$D_y := \partial_y + \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)}{\alpha_s(x)} (s - 1).$$

We have the following very important theorem.

THEOREM 2.6 ([Du]). *Let $y, z \in \mathfrak{h}$. Then*

$$[D_y, D_z] = 0.$$

Proof. See [Du], [Op]. \square

PROPOSITION 2.7 (Heckman [He]). *Let $\{y_1, \dots, y_n\}$ be an orthonormal basis of \mathfrak{h} . Then we have*

$$m\left(\sum_{i=1}^n D_{y_i}^2\right) = L.$$

Proof. Observe that $m(\sum_{i=1}^n D_{y_i}^2) = \sum_{i=1}^n m(D_{y_i}^2)$, so we need to compute $m(D_y^2)$ for $y \in \mathfrak{h}$. We have $m(D_y^2) = m(D_y m(D_y)) = m(D_y \partial_y)$. A simple computation shows that

$$D_y \partial_y = \partial_y^2 + \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)}{\alpha_s(x)} (\partial_y (s - 1) - \frac{2(\alpha_s, y)}{(\alpha_s, \alpha_s)} \partial_{\alpha_s}).$$

Thus

$$m(D_y^2) = \partial_y^2 - 2 \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)^2}{(\alpha_s, \alpha_s) \alpha_s(x)} \partial_{\alpha_s}.$$

We get

$$m\left(\sum_{i=1}^n D_{y_i}^2\right) = \sum_i \partial_{y_i}^2 - 2 \sum_{s \in \Sigma} c_s \frac{\sum_{i=1}^n (\alpha_s, y_i)^2}{(\alpha_s, \alpha_s) \alpha_s(x)} \partial_{\alpha_s} = L,$$

since $\sum_{i=1}^n (\alpha_s, y_i)^2 = (\alpha_s, \alpha_s)$. \square

We are now ready to give the construction of quantum integrals of L . Consider the symmetric algebra $S\mathfrak{h} = \mathbf{C}[y_1, \dots, y_n]$ which we can identify, using the fact that the Dunkl operators commute, with the polynomial ring $\mathbf{C}[D_{y_1}, \dots, D_{y_n}] \subset \mathcal{A}$. The restriction of m to $S\mathfrak{h}^W$ is an algebra homomorphism into the ring $\mathcal{D}(U)$ (and in fact into $\mathcal{D}(U/W)$). Since $S\mathfrak{h}^W$ is itself a polynomial ring $\mathbf{C}[q_1, \dots, q_n]$, with q_1, \dots, q_n of degree d_1, \dots, d_n ,

d_i being the degrees of basic W -invariants, we obtain a polynomial ring of commuting differential operators in $\mathcal{D}(U)$. Given $q \in \mathbf{C}[q_1, \dots, q_n]$ we will denote by L_q the corresponding differential operator. We may assume that $q_1 = \sum_{i=1}^n y_i^2$ so that $L = L_{q_1}$. Thus for every $q \in \mathbf{C}[q_1, \dots, q_n]$, L_q is a quantum integral of the quantum Calogero-Moser system. In particular, the operators L_{q_1}, \dots, L_{q_n} are n algebraically independent pairwise commuting quantum integrals.

Now the eigenvalue problem (4) may be replaced by

$$L_p \psi = \lambda_p \psi$$

for $p \in \mathbf{C}[q_1, \dots, q_n]$, where the assignment $p \rightarrow \lambda_p$ is an algebra homomorphism $\mathbf{C}[q_1, \dots, q_n] \rightarrow \mathbf{C}$.

In other words, we may say that since $\mathbf{C}[q_1, \dots, q_n] = \mathbf{C}[\mathfrak{h}^*/W] = \mathbf{C}[\mathfrak{h}/W]$, for every point $k \in \mathfrak{h}/W$, we have the eigenvalue problem

$$(5) \quad L_p \psi = p(k) \psi.$$

PROPOSITION 2.8. *Near a generic point $x_0 \in \mathfrak{h}$, the system $L_p \psi = p(k) \psi$ has a space of solutions of dimension $|W|$.*

Proof. The proposition follows easily from the fact that the symbols of L_{q_i} are $q_i(\partial)$, and that $\mathbf{C}[y_1, \dots, y_n]$ is a free module over $\mathbf{C}[q_1, \dots, q_n]$ of rank $|W|$. \square

2.6 ADDITIONAL INTEGRALS FOR INTEGER VALUED c

If $c_s \notin \mathbf{Z}$, the analysis of the solutions of the equations $L_p \psi = p(k) \psi$ is rather difficult (see [HO]). However, in the case $c: \Sigma \rightarrow \mathbf{Z}$, the system can be simplified. Let us consider this case. First remark that, since $\beta_s = c_s(c_s + 1)$, by changing c_s to $-1 - c_s$ if necessary, we may assume that c is non-negative. So we will assume that c takes non-negative integral values and we will denote it by m .

System (5) can be further simplified, if we can find a differential operator M (not a polynomial of L_{q_1}, \dots, L_{q_n}) such that $[M, L_p] = 0$ for all $p \in \mathbf{C}[q_1, \dots, q_n]$. Then the operator M will act on the space of solutions of (5), hopefully with distinct eigenvalues. So if μ is such an eigenvalue, the system

$$\begin{cases} L_p \psi = p(k) \psi \\ M \psi = \mu \psi \end{cases}$$