

2.5 Dunkl operators and symmetric quantum integrals

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

In general, m is not a homomorphism. However:

PROPOSITION 2.4. *Let $\mathcal{A}^W \subset \mathcal{A}$ denote the subalgebra of elements invariant under conjugation by W . Then the restriction of m to \mathcal{A}^W is an algebra homomorphism.*

Proof. If $A \in \mathcal{A}^W$, then clearly $m(A)$ is W -invariant. Now if we take $A, B \in \mathcal{A}^W$ and f a W -invariant function we have that $B(f)$ is also W -invariant. So

$$m(AB)(f) = (AB)(f) = A(B(f)) = A(m(B)(f)) = m(A)(m(B)(f)).$$

Thus $m(AB)$ and $m(A)m(B)$ coincide on W -invariant functions and hence coincide. \square

2.5 DUNKL OPERATORS AND SYMMETRIC QUANTUM INTEGRALS

In this subsection we will construct quantum integrals of the Calogero-Moser operator. This construction is due to Heckman [He] and is based on the Dunkl operators, introduced in [Du].

Fix a W -invariant function $c: \Sigma \rightarrow \mathbf{C}$ such that $\beta_s = c_s(c_s + 1)$ for each $s \in \Sigma$. Set $\delta_c := \prod_{s \in \Sigma} \alpha_s(x)^{c_s}$ and define

$$L = \delta_c(x) H \delta_c(x)^{-1}.$$

Then an easy computation shows that

$$L = \Delta - \sum_{s \in \Sigma} \frac{2c_s}{\alpha_s(x)} \partial_{\alpha_s},$$

where, for a vector $y \in \mathfrak{h}$, the symbol ∂_y denotes, as usual, the partial derivative in the y direction (notice that using the scalar product we are viewing α_s as a vector in \mathfrak{h} orthogonal to the hyperplane fixed by s).

From now on we will work with L instead of H and study the eigenvalue problem

$$(4) \quad L\psi = \lambda\psi.$$

It is clear that ψ is a solution of this equation if and only if $\delta_c(x)^{-1}\psi$ is a solution of (3).

Since for any $s \in \Sigma$ and $f \in \mathbf{C}[\mathfrak{h}]$ we have that $f(sx) - f(x)$ is divisible by $\alpha_s(x)$, the operator

$$\frac{1}{\alpha_s(x)}(s - 1) \in \mathcal{A}$$

maps $\mathbf{C}[\mathfrak{h}]$ to itself.

DEFINITION 2.5. Given $y \in \mathfrak{h}$, we define the *Dunkl operator* D_y on $\mathbf{C}[\mathfrak{h}]$ by

$$D_y := \partial_y + \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)}{\alpha_s(x)} (s - 1).$$

We have the following very important theorem.

THEOREM 2.6 ([Du]). *Let $y, z \in \mathfrak{h}$. Then*

$$[D_y, D_z] = 0.$$

Proof. See [Du], [Op]. \square

PROPOSITION 2.7 (Heckman [He]). *Let $\{y_1, \dots, y_n\}$ be an orthonormal basis of \mathfrak{h} . Then we have*

$$m\left(\sum_{i=1}^n D_{y_i}^2\right) = L.$$

Proof. Observe that $m(\sum_{i=1}^n D_{y_i}^2) = \sum_{i=1}^n m(D_{y_i}^2)$, so we need to compute $m(D_y^2)$ for $y \in \mathfrak{h}$. We have $m(D_y^2) = m(D_y m(D_y)) = m(D_y \partial_y)$. A simple computation shows that

$$D_y \partial_y = \partial_y^2 + \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)}{\alpha_s(x)} (\partial_y(s - 1) - \frac{2(\alpha_s, y)}{(\alpha_s, \alpha_s)} \partial_{\alpha_s} s).$$

Thus

$$m(D_y^2) = \partial_y^2 - 2 \sum_{s \in \Sigma} c_s \frac{(\alpha_s, y)^2}{(\alpha_s, \alpha_s) \alpha_s(x)} \partial_{\alpha_s}.$$

We get

$$m\left(\sum_{i=1}^n D_{y_i}^2\right) = \sum_i \partial_{y_i}^2 - 2 \sum_{s \in \Sigma} c_s \frac{\sum_{i=1}^n (\alpha_s, y_i)^2}{(\alpha_s, \alpha_s) \alpha_s(x)} \partial_{\alpha_s} = L,$$

since $\sum_{i=1}^n (\alpha_s, y_i)^2 = (\alpha_s, \alpha_s)$. \square

We are now ready to give the construction of quantum integrals of L . Consider the symmetric algebra $S\mathfrak{h} = \mathbf{C}[y_1, \dots, y_n]$ which we can identify, using the fact that the Dunkl operators commute, with the polynomial ring $\mathbf{C}[D_{y_1}, \dots, D_{y_n}] \subset \mathcal{A}$. The restriction of m to $S\mathfrak{h}^W$ is an algebra homomorphism into the ring $\mathcal{D}(U)$ (and in fact into $\mathcal{D}(U/W)$). Since $S\mathfrak{h}^W$ is itself a polynomial ring $\mathbf{C}[q_1, \dots, q_n]$, with q_1, \dots, q_n of degree d_1, \dots, d_n ,

d_i being the degrees of basic W -invariants, we obtain a polynomial ring of commuting differential operators in $\mathcal{D}(U)$. Given $q \in \mathbf{C}[q_1, \dots, q_n]$ we will denote by L_q the corresponding differential operator. We may assume that $q_1 = \sum_{i=1}^n y_i^2$ so that $L = L_{q_1}$. Thus for every $q \in \mathbf{C}[q_1, \dots, q_n]$, L_q is a quantum integral of the quantum Calogero-Moser system. In particular, the operators L_{q_1}, \dots, L_{q_n} are n algebraically independent pairwise commuting quantum integrals.

Now the eigenvalue problem (4) may be replaced by

$$L_p \psi = \lambda_p \psi$$

for $p \in \mathbf{C}[q_1, \dots, q_n]$, where the assignment $p \rightarrow \lambda_p$ is an algebra homomorphism $\mathbf{C}[q_1, \dots, q_n] \rightarrow \mathbf{C}$.

In other words, we may say that since $\mathbf{C}[q_1, \dots, q_n] = \mathbf{C}[\mathfrak{h}^*/W] = \mathbf{C}[\mathfrak{h}/W]$, for every point $k \in \mathfrak{h}/W$, we have the eigenvalue problem

$$(5) \quad L_p \psi = p(k) \psi.$$

PROPOSITION 2.8. *Near a generic point $x_0 \in \mathfrak{h}$, the system $L_p \psi = p(k) \psi$ has a space of solutions of dimension $|W|$.*

Proof. The proposition follows easily from the fact that the symbols of L_{q_i} are $q_i(\partial)$, and that $\mathbf{C}[y_1, \dots, y_n]$ is a free module over $\mathbf{C}[q_1, \dots, q_n]$ of rank $|W|$. \square

2.6 ADDITIONAL INTEGRALS FOR INTEGER VALUED c

If $c_s \notin \mathbf{Z}$, the analysis of the solutions of the equations $L_p \psi = p(k) \psi$ is rather difficult (see [HO]). However, in the case $c: \Sigma \rightarrow \mathbf{Z}$, the system can be simplified. Let us consider this case. First remark that, since $\beta_s = c_s(c_s+1)$, by changing c_s to $-1 - c_s$ if necessary, we may assume that c is non-negative. So we will assume that c takes non-negative integral values and we will denote it by m .

System (5) can be further simplified, if we can find a differential operator M (not a polynomial of L_{q_1}, \dots, L_{q_n}) such that $[M, L_p] = 0$ for all $p \in \mathbf{C}[q_1, \dots, q_n]$. Then the operator M will act on the space of solutions of (5), hopefully with distinct eigenvalues. So if μ is such an eigenvalue, the system

$$\begin{cases} L_p \psi = p(k) \psi \\ M\psi = \mu \psi \end{cases}$$