Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 49 (2003)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND

THE CHEREDNIK ALGEBRA

Autor: Etingof, Pavel / Strickland, Elisabetta

Kapitel: 2.6 Additional integrals for integer valued c

DOI: https://doi.org/10.5169/seals-66677

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

 d_i being the degrees of basic W-invariants, we obtain a polynomial ring of commuting differential operators in $\mathcal{D}(U)$. Given $q \in \mathbb{C}[q_1,\ldots,q_n]$ we will denote by L_q the corresponding differential operator. We may assume that $q_1 = \sum_{i=1}^n y_1^2$ so that $L = L_{q_1}$. Thus for every $q \in \mathbb{C}[q_1,\ldots,q_n]$, L_q is a quantum integral of the quantum Calogero-Moser system. In particular, the operators L_{q_1},\ldots,L_{q_n} are n algebraically independent pairwise commuting quantum integrals.

Now the eigenvalue problem (4) may be replaced by

$$L_p\psi=\lambda_p\psi$$

for $p \in \mathbb{C}[q_1, \dots, q_n]$, where the assignment $p \to \lambda_p$ is an algebra homomorphism $\mathbb{C}[q_1, \dots, q_n] \to \mathbb{C}$.

In other words, we may say that since $\mathbb{C}[q_1,\ldots,q_n] = \mathbb{C}[\mathfrak{h}^*/W] = \mathbb{C}[\mathfrak{h}/W]$, for every point $k \in \mathfrak{h}/W$, we have the eigenvalue problem

$$(5) L_p \psi = p(k)\psi.$$

PROPOSITION 2.8. Near a generic point $x_0 \in \mathfrak{h}$, the system $L_p \psi = p(k) \psi$ has a space of solutions of dimension |W|.

Proof. The proposition follows easily from the fact that the symbols of L_{q_i} are $q_i(\partial)$, and that $\mathbf{C}[y_1, \ldots, y_n]$ is a free module over $\mathbf{C}[q_1, \ldots, q_n]$ of rank |W|.

2.6 Additional integrals for integer valued c

If $c_s \notin \mathbb{Z}$, the analysis of the solutions of the equations $L_p \psi = p(k) \psi$ is rather difficult (see [HO]). However, in the case $c: \Sigma \to \mathbb{Z}$, the system can be simplified. Let us consider this case. First remark that, since $\beta_s = c_s(c_s+1)$, by changing c_s to $-1 - c_s$ if necessary, we may assume that c is non-negative. So we will assume that c takes non-negative integral values and we will denote it by m.

System (5) can be further simplified, if we can find a differential operator M (not a polynomial of L_{q_1}, \ldots, L_{q_n}) such that $[M, L_p] = 0$ for all $p \in \mathbb{C}[q_1, \ldots, q_n]$. Then the operator M will act on the space of solutions of (5), hopefully with distinct eigenvalues. So if μ is such an eigenvalue, the system

$$\begin{cases} L_p \psi = p(k) \, \psi \\ M \psi = \mu \, \psi \end{cases}$$

will have a one dimensional space of solutions and we can find the unique up to scaling solution ψ using Euler's formula.

Such an M exists if and only if c=m has integer values. Namely, we will see that one can extend the homomorphism $\mathbb{C}[q_1,\ldots,q_n]\to \mathcal{D}(U)$ mapping $q\to L_q$ to the ring of m-quasi-invariants Q_m .

We start by remarking that under some natural homogeneity assumptions, if such an extension exists, it is unique.

PROPOSITION 2.9. 1) Assume that $q \in \mathbb{C}[y_1, \dots, y_n]$ is a homogeneous polynomial of degree d. If there exists a differential operator M_q with coefficients in $\mathbb{C}(\mathfrak{h})$, of the form

$$M_q = q(\partial_{y_1}, \ldots, \partial_{y_n}) + l.o.t.$$

such that $[M_q, L] = 0$, whose homogeneity degree is -d, then M_q is unique.

2) Let $\mathbb{C}[q_1,\ldots,q_n]\subseteq B\subseteq \mathbb{C}[y_1,\ldots,y_n]$ be a graded ring. Assume that we have a linear map $M\colon B\to \mathcal{D}(U)$ such that, if $q\in B$ is homogeneous of degree d, then $[M_q,L]=0$, M_q has homogeneity degree -d, and

$$M_q = q(\partial_{y_1}, \ldots, \partial_{y_n}) + l.o.t.$$

Then M is a ring homomorphism and $M_q = L_q$ for all $q \in \mathbb{C}[q_1, \ldots, q_n]$.

Proof. 1) If there exist two different operators M_q and M'_q with these properties, take $M_q - M'_q$. This operator has degree of homogeneity -d, but order smaller than d. Therefore, its symbol S(x,y) is not a polynomial. On the other hand, since the symbol of L is $\sum y_i^2$, we get that $[L, M_q - M'_q] = 0$ implies $\{\sum y_i^2, S(x,y)\} = 0$. Write S in the form K(x,y)/H(x) with K is a polynomial, and H(x) a homogeneous polynomial of positive degree t (we assume that K(x,y) and H(x) have no common irreducible factors). Then

$$0 = \left\{ \sum y_i^2, S(x, y) \right\} = 2 \frac{\sum_{i=1}^n y_i K_{x_i}(x, y) H(x) - \sum_{i=1}^n y_i H_{x_i}(x) K(x, y)}{H(x)^2}.$$

Since $\sum_{i=1}^{n} x_i H_{x_i}(x) = tH(x)$, we have $\sum_{i=1}^{n} y_i H_{x_i}(x) K(x, y) \neq 0$. So H(x) must divide this polynomial and, by our assumptions, this implies that it must divide the polynomial $\sum_{i=1}^{n} y_i H_{x_i}(x)$ whose degree in x is t-1. This is a contradiction.

2) Let $q, p \in B$ be two homogeneous elements. Then $M_q M_p$ and M_{pq} both satisfy the same homogeneity assumptions. Hence they are equal by 1).

Finally if $q \in \mathbb{C}[q_1, \dots, q_n]$, both M_q and L_q satisfy the same homogeneity assumptions. Hence they are equal by 1). \square

The required extension to the ring of m-quasi-invariants is then provided by the following

THEOREM 2.10 ([CV1, CV2]). Let $c = m \colon \Sigma \to \mathbf{Z}_+$. The following two conditions are equivalent for a homogeneous polynomial $q \in \mathbf{C}[\mathfrak{h}^*]$ of degree d.

1) There exists a differential operator

$$L_q = q(\partial_{y_1}, \dots, \partial_{y_n}) + l.o.t.$$

of homogeneity degree -d, such that $[L_q, L] = 0$.

2) q is an m-quasi-invariant homogeneous of degree d.

Using this, we can extend system (5) to the system

(6)
$$L_p \psi = p(k)\psi, \quad p \in Q_m, \quad k \in \operatorname{Spec} Q_m = X_m.$$

(Recall that, as a set, $X_m = \mathfrak{h}$.) Near a generic point $x_0 \in \mathfrak{h}$, system (6) has a one dimensional space of solutions, thus there exists a unique up to scaling solution $\psi(k,x)$, which can be expressed in elementary functions. This solution is called the *Baker-Akhiezer function*, and has the form

$$\psi(k, x) = P(k, x) e^{(k,x)}$$

with P(k,x) a polynomial of the form $\delta(x)\delta(k)+l.o.t.$ and $e^{(k,x)}$ denotes the exponential function computed in the scalar product (k,x). Furthermore, it can be shown that $\psi(k,x)=\psi(x,k)$ (see [CV1, CV2, FV]).

These results motivate the following terminology. The variety X_m is called the spectral variety of the Calogero-Moser system for the multiplicity function m, and Q_m is called the spectral ring of this system.

2.7 AN EXAMPLE

EXAMPLE 2.11. Let $W = \mathbb{Z}/2$, $\mathfrak{h} = \mathbb{C}$, m = 1. As we have seen, Q_m has a basis given by the monomials $\{x^{2i}\} \cup \{x^{2i+3}\}$, $i \geq 0$. Let us set for such a monomial, $L_{x^r} = L_r$, and $\partial = \frac{d}{dx}$. Then we have

$$L_0 = 1$$
, $L_2 = \partial^2 - \frac{2}{x}\partial$, $L_3 = \partial^3 - \frac{3}{x}\partial^2 + \frac{3}{x^2}\partial$.

As for the others, $L_{2j} = L_2^j$, $L_{2j+3} = L_2^j L_3$. (Note that L_1 is not defined). The system (6) in this case is

$$\begin{cases} \psi'' - \frac{2}{x}\psi' = k^2\psi, \\ \psi''' - \frac{3}{x}\psi'' + \frac{3}{x^2}\psi' = k^3\psi. \end{cases}$$