

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 3.3 Differential operators on X_m
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We now pass to a general L_q , $q \in Q_m$. We may assume that q is homogeneous of, say, degree d . By Corollary 3.3 we have that L_q is a non zero multiple of $(adL)^d(q)$. Since both q and L preserve Q_m , our claim follows. \square

3.3 DIFFERENTIAL OPERATORS ON X_m

Now let us return to the algebra of differential operators $\mathcal{D}(X_m)$. Notice that $\mathcal{D}(X_m)$ contains two commutative subalgebras (both isomorphic to Q_m). The first is Q_m itself, the second is the subalgebra Q_m^\dagger consisting of the differential operators of the form L_q with $q \in Q_m$. It is possible to prove

THEOREM 3.6 ([BEG]). $\mathcal{D}(X_m)$ is generated by Q_m and Q_m^\dagger .

Notice that by Corollary 3.3 we in fact have that $\mathcal{D}(X_m)$ is generated by Q_m and by L .

EXAMPLE 3.7. If $W = \mathbf{Z}/2$, $\mathfrak{h} = \mathbf{C}$ we get that $\mathcal{D}(X_m)$ is generated by the operators

$$x^2, \quad x^{2m+1}, \quad \frac{d^2}{dx^2} - \frac{2m}{x} \frac{d}{dx}.$$

Theorem 3.6 together with Proposition 3.4, imply

COROLLARY 3.8 ([BEG]). $\mathcal{D}(X_m)$ is locally finite dimensional under the action of the Lie algebra $\mathfrak{sl}(2)$ defined in (8).

This Corollary implies that our $\mathfrak{sl}(2)$ action on $\mathcal{D}(X_m)$ can be integrated to an action of the group $SL(2)$. In particular we have

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} q = L_q$$

for all $q \in Q_m$. This transformation is a generalization of the Fourier transform, since it reduces to the usual Fourier transform on differential operators on \mathfrak{h} when $m = 0$.

EXAMPLE 3.9. If $W = \mathbf{Z}/2$, $\mathfrak{h} = \mathbf{C}$, we get that the monomials $\{x^{2i}\} \cup \{x^{2i+2m+1}\}$ are (up to constants) all lowest weight vectors for the $\mathfrak{sl}(2)$ action on $\mathcal{D}(X_m)$. x^n has weight $-n$. We deduce that $\mathcal{D}(X_m)$ is isomorphic as a $\mathfrak{sl}(2)$ -module to the direct sum of the irreducible representations of dimension $n + 1$ for n even or $n = 2(m + i) + 1$, each with multiplicity one.