

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	49 (2003)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 <b>Artikel:</b>	LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
<b>Autor:</b>	Etingof, Pavel / Strickland, Elisabetta
<b>Kapitel:</b>	3.6 Category O
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-66677">https://doi.org/10.5169/seals-66677</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

DEFINITION 3.13. The *spherical subalgebra* of  $H_c$  is the algebra  $eH_ce$ .

Notice that  $1 \notin eH_ce$ . On the other hand, since  $ex = xe = e$  for  $x \in eH_ce$ ,  $e$  is the unit for the spherical subalgebra. We can embed both  $\mathbf{C}[\mathfrak{h}^*]^W$  and  $\mathbf{C}[\mathfrak{h}]^W$  in the spherical subalgebra as follows. Take  $f \in \mathbf{C}[\mathfrak{h}^*]^W$  (the other case is identical) and set  $m_e(f) = fe$ . Since  $f$  is invariant, we have  $efe = fe^2 = fe = m_e(f)$ , so that  $m_e$  actually maps  $\mathbf{C}[\mathfrak{h}^*]^W$  to  $eH_ce$ . The injectivity is clear from the PBW-theorem. As for the fact that  $m_e$  is a homomorphism, we have  $m_e(fg) = fge = fge^2 = fege = m_e(f)m_e(g)$ . From now on, we will consider both  $\mathbf{C}[\mathfrak{h}^*]^W$  and  $\mathbf{C}[\mathfrak{h}]^W$  as subalgebras of the spherical subalgebra.

### 3.6 CATEGORY $\mathcal{O}$

We are now going to study representations of the algebras  $H_c$  and  $eH_ce$ .

DEFINITION 3.14. The category  $\mathcal{O}(H_c)$  (resp.  $\mathcal{O}(eH_ce)$ ) is the full subcategory of the category of  $H_c$ -modules (resp.  $eH_ce$ -modules) whose objects are the modules  $M$  such that

- 1)  $M$  is finitely generated.
- 2) For all  $v \in M$ , the subspace  $\mathbf{C}[\mathfrak{h}^*]^Wv \subset M$  is finite dimensional.

We can define a functor

$$F: \mathcal{O}(H_c) \rightarrow \mathcal{O}(eH_ce)$$

by setting  $F(M) = eM$ . It is easy to show that  $F(M)$  is an object of  $\mathcal{O}(eH_ce)$ .

We are now going to explain how to construct some modules in  $\mathcal{O}(H_c)$  which, by analogy with the case of enveloping algebras of semisimple Lie algebras, we will call Whittaker and Verma modules. First, take  $\lambda \in \mathfrak{h}^*$ . Denote by  $W_\lambda \subset W$  the stabilizer of  $\lambda$ . Take an irreducible  $W_\lambda$ -module  $\tau$ . We define a structure of  $\mathbf{C}[\mathfrak{h}^*] \rtimes \mathbf{C}[W_\lambda]$ -module on  $\tau$  by

$$(fw)v = f(\lambda)(wv) \quad \forall v \in \tau, w \in W_\lambda, f \in \mathbf{C}[\mathfrak{h}^*].$$

It is easy to see that this action is well defined and we denote this module by  $\lambda\#\tau$ . We can then consider the  $H_c$ -module

$$M(\lambda, \tau) = H_c \otimes_{\mathbf{C}[\mathfrak{h}^*] \rtimes \mathbf{C}[W_\lambda]} \lambda\#\tau.$$

This is called a Whittaker module. In the special case  $\lambda = 0$  (and hence  $W_\lambda = W$ ), the module  $M(0, \tau)$  is called a Verma module. It is clear that these are objects of  $\mathcal{O}$ . Notice that as  $\mathbf{C}[\mathfrak{h}] \rtimes \mathbf{C}[W]$ -module,  $M(\lambda, \tau) = \mathbf{C}[\mathfrak{h}] \otimes_{\mathbf{C}} \mathbf{C}[W] \otimes_{\mathbf{C}[W_\lambda]} \tau$ .

EXAMPLE 3.15. If  $\lambda = 0$  and  $\tau = \mathbf{1}$  is the trivial representation of  $W$ , the Verma module  $M(0, \mathbf{1}) = \mathbf{C}[\mathfrak{h}]$ . The action of  $\mathbf{C}[\mathfrak{h}]$  is given by multiplication, that of  $\mathbf{C}[\mathfrak{h}^*]$  is generated by the Dunkl operators and  $W$  acts in the usual way.

### 3.7 GENERIC $c$

Opdam and Rouquier have recently studied the structure of the categories  $\mathcal{O}(H_c)$ ,  $\mathcal{O}(eH_ce)$ , and found that it is especially simple if  $c$  is “generic” in a certain sense. Namely, recall that for a  $W$ -invariant function  $q: \Sigma \rightarrow \mathbf{C}^*$  one can define the *Hecke algebra*  $\text{He}_q(W)$  to be the quotient of the group algebra of the fundamental group of  $U/W$  by the relations  $(T_s - 1)(T_s + q_s) = 0$ , where  $T_s$  is the image in  $U/W$  of a small half-circle around the hyperplane of  $s$  in the counterclockwise direction. It is well known that  $\text{He}_q(W)$  is an algebra of dimension  $|W|$ , which coincides with  $\mathbf{C}[W]$  if  $q = 1$ . It is also known that  $\text{He}_q(W)$  is semisimple (and isomorphic to  $\mathbf{C}[W]$  as an algebra) unless  $q_s$  belongs for some  $s$  to a finite set of roots of unity depending on  $W$  (see [Hu]).

DEFINITION 3.16. The function  $c$  is said to be *generic* if for  $q = e^{2\pi i c}$ , the Hecke algebra  $\text{He}_q(W)$  is semisimple.

In particular, any irrational  $c$  is generic, and (more important for us) an integer valued  $c$  is generic (since in this case  $q = 1$ ). We can now state the following central result:

**THEOREM 3.17** (Opdam-Rouquier [OR]; see also [BEG] for an exposition). *If  $c$  is generic (in particular, if  $c$  takes non negative integer values), then the irreducible objects in  $\mathcal{O}$  are exactly the modules  $M(\lambda, \tau)$ . Moreover, the category  $\mathcal{O}$  is semisimple.*

We also have

**THEOREM 3.18** ([OR]). *If  $c$  is generic then the functor  $F$  is an equivalence of categories.*

From Theorem 3.17 we can deduce

**THEOREM 3.19** ([BEG]). *If  $c$  is generic, then  $H_c$  is a simple algebra.*

In the case  $c = 0$ , we get the simplicity of  $\mathbf{C}[\mathfrak{h} \oplus \mathfrak{h}^*] \rtimes \mathbf{C}[W]$ , which is well known.