

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
Autor: Etingof, Pavel / Strickland, Elisabetta
Kapitel: 3.9 The action of the Cherednik algebra to quasi-invariants
DOI: <https://doi.org/10.5169/seals-66677>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3.8 THE LEVASSEUR-STAFFORD THEOREM AND ITS GENERALIZATION

Let us now recall a result of Levasseur and Stafford:

THEOREM 3.20 ([LS]). *If G is a finite group acting on a finite dimensional vector space V over the complex numbers, then the ring $\mathcal{D}(V)^G$ is generated by the subrings $\mathbf{C}[V]^G$ and $\mathbf{C}[V^*]^G$.*

As an example, notice that if we let $\mathbf{Z}/n\mathbf{Z}$ act on the complex line by multiplication by the n^{th} roots of 1, we deduce that the operator $x \frac{d}{dx}$ can be expressed as a non commutative polynomial in the operators x^n and $\frac{d^n}{dx^n}$, a non-obvious fact. We note also that this theorem has a purely “quantum” nature, i.e. the corresponding “classical” statement, saying that the Poisson algebra $\mathbf{C}[V \times V^*]^G$ is generated, as a Poisson algebra, by $\mathbf{C}[V]^G$ and $\mathbf{C}[V^*]^G$, is in fact false, already for $V = \mathbf{C}$ and $G = \mathbf{Z}/n\mathbf{Z}$.

One can prove a similar result for the algebra $eH_c e$. Namely, recall that the algebra $eH_c e$ contains the subalgebras $\mathbf{C}[\mathfrak{h}]^W$, and $\mathbf{C}[\mathfrak{h}^*]^W$.

THEOREM 3.21 ([BEG]). *If c is generic then the two subalgebras $\mathbf{C}[\mathfrak{h}]^W$ and $\mathbf{C}[\mathfrak{h}^*]^W$ generate $eH_c e$.*

Notice that if $c = 0$, then $eH_0 e = \mathcal{D}(\mathfrak{h})^W$, so Theorem 3.21 reduces to the Levasseur-Stafford theorem.

REMARK. It is believed that this result holds without the assumption of generic c . Moreover, it is known to be true for all c if W is a Weyl group not of type E and F , since in this case Wallach proved that the corresponding classical statement for Poisson algebras holds true. Nevertheless, the genericity assumption is needed for the proof, because, similarly to the proof of the Levasseur-Stafford theorem, it is based on the simplicity of H_c .

3.9 THE ACTION OF THE CHEREDNIK ALGEBRA TO QUASI-INVARIANTS

We now go back to the study of Q_m . Notice that the algebra $eH_m e$ acts on $\mathbf{C}[\mathfrak{h}]^W$, since e gives the W -equivariant projection of $\mathbf{C}[\mathfrak{h}]$ onto $\mathbf{C}[\mathfrak{h}]^W$. It is clear that this action is by differential operators. For instance, the subalgebra $\mathbf{C}[\mathfrak{h}]^W \subset eH_m e$ acts by multiplication. Also, an element $q \in \mathbf{C}[\mathfrak{h}^*]^W \subset eH_m e$ acts via the operator $q(D_{x_1}, \dots, D_{x_n})$. By definition this operator coincides with L_q on $\mathbf{C}[\mathfrak{h}]^W$.

The following important theorem shows that this action extends to Q_m .

THEOREM 3.22 ([BEG]). *There exists a unique representation of the algebra eH_me on Q_m in which an element $q \in \mathbf{C}[\mathfrak{h}]^W$ acts by multiplication and an element $q \in \mathbf{C}[\mathfrak{h}^*]^W$ by L_q .*

Proof. Since by Proposition 3.5, L_q preserves Q_m , we get a uniquely defined representation of the subalgebra of eH_me generated by $\mathbf{C}[\mathfrak{h}]^W$ and $\mathbf{C}[\mathfrak{h}^*]^W$ on Q_m . The result now follows from Theorem 3.21. \square

3.10 PROOF OF THEOREM 1.8

Finally we can prove Theorem 1.8.

To do this, observe that as an eH_me -module, Q_m is in the category $\mathcal{O}(eH_me)$, and $\mathbf{C}[\mathfrak{h}^*]^W$ acts locally nilpotently in Q_m (by degree arguments). We can now apply Theorem 3.18 and Theorem 3.17 and deduce that Q_m is a direct sum of modules of the form $eM(0, \tau)$. As a $\mathbf{C}[\mathfrak{h}] \rtimes \mathbf{C}[W]$ -module, $M(0, \tau) = \mathbf{C}[\mathfrak{h}] \otimes \tau$. On the other hand, by Chevalley's theorem, there is an isomorphism $\mathbf{C}[\mathfrak{h}] \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \mathbf{C}[W]$, commuting with the action of W and $\mathbf{C}[\mathfrak{h}]^W$. Thus we get an isomorphisms of $\mathbf{C}[\mathfrak{h}]^W$ -modules

$$eM(0, \tau) \simeq (M(0, \tau))^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes (\mathbf{C}[W] \otimes \tau)^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \tau,$$

proving that $eM(0, \tau)$ and hence Q_m is a free $\mathbf{C}[\mathfrak{h}]^W$ -module. \square

EXAMPLE 3.23. For $W = \mathbf{Z}/2$ and $\mathfrak{h} = \mathbf{C}$, take the polynomials $1, x^{2m+1}$. Notice that $L(1) = L(x^{2m+1}) = 0$ while $s(1) = 1, s(x^{2m+1}) = -x^{2m+1}, s \in \mathbf{Z}/2$ being the element of order two. It follows that Q_m as a eH_me -module is the direct sum of $\mathbf{C}[x^2] \oplus x^{2m+1}\mathbf{C}[x^2]$. These modules are irreducible. Moreover, $\mathbf{C}[x^2] \simeq eM(0, \mathbf{1}), x^{2m+1}\mathbf{C}[x^2] \simeq eM(0, \varepsilon), \varepsilon$ being the sign representation.

3.11 PROOF OF THEOREM 1.15

Let I be a nonzero two-sided ideal in $\mathcal{D}(X_m)$. First we claim that I nontrivially intersects Q_m . Indeed, otherwise let $K \in I$ be a lowest order nonzero element in I . Since the order of K is positive, there exists $f \in Q_m$ such that $[K, f] \neq 0$. Then $[K, f] \in I$ is of smaller order than K , a contradiction.

Now let $f \in Q_m$ be an element of I . Then $g = \prod_{w \in W} {}^w f \in I$. But g is W -invariant. This shows that the intersection J of I with the subalgebra H_m in $\mathcal{D}(X_m)$ is nonzero. But H_m is simple by Theorem 3.19, so $J = H_m$. Hence, $1 \in J \subset I$, and $I = \mathcal{D}(X_m)$. \square