

<b>Zeitschrift:</b>	L'Enseignement Mathématique
<b>Herausgeber:</b>	Commission Internationale de l'Enseignement Mathématique
<b>Band:</b>	49 (2003)
<b>Heft:</b>	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
<b>Artikel:</b>	LECTURES ON QUASI-INVARIANTS OF COXETER GROUPS AND THE CHEREDNIK ALGEBRA
<b>Autor:</b>	Etingof, Pavel / Strickland, Elisabetta
<b>Kapitel:</b>	3.11 Proof of Theorem 1.15
<b>DOI:</b>	<a href="https://doi.org/10.5169/seals-66677">https://doi.org/10.5169/seals-66677</a>

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

The following important theorem shows that this action extends to  $Q_m$ .

**THEOREM 3.22 ([BEG]).** *There exists a unique representation of the algebra  $eH_m e$  on  $Q_m$  in which an element  $q \in \mathbf{C}[\mathfrak{h}]^W$  acts by multiplication and an element  $q \in \mathbf{C}[\mathfrak{h}^*]^W$  by  $L_q$ .*

*Proof.* Since by Proposition 3.5,  $L_q$  preserves  $Q_m$ , we get a uniquely defined representation of the subalgebra of  $eH_m e$  generated by  $\mathbf{C}[\mathfrak{h}]^W$  and  $\mathbf{C}[\mathfrak{h}^*]^W$  on  $Q_m$ . The result now follows from Theorem 3.21.  $\square$

### 3.10 PROOF OF THEOREM 1.8

Finally we can prove Theorem 1.8.

To do this, observe that as an  $eH_m e$ -module,  $Q_m$  is in the category  $\mathcal{O}(eH_m e)$ , and  $\mathbf{C}[\mathfrak{h}^*]^W$  acts locally nilpotently in  $Q_m$  (by degree arguments). We can now apply Theorem 3.18 and Theorem 3.17 and deduce that  $Q_m$  is a direct sum of modules of the form  $eM(0, \tau)$ . As a  $\mathbf{C}[\mathfrak{h}] \rtimes \mathbf{C}[W]$ -module,  $M(0, \tau) = \mathbf{C}[\mathfrak{h}] \otimes \tau$ . On the other hand, by Chevalley's theorem, there is an isomorphism  $\mathbf{C}[\mathfrak{h}] \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \mathbf{C}[W]$ , commuting with the action of  $W$  and  $\mathbf{C}[\mathfrak{h}]^W$ . Thus we get an isomorphisms of  $\mathbf{C}[\mathfrak{h}]^W$ -modules

$$eM(0, \tau) \simeq (M(0, \tau))^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes (\mathbf{C}[W] \otimes \tau)^W \simeq \mathbf{C}[\mathfrak{h}]^W \otimes \tau,$$

proving that  $eM(0, \tau)$  and hence  $Q_m$  is a free  $\mathbf{C}[\mathfrak{h}]^W$ -module.  $\square$

**EXAMPLE 3.23.** For  $W = \mathbf{Z}/2$  and  $\mathfrak{h} = \mathbf{C}$ , take the polynomials  $1, x^{2m+1}$ . Notice that  $L(1) = L(x^{2m+1}) = 0$  while  $s(1) = 1, s(x^{2m+1}) = -x^{2m+1}, s \in \mathbf{Z}/2$  being the element of order two. It follows that  $Q_m$  as a  $eH_m e$ -module is the direct sum of  $\mathbf{C}[x^2] \oplus x^{2m+1}\mathbf{C}[x^2]$ . These modules are irreducible. Moreover,  $\mathbf{C}[x^2] \simeq eM(0, \mathbf{1}), x^{2m+1}\mathbf{C}[x^2] \simeq eM(0, \varepsilon)$ ,  $\varepsilon$  being the sign representation.

### 3.11 PROOF OF THEOREM 1.15

Let  $I$  be a nonzero two-sided ideal in  $\mathcal{D}(X_m)$ . First we claim that  $I$  nontrivially intersects  $Q_m$ . Indeed, otherwise let  $K \in I$  be a lowest order nonzero element in  $I$ . Since the order of  $K$  is positive, there exists  $f \in Q_m$  such that  $[K, f] \neq 0$ . Then  $[K, f] \in I$  is of smaller order than  $K$ , a contradiction.

Now let  $f \in Q_m$  be an element of  $I$ . Then  $g = \prod_{w \in W} w f \in I$ . But  $g$  is  $W$ -invariant. This shows that the intersection  $J$  of  $I$  with the subalgebra  $H_m$  in  $\mathcal{D}(X_m)$  is nonzero. But  $H_m$  is simple by Theorem 3.19, so  $J = H_m$ . Hence,  $1 \in J \subset I$ , and  $I = \mathcal{D}(X_m)$ .  $\square$