

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	49 (2003)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	SOME REMARKS ON NONCONNECTED COMPACT LIE GROUPS
Autor:	HÄMMERLI, Jean-François
Kapitel:	2. Compact Lie groups :a review
DOI:	https://doi.org/10.5169/seals-66678

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

is shown that centralizers of principal subgroups are extensions of Γ by the center Z_o of G_o that completely control the situation. The Main Theorem is proved in Section 4 and as an illustration, two examples are then given. The final section relates the approach taken in the present work with the natural question of the splitting of the extension associated to a compact Lie group. As an application of principal subgroups, a revisited “Sandwich” Theorem is proved. Particular cases where the extension is always split are also described, and, finally, a “minimal” extension failing to be split is exhibited.

ACKNOWLEDGMENTS. The material in this note is taken from my Ph.D. thesis [13]. It is a pleasure to thank warmly my advisor Professor U. Suter for his guidance and constant encouragement. I am also indebted to M. Matthey for his careful reading of an earlier version of this paper and for his useful comments.

2. COMPACT LIE GROUPS: A REVIEW

In this section, we recall the existence of subgroups of G whose related extensions have close relationships to that corresponding to G . First, we introduce more notation. Let T be a fixed maximal torus in G_o , and let LT denote its Lie algebra. Let B be a basis of the root system $R = R(G_o, T)$ of G_o associated to T . Let \mathcal{H} denote the maximal semisimple ideal of the Lie algebra LG_o of G_o ; the *principal diagonal* of G_o with respect to B is the 1-dimensional subspace given by $D(B) = \{X \in LT : \alpha(X) = \beta(X), \text{ for all } \alpha, \beta \in B\} \cap \mathcal{H} \subset LT$. The image $\Delta = \Delta(B) = \exp(D(B))$ of this subspace under the exponential map is easily seen to be a closed subgroup of T , isomorphic to the circle group S^1 . With a slight abuse of language, we will also call this subgroup a *principal diagonal*. We are now ready to recall the definition of one of the key notions of the present work.

DEFINITION 2.1 (de Siebenthal). A *principal subgroup* of G_o (associated to T) is a connected closed subgroup H such that H is not contained in any proper connected closed subgroup of maximal rank, and such that $\Delta(B) \subset H$ for some basis B of R .

The work of de Siebenthal shows that any compact Lie group possesses a principal subgroup of rank 1, thus isomorphic to $SU(2)$ or $SO(3)$, and that two such principal subgroups are conjugate [8].

NOTE. For the rest of this work, “principal subgroup” will always mean principal subgroup *of rank 1*.

Before stating the main result of this section, which is a direct consequence of the results of de Siebenthal, we introduce three subgroups of G . Let H_T be a principal subgroup associated to T , and let $Z = Z_G(H_T)$ denote its centralizer in G . This subgroup will play a crucial role in the paper. Let also $N = N_G(T)$ be the normalizer of T in G . We will use the convenient notation $N_o = N_{G_o}(T)$ for the intersection of N with G_o , but one should not be confused, N_o is *not* connected (its group of components being the Weyl group of G_o). Finally, we will consider the centralizer $Q = Z_G(\Delta)$ in G of a principal diagonal Δ .

THEOREM 2.2. *For any compact Lie group G there exists a commutative diagram*

$$\begin{array}{ccccc}
 Z_o & \hookrightarrow & Z & \twoheadrightarrow & \Gamma \\
 \downarrow & & \downarrow & & \parallel \\
 T & \hookrightarrow & Q & \twoheadrightarrow & \Gamma \\
 \downarrow & & \downarrow & & \parallel \\
 N_o & \hookrightarrow & N & \twoheadrightarrow & \Gamma \\
 \downarrow & & \downarrow & & \parallel \\
 G_o & \hookrightarrow & G & \twoheadrightarrow & \Gamma
 \end{array}$$

where each row is a group extension.

Proof. The centralizer of H_T in G_o is equal to the center Z_o (by a theorem of Borel and de Siebenthal [3], this property characterizes the closed subgroups of G_o that are not contained in any proper connected closed subgroup of maximal rank [5, Ex. 15, p. 116]). As Z intersects every component of G ([8], Théorème 4, pp. 253–254), we get an extension $Z_o \hookrightarrow Z \twoheadrightarrow \Gamma$. The other statements are deduced from the fact that $\Delta \subset T$ contains a regular element, i.e. an element that is contained in exactly one maximal torus, namely T in the present case (see [12] or [17] for more details). \square

REMARK 2.3. We call the subgroup Q an *extended maximal torus* of G . These subgroups share some important properties with maximal tori: they are all conjugate, and fixing one of them, its conjugates by the elements of G_o cover the whole group G . They appear in the literature under various disguises (see for instance Oliver [19], Section 1, and Segal [22], §4), as explained in [12].

In the final section, we will see how the splitting of the extension associated to G is related to the splitting of the extensions associated to N and Q that appear in Theorem 2.2.

We end this section by recalling a very important result relating the inner, “usual”, and outer automorphism groups of a connected compact Lie group. This result is one of the main reasons why the case of compact Lie groups is well controlled when applying the theory of group extensions, as we will see in Section 3. For the proof, we refer to de Siebenthal [9, Théorème, pp. 46–47] (for another approach consult Bourbaki [5], §4.10).

THEOREM 2.4 (de Siebenthal). *Let G_0 be a connected compact Lie group and let $H \subset G_0$ be a principal subgroup. Then the extension*

$$\text{Inn}(G_0) \xhookrightarrow{\iota} \text{Aut}(G_0) \xrightarrow{\pi} \text{Out}(G_0)$$

is split, i.e.

$$\text{Aut}(G_0) \cong \text{Inn}(G_0) \rtimes \text{Out}(G_0).$$

A possible splitting is given by $s: \text{Out}(G_0) \rightarrow \text{Aut}(G_0)$, where $s(\alpha)$ is the unique automorphism in $\pi^{-1}(\alpha)$ fixing H pointwise.

REMARK 2.5. The fact that the extension associated to $\text{Aut}(G_0)$ is split was known before the work of de Siebenthal, at least in the semisimple case, and appeared in a paper of Dynkin [10].

3. COMPACT LIE GROUPS AND EXTENSIONS

We assume knowledge of the classical relationship between group extensions and related cohomology groups of low degree, as first introduced by Eilenberg and MacLane [11]. For readers not familiar with it, the textbooks by MacLane [16], Robinson [21], or Adem-Milgram [2], provide a thorough treatment; a more concise approach can be found in Kirillov’s book [15], and a sketch in Brown’s [6]. We now want to apply this relationship to the case of compact Lie groups. We fix a nonabelian connected compact Lie group G_0 , a finite group Γ , and a homomorphism $\varphi: \Gamma \rightarrow \text{Out}(G_0)$. Recall that Z_0 denotes the center of G_0 . Choosing a principal subgroup $H \subset G_0$ and fixing s as in Theorem 2.4, we get the commutative diagram