Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

**Band:** 49 (2003)

**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME REMARKS ON NONCONNECTED COMPACT LIE GROUPS

Autor: HÄMMERLI, Jean-François

**Kapitel:** 3. Compact Lie groups and extensions

**DOI:** https://doi.org/10.5169/seals-66678

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 15.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

In the final section, we will see how the splitting of the extension associated to G is related to the splitting of the extensions associated to N and Q that appear in Theorem 2.2.

We end this section by recalling a very important result relating the inner, "usual", and outer automorphism groups of a connected compact Lie group. This result is one of the main reasons why the case of compact Lie groups is well controlled when applying the theory of group extensions, as we will see in Section 3. For the proof, we refer to de Siebenthal [9, Théorème, pp. 46–47] (for another approach consult Bourbaki [5], §4.10).

THEOREM 2.4 (de Siebenthal). Let  $G_o$  be a connected compact Lie group and let  $H \subset G_o$  be a principal subgroup. Then the extension

$$\operatorname{Inn}(G_0) \stackrel{\iota}{\hookrightarrow} \operatorname{Aut}(G_0) \stackrel{\pi}{\twoheadrightarrow} \operatorname{Out}(G_0)$$

is split, i.e.


$$\operatorname{Aut}(G_{o}) \cong \operatorname{Inn}(G_{o}) \rtimes \operatorname{Out}(G_{o})$$
.

A possible splitting is given by  $s: Out(G_o) \to Aut(G_o)$ , where  $s(\alpha)$  is the unique automorphism in  $\pi^{-1}(\alpha)$  fixing H pointwise.

REMARK 2.5. The fact that the extension associated to  $Aut(G_0)$  is split was known before the work of de Siebenthal, at least in the semisimple case, and appeared in a paper of Dynkin [10].

## 3. COMPACT LIE GROUPS AND EXTENSIONS

We assume knowledge of the classical relationship between group extensions and related cohomology groups of low degree, as first introduced by Eilenberg and Mac Lane [11]. For readers not familiar with it, the textbooks by Mac Lane [16], Robinson [21], or Adem-Milgram [2], provide a thorough treatment; a more concise approach can be found in Kirillov's book [15], and a sketch in Brown's [6]. We now want to apply this relationship to the case of compact Lie groups. We fix a nonabelian connected compact Lie group  $G_0$ , a finite group  $\Gamma$ , and a homomorphism  $\varphi \colon \Gamma \to \operatorname{Out}(G_0)$ . Recall that  $Z_0$  denotes the center of  $G_0$ . Choosing a principal subgroup  $H \subset G_0$  and fixing s as in Theorem 2.4, we get the commutative diagram



In the sequel, we will use the notation  $\sigma_{\gamma} = (s \circ \varphi)(\gamma)$ , for  $\gamma \in \Gamma$ . Let  $\mathcal{E}(\Gamma, G_0, \varphi) \subset \mathcal{E}$  denote the subset of equivalence classes giving rise to  $\varphi$ . In the particular case of compact Lie groups, one has the following results.

### PROPOSITION 3.1.

- (i) The set of equivalence classes of extensions  $\mathcal{E}(\Gamma, G_o, \varphi)$  is in bijection with the cohomology group  $H^2_{\overline{\varphi}}(\Gamma; Z_o)$ .
- (ii) For all  $u \in H^2_{\overline{\varphi}}(\Gamma; Z_o)$  the corresponding extension  $G_o \hookrightarrow G \twoheadrightarrow \Gamma$  carries a natural structure of Lie group.

*Proof.* It suffices to check that  $\mathcal{E}(\Gamma, G_o, \varphi) \neq \emptyset$  to prove (i). But this follows from Theorem 2.4: the semidirect product  $G = G_o \rtimes_{so\varphi} \Gamma$  exists for any  $\varphi$ . The second statement is easily deduced from classical Lie group theory.  $\square$ 

The bijection in the latter proposition is not canonical, as it depends on the choice of a particular element in  $\mathcal{E}(\Gamma,G_0,\varphi)$ . On the other hand, there is a canonical bijection between  $H^2_{\overline{\varphi}}(\Gamma;Z_0)$  and the set  $\mathcal{E}(\Gamma,Z_0,\overline{\varphi})$  of equivalence classes of extension of  $\Gamma$  by  $Z_0$  with action given by  $\overline{\varphi}$ . Therefore, there is a bijection  $\Lambda\colon \mathcal{E}(\Gamma,Z_0,\overline{\varphi})\to \mathcal{E}(\Gamma,G_0,\varphi)$  still depending on the previous choice. Let us describe this bijection by first expliciting the cohomology group  $H^2_{\overline{\varphi}}(\Gamma;Z_0)=Z^2_{\overline{\varphi}}(\Gamma;Z_0)/B^2_{\overline{\varphi}}(\Gamma;Z_0)$ . Keeping the multiplicative notation in  $Z_0$ , the cocycles, i.e. the elements of  $Z^2_{\overline{\varphi}}(\Gamma;Z_0)$ , are functions  $h\colon \Gamma\times\Gamma\to Z_0$  satisfying  $h(\gamma_1,e)=h(e,\gamma_2)=e$  (normalization), and

$$(\delta h)(\gamma_1, \gamma_2, \gamma_3) = \sigma_{\gamma_1}(h(\gamma_2, \gamma_3)) \cdot h(\gamma_1 \gamma_2, \gamma_3)^{-1} \cdot h(\gamma_1, \gamma_2 \gamma_3) \cdot h(\gamma_1, \gamma_2)^{-1} = e$$

for all  $\gamma_1, \gamma_2, \gamma_3 \in \Gamma$ . The coboundaries, i.e. the elements of  $B^2_{\overline{\varphi}}(\Gamma; Z_0)$ , are functions  $h: \Gamma \times \Gamma \to Z_0$  such that there exists a function  $k: \Gamma \to Z_0$ , with k(e) = e, satisfying

$$h(\gamma_1, \gamma_2) = (\delta k)(\gamma_1, \gamma_2) = \sigma_{\gamma_1}(k(\gamma_2)) \cdot k(\gamma_1 \gamma_2)^{-1} \cdot k(\gamma_1)$$

for all  $\gamma_1, \gamma_2 \in \Gamma$ . Let us choose the semidirect product  $G_o \rtimes \Gamma$  associated to the section s as the extension corresponding to the neutral element in  $H^2_{\overline{\varphi}}(\Gamma; Z_o)$ . Then, for  $u = [h] \in H^2_{\overline{\varphi}}(\Gamma; Z_o)$ , the corresponding class of extensions is

given by  $[G_o \hookrightarrow G_h \twoheadrightarrow \Gamma]$ , where  $G_h$  is the set  $G_o \times \Gamma$  equipped with the multiplication

$$(g, \gamma) *_h (g', \gamma') = (g \cdot \sigma_{\gamma}(g') \cdot h(\gamma, \gamma'), \gamma \cdot \gamma')$$

(see [16], Chapter IV, §4 and §8). We will also denote by  $G_o \hookrightarrow G_u \twoheadrightarrow \Gamma$  any representative of the class of extensions corresponding to  $u \in H^2_{\overline{\varphi}}(\Gamma; Z_o)$ . We now give a canonical description of the inverse of  $\Lambda$ , i.e. a description that does *not* depend on the choice of a particular element in  $\mathcal{E}(\Gamma, G_o, \varphi)$ .

LEMMA 3.2. For any principal subgroup H in  $G_0$ , the map

$$\Theta \colon \mathcal{E}(\Gamma, G_{o}, \varphi) \longrightarrow \mathcal{E}(\Gamma, Z_{o}, \overline{\varphi}), \ [G_{o} \hookrightarrow G \twoheadrightarrow \Gamma] \longmapsto [Z_{o} \hookrightarrow Z_{G}(H) \twoheadrightarrow \Gamma]$$

is the inverse of  $\Lambda$  (and does not depend on the choice of H). In particular it is a bijection.

*Proof.* As centralizers of principal subgroups are preserved by isomorphisms of G,  $\Theta$  does not depend on the choice of a representative in  $[G_o \hookrightarrow G \twoheadrightarrow \Gamma]$ . Let  $u = [Z_o \hookrightarrow E_h \twoheadrightarrow \Gamma] = [h] \in H^2_{\overline{\varphi}}(\Gamma, Z_o)$ . Then, we have the commutative diagram

$$Z_{0} \xrightarrow{} E_{h} \xrightarrow{} \Gamma$$

$$\downarrow \qquad \qquad \qquad \parallel$$

$$G_{0} \xrightarrow{} G = G_{h} \xrightarrow{} \Gamma$$

where  $E_h$  is  $Z_o \times \Gamma$  as a set. Let us show that  $E_h = Z_{G_h}(H)$ , H being the principal subgroup of  $G_o$  corresponding to the fixed section s. By Theorem 2.2, it is enough to check that  $E_h$  is contained in  $Z_{G_h}(H)$ . Let  $(z, \gamma) \in E_h$  and  $(x, e) \in H \subset G_o \subset G_h$ . We calculate

$$(z, \gamma) *_h (x, e) = (z \cdot \sigma_{\gamma}(x) \cdot h(\gamma, e), \gamma)$$
  
=  $(z \cdot x, \gamma)$ ,

and


$$(x, e) *_h (z, \gamma) = (x \cdot \sigma_e(z) \cdot h(e, \gamma), \gamma)$$
$$= (x \cdot z, \gamma)$$
$$= (z \cdot x, \gamma),$$

by normalization, and because the restriction of  $\sigma_{\gamma}$  to H is the identity by the choice of the section s.

Now, as the principal subgroups are all conjugate by an element of  $G_0$  (see [8], Théorème, pp. 46–47), so are their centralizers. Therefore, the extensions  $Z_0 \hookrightarrow Z_G(H) \twoheadrightarrow \Gamma$ , for H running through the family of principal subgroups, all belong to the same class. This shows that  $\Theta$  is well defined and satisfies  $\Theta \circ \Lambda = id_{\mathcal{E}(\Gamma, Z_0, \overline{\varphi})}$ . As  $\Lambda$  is bijective, this shows that  $\Theta = \Lambda^{-1}$ .  $\square$ 

We summarize the situation exposed in this section.

THEOREM 3.3. Suppose given  $G_o$ , a homomorphism  $\varphi \colon \Gamma \to \operatorname{Out}(G_o)$  and an extension  $Z_o \hookrightarrow Z \twoheadrightarrow \Gamma$ , for which the homomorphism  $\Gamma \to \operatorname{Aut}(Z_o)$  coincides with  $\overline{\varphi}$ . Then, up to equivalence of extensions, there exists a unique compact Lie group G fitting into the commutative diagram



where the rows are group extensions. Moreover the given data allow the construction of an extension  $G_o \hookrightarrow G \twoheadrightarrow \Gamma$ , in which the subgroup Z is the centralizer of a principal subgroup.

Conversely, the class of the extension  $Z_o \hookrightarrow Z \twoheadrightarrow \Gamma$  in  $G_o \hookrightarrow G \twoheadrightarrow \Gamma$  can be recovered by taking the centralizer of any principal subgroup.

# 4. PROOF OF THE MAIN THEOREM AND EXAMPLES

We are almost ready to show that the map described in the Introduction is an action of  $\operatorname{Out}(G_0) \times \operatorname{Aut}(\Gamma)$  on the set

$$\mathcal{E} pprox \coprod_{\varphi \in \operatorname{Hom}(\Gamma,\operatorname{Out}(G_{\mathsf{o}}))} H^2_{\overline{\varphi}}(\Gamma; Z_{\mathsf{o}}) \,.$$

We first introduce some notation. For an element g in a group K, we will write  $c_g$  for conjugation by g, i.e.  $c_g(x) = gxg^{-1}$ , for all x in K. For  $\alpha \in \text{Out}(G_0)$ , we will choose  $\widetilde{\alpha} \in \text{Aut}(G_0)$  such that  $\pi(\widetilde{\alpha}) = \alpha$ , and we will denote the restricted automorphism by  $\overline{\alpha} \in \text{Aut}(Z_0)$ . Finally,