Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 49 (2003)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SOME REMARKS ON NONCONNECTED COMPACT LIE GROUPS

Autor: HÄMMERLI, Jean-François

Kapitel: 5. Splitting of the extension ASSOCIATED TO A NONCONNECTED

COMPACT LIE GROUP

DOI: https://doi.org/10.5169/seals-66678

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

5. SPLITTING OF THE EXTENSION ASSOCIATED TO A NONCONNECTED COMPACT LIE GROUP

Let G still denote a compact Lie group with a nonabelian connected component; we also assume the other notations introduced previously. In this final section, we make a few observations on the following problem: when is the natural extension associated to G split, i.e. when is G isomorphic to a semidirect product $G_0 \times \Gamma$? Our aim is to relate this problem to the rest of this work. For a deeper analysis one should consult Chapter 6 in the book by Hofmann and Morris [14]. We start with a structure theorem for compact Lie groups based on centralizers of principal subgroups, similar to the "Sandwich Theorem for compact Lie groups" (see [14], Corollary 6.75, p. 272). This theorem shows that any compact Lie group is "sandwiched" in between two semidirect products closely related to it. We then recall a theorem of de Siebenthal and compare, in some particular cases, the question of the splitting of the extension associated to G to that of the extensions associated to the normalizer N of a maximal torus and to the centralizer Q of a principal diagonal, both introduced in Section 2. As an application of the "Sandwich" Theorem, the final proposition presents a "minimal" compact Lie group Gsuch that the associated extension is *not* split.

Let \overline{G}_{o} denote the adjoint group G_{o}/Z_{o} ; it is well-known that the center of \overline{G}_{o} is trivial.

THEOREM 5.1. Let G be a compact Lie group with a nonabelian connected component. Then there exist two surjective homomorphisms

$$G_s = G_o \rtimes Z \xrightarrow{\pi_1} G \xrightarrow{\pi_2} \overline{G} = G/Z_o \cong \overline{G}_o \rtimes \Gamma,$$
 $(g_o, z) \longmapsto g_o \cdot z$

where the centralizer Z of a fixed principal subgroup acts on G_o by conjugation, and where π_2 is the canonical projection corresponding to the normal subgroup Z_o of G.

For the kernels, we have $\ker \pi_1 \cong Z_o$ and $\ker \pi_2 = Z_o$; in particular


$$G \cong (G_0 \rtimes Z)/Z_0$$
 and $G/Z_0 \cong \overline{G}_0 \rtimes \Gamma$.

Proof. As \overline{G}_0 is centerless, \overline{G} must be isomorphic to $\overline{G}_0 \rtimes \Gamma$ by the proof of Proposition 3.1. The other assertions about π_2 are clear.

The map π_1 is well-defined and surjective (because Z intersects every component of G). Straightforward computations show that it is a homomorphism and that $\ker \pi_1 \cong Z_0$. \square

REMARK 5.2. 1) The component of the identity of G_s is equal to G_o if and only if G_o is semisimple.

- 2) The present version of the "Sandwich" Theorem has the advantage of being more explicit than the one in [14] (the result therein is an existence theorem). Its drawbacks are the fact that Z is not finite if G_0 is not semisimple, and that it obviously makes no sense for compact Lie groups with an abelian connected component.
- 3) Given a homomorphism $\varphi \colon \Gamma \to \operatorname{Out}(G_0)$, and an extension $Z_0 \stackrel{\mu}{\hookrightarrow} E \stackrel{\nu}{\twoheadrightarrow} \Gamma$ for which the action coincides with the "restriction" $\overline{\varphi}$, there is a more direct way than the cohomological one to recover the corresponding compact Lie group G, i.e. the one that fits into the commutative diagram



Let us define the composition $\overline{\sigma} \colon E \xrightarrow{\nu} \Gamma \xrightarrow{\varphi} \operatorname{Out}(G_0) \xrightarrow{s} \operatorname{Aut}(G_0)$, where s is as in Theorem 2.4. Then by Bourbaki (see [4], Lemme 7, pp. 210–211), we have

$$G = (G_0 \rtimes_{\bar{\sigma}} E)/\Delta Z_0$$

where ΔZ_0 is the image of the injection $z_0 \mapsto (z_0^{-1}, \mu(z_0))$. Taking E = Z, this gives another proof of the assertions concerning π_1 in Theorem 5.1.

Using Cartan subgroups (in the sense of Segal [22], i.e. those Adams called "SS subgroups" in honour of Segal and de Siebenthal [1]), de Siebenthal gave some explicit sufficient conditions for the splitting of the extension associated to G ([9], Théorème p. 74).

Theorem 5.3 (de Siebenthal). Let G be a compact Lie group with G_o simply connected, or of adjoint type (i.e. Z_o is trivial). If $\Gamma = \pi_o(G)$ is cyclic then G is a semidirect product, i.e. $G \cong G_o \rtimes \Gamma$.

A relationship with the splitting of the extension associated to the normalizer of a maximal torus N in G is given in the next proposition.

PROPOSITION 5.4. If the group of components Γ of G is nilpotent, then the extension $G_o \hookrightarrow G \twoheadrightarrow \Gamma$ is split if and only if the extension $N_o \hookrightarrow N \twoheadrightarrow \Gamma$ is split.

Proof. The "if" part is clear. Conversely, let $s: \Gamma \to G$ be a section. By a result in Bourbaki (see [5], Corollaire 4, p. 49), any nilpotent subgroup of a compact Lie group is contained in the normalizer of some maximal torus. Therefore, if needed after conjugation by an element in G_0 , we have $s(\Gamma) \subset N$, and we can conclude that the extension associated to N is split. \square

For an extended maximal torus Q, the extensions can be related as follows.

PROPOSITION 5.5. If the group of components Γ of G is cyclic, then the extension $G_0 \hookrightarrow G \twoheadrightarrow \Gamma$ is split if and only if the extension $T \hookrightarrow Q \twoheadrightarrow \Gamma$ is split.

Proof. The proposition readily follows from the fact that the conjugates of Q cover G. \square

REMARK 5.6. This latter proposition fails in general. An obstruction to the splitting of the extension associated to the extended maximal torus Q can be found in a paper by Oliver; this obstruction involves the representation ring of G and its relation with the family of all p-toral subgroups of G (see [19], Corollary 3.11). In particular, Oliver constructs a compact Lie group $G = SU(2) \times (\mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/3)$ such that the extension corresponding to the extended maximal torus Q is *not* split ([19], pp. 376–377).

We conclude with the promised example.

PROPOSITION 5.7. Let $D_8 = \langle r, s | r^4 = s^2 = e, srs^{-1} = r^{-1} \rangle$ be a presentation of the dihedral group. Then the quotient

$$G = (SU(2) \times D_8)/\Delta \mathbf{Z}/2$$
,

where $\Delta \mathbf{Z}/2$ denotes the central subgroup generated by $(-1, r^2)$, is a compact Lie group with $G_0 \cong SU(2)$ and $\Gamma \cong \mathbf{Z}/2 \times \mathbf{Z}/2$, the 4-group of Klein. The associated extension

$$SU(2) \hookrightarrow G \twoheadrightarrow \mathbf{Z}/2 \times \mathbf{Z}/2$$

is not split. Among the extensions associated to a compact Lie group with a nonabelian connected component, it is a minimal one having the property of being non-split, in the sense that the rank of the connected component and the size of the group of components are minimal. Moreover replacing the

connected component by SO(3) (i.e. by "any" group of the same rank), or the group of components by $\mathbb{Z}/4$ (i.e. by "any" group of the same size), will force the extension to be split.

Proof. The assertions about the connected component and the group of components are clear. Let us show that the extension associated to G is not split. Let us denote $[g, \gamma] \in G$ the image of $(g, \gamma) \in SU(2) \times D_8$ under the canonical projection. Let S^1 denote the standard maximal torus in SU(2), and let N denote its normalizer in G. We have

$$N = \left\{ [t, e] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [jt, e] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [t, r] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [jt, r] : t \in \mathbf{S}^1 \right\}$$

$$\coprod \left\{ [t, s] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [jt, s] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [t, rs] : t \in \mathbf{S}^1 \right\} \coprod \left\{ [jt, rs] : t \in \mathbf{S}^1 \right\}.$$

By contradiction, suppose that the extension associated to G is split, i.e. there exists a section. As $\mathbb{Z}/2 \times \mathbb{Z}/2$ is abelian, thus nilpotent, we deduce, by Proposition 5.4, that the extension associated to N is also split. We want to show that this is not possible by considering the elements of order 2 in N. For n=0,1, a straightforward calculation shows that in the component corresponding to $r^n s$, an element $[t,r^n s]$ is of order 2 if and only if $t=\pm 1$, and that the sub-component $\{[jt,r^n s]:t\in \mathbf{S}^1\}$ does not contain any element of order 2. Two of the three non-trivial elements in $\Gamma\cong \mathbb{Z}/2\times \mathbb{Z}/2$ must thus be mapped by the section to $[\pm 1,s]$ and $[\pm 1,rs]$. Therefore, as the section is a homomorphism, the image of the third non-trivial element is

$$[\pm \mathbf{1}, rs] \cdot [\pm \mathbf{1}, s] = [\pm \mathbf{1}, r] ,$$

which is not of order 2. A contradiction that shows that the extension associated to G is not split.

The property of minimality follows by Theorem 5.3, and by the fact that any extension with SO(3) as normal subgroup is a direct product (because SO(3) is complete, i.e. centerless and with trivial outer automorphism group).

REFERENCES

- [1] ADAMS, J. F. Maps between classifying spaces, II. *Invent. Math.* 49 (1978), 1–65.
- [2] ADEM, A. and R. J. MILGRAM. Cohomology of Finite Groups. Springer, 1994.
- [3] BOREL, A. and J. DE SIEBENTHAL Les sous-groupes fermés de rang maximum des groupes de Lie clos. *Comment. Math. Helv. 23* (1949), 200–221.
- [4] BOURBAKI, N. Groupes et algèbres de Lie (Chapitres 2 et 3). Hermann, 1972.
- [5] Groupes et algèbres de Lie (Chapitre 9). Masson, 1983.