

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 49 (2003)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** ATIYAH'S  $L^2$ -INDEX THEOREM  
**Autor:** Chatterji, Indira / Mislin, Guido  
**DOI:** <https://doi.org/10.5169/seals-66679>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## ATIYAH'S $L^2$ -INDEX THEOREM

by Indira CHATTERJI and Guido MISLIN

### 1. INTRODUCTION

The  $L^2$ -Index Theorem of Atiyah [1] expresses the index of an elliptic operator on a closed manifold  $M$  in terms of the  $G$ -equivariant index of some regular covering  $\tilde{M}$  of  $M$ , with  $G$  the group of covering transformations. Atiyah's proof is analytic in nature. Our proof is algebraic and involves an embedding of a given group into an acyclic one, together with naturality properties of the indices.

### 2. REVIEW OF THE $L^2$ -INDEX THEOREM

The main reference for this section is Atiyah's paper [1]. All manifolds considered are smooth Riemannian, without boundary. Covering spaces of manifolds carry the induced smooth and Riemannian structure. Let  $M$  be a closed manifold and let  $E, F$  denote two complex (Hermitian) vector bundles over  $M$ . Consider an elliptic pseudo-differential operator

$$D: C^\infty(M, E) \rightarrow C^\infty(M, F)$$

acting on the smooth sections of the vector bundles. One defines its space of solutions

$$S_D = \{s \in C^\infty(M, E) \mid Ds = 0\}.$$

The complex vector space  $S_D$  has finite dimension (see [13]), and so has  $S_{D^*}$  the space of solutions of the adjoint  $D^*$  of  $D$  where

$$D^*: C^\infty(M, F) \rightarrow C^\infty(M, E)$$

is the unique continuous linear map satisfying

$$\langle Ds, s' \rangle = \int_M \langle Ds(m), s'(m) \rangle_F dm = \langle s, D^* s' \rangle = \int_M \langle s(m), D^* s'(m) \rangle_E dm$$

for all  $s \in C^\infty(M, E)$ ,  $s' \in C^\infty(M, F)$ . One now defines the *index* of  $D$  as follows :

$$\text{Index}(D) = \dim_{\mathbf{C}}(S_D) - \dim_{\mathbf{C}}(S_{D^*}) \in \mathbf{Z}.$$

An explicit formula for  $\text{Index}(D)$  is given by the famous Atiyah-Singer Theorem (cf. [2]). Consider a not necessarily connected, regular covering  $\pi: \tilde{M} \rightarrow M$  with countable covering transformation group  $G$ . The projection  $\pi$  can be used to define an elliptic operator

$$\tilde{D} := \pi^*(D): C_c^\infty(\tilde{M}, \pi^* E) \rightarrow C_c^\infty(\tilde{M}, \pi^* F).$$

Denote by  $S_{\tilde{D}}$  the closure of  $\{s \in C_c^\infty(\tilde{M}, \pi^* E) \mid \tilde{D}s = 0\}$  in  $L^2(\tilde{M}, \pi^* E)$ . Let  $\tilde{D}^*$  denote the adjoint of  $\tilde{D}$ . The space  $S_{\tilde{D}}$  is not necessarily finite dimensional, but being a closed  $G$ -invariant subspace of the  $L^2$ -completion  $L^2(\tilde{M}, \pi^* E)$  of the space of smooth sections with compact supports  $C_c^\infty(\tilde{M}, \pi^* E)$ , its von Neumann dimension is therefore defined as follows. Write

$$\mathcal{N}(G) = \{P: \ell^2(G) \rightarrow \ell^2(G) \text{ bounded and } G\text{-invariant}\}$$

for the group von Neumann algebra of  $G$ , where  $G$  acts on  $\ell^2(G)$  via the right regular representation. Then  $S_{\tilde{D}}$  is a finitely generated Hilbert  $G$ -module and hence can be represented by an idempotent matrix  $P = (p_{ij}) \in M_n(\mathcal{N}(G))$  (recall that a finitely generated Hilbert  $G$ -module is isometrically  $G$ -isomorphic to a Hilbert  $G$ -subspace of the Hilbert space  $\ell^2(G)^n$  for some  $n \geq 1$ , see [9]). One then sets

$$\dim_G(S_{\tilde{D}}) = \sum_{i=1}^n \langle p_{ii}(e), e \rangle = \kappa(P) \in \mathbf{R},$$

where by abuse of notation  $e$  denotes the element in  $\ell^2(G)$  taking value 1 on the neutral element  $e \in G$  and 0 elsewhere (see Eckmann's survey [9] on  $L^2$ -cohomology for more on von Neumann dimensions). The map  $\kappa: M_n(\mathcal{N}(G)) \rightarrow \mathbf{C}$  is the Kaplansky trace. One defines the  $L^2$ -index of  $\tilde{D}$  by

$$\text{Index}_G(\tilde{D}) = \dim_G(S_{\tilde{D}}) - \dim_G(S_{\tilde{D}^*}).$$

We can now state Atiyah's  $L^2$ -Index Theorem.

THEOREM 2.1 (Atiyah [1]). *For  $D$  an elliptic pseudo-differential operator on a closed Riemannian manifold  $M$*

$$\text{Index}(D) = \text{Index}_G(\tilde{D})$$

for any countable group  $G$  and any lift  $\tilde{D}$  of  $D$  to a regular  $G$ -cover  $\tilde{M}$  of  $M$ .

In particular, the  $L^2$ -index of  $\tilde{D}$  is always an integer, even though it is a priori given in terms of real numbers. The following serves as an illustration of the  $L^2$ -Index Theorem.

EXAMPLE 2.2 (Atiyah's formula [1]). Let  $\Omega^\bullet$  be the de Rham complex of complex valued differential forms on the closed connected manifold  $M$  and consider the de Rham differential  $D = d + d^* : \Omega^{ev} \rightarrow \Omega^{odd}$ . Let  $\pi : \tilde{M} \rightarrow M$  be the universal cover of  $M$  so that  $G = \pi_1(M)$ . Then

- $\text{Index}(D) = \chi(M)$ , the ordinary Euler characteristic of  $M$ .
- $\text{Index}_G(\tilde{D}) = \sum_j (-1)^j \beta^j(M)$ , the  $L^2$ -Euler characteristic of  $M$ .

The  $\beta^j(M)$ 's denote the  $L^2$ -Betti numbers of  $M$ . Thus the  $L^2$ -Index Theorem translates into Atiyah's formula

$$\chi(M) = \sum_j (-1)^j \beta^j(M).$$

We recall that the  $L^2$ -Betti numbers  $\beta^j(M)$  are in general not integers. For instance, if  $\pi_1(M)$  is a finite group, one checks that

$$\beta^j(M) = \frac{1}{|\pi_1(M)|} b^j(\tilde{M}),$$

where  $b^j(\tilde{M})$  stands for the ordinary  $j$ 'th Betti number of the universal cover  $\tilde{M}$  of  $M$ . In particular, for  $1 < |\pi_1(M)| < \infty$ ,  $\beta^0(M) = 1/|\pi_1(M)|$  is not an integer and the  $L^2$ -Index Theorem reduces to the well-known fact that

$$\chi(M) = \frac{\chi(\tilde{M})}{|\pi_1(M)|}.$$

It is a conjecture (Atiyah Conjecture) that for a general closed connected manifold  $M$  the  $L^2$ -Betti numbers  $\beta^j(M)$  are always rational numbers, and even integers in case that  $\pi_1(M)$  is torsion-free. For some interesting examples, which might lead to counterexamples, see Dicks and Schick [8].

## 3. HILBERT MODULES

Recall that for  $H < G$  and  $X$  an  $H$ -space, the *induced*  $G$ -space is

$$G \times_H X = (G \times X)/H$$

where  $H$  acts on  $G \times X$  via  $h \cdot (g, x) = (gh^{-1}, hx)$  and the left  $G$ -action on  $G \times_H X$  is given by  $g \cdot [k, x] = [gk, x]$  (where  $[k, x]$  denotes the class of the pair  $(k, x) \in G \times X$  in  $G \times_H X$ ). For  $A \subseteq \ell^2(H)^n$  a Hilbert  $H$ -module one defines  $\text{Ind}_H^G(A)$ , the *induced* Hilbert  $G$ -module, as follows:

$$\text{Ind}_H^G(A) = \left\{ f: G \rightarrow A, \quad f(gh) = h^{-1}f(g), \quad \sum_{\gamma \in G/H} \|f(\gamma)\|^2 < \infty \right\}.$$

On  $\text{Ind}_H^G(A)$  the action of  $G$  is given as follows:

$$(\gamma \cdot f)(\mu) = f(\gamma^{-1}\mu), \quad \gamma, \mu \in G \text{ and } f \in \text{Ind}_H^G(A).$$

For  $\tilde{M}$  an  $H$ -free, cocompact Riemannian manifold and  $\tilde{D}$  an  $H$ -equivariant pseudo-differential operator on  $\tilde{M}$ , one can express the lift  $\bar{D}$  of  $\tilde{D}$  to  $\bar{M} = G \times_H \tilde{M}$  as follows. Fix a set  $R$  of representatives for  $G/H$  and write  $\pi: \bar{M} \rightarrow \tilde{M}$  for the projection; a section  $\bar{s} \in C_c^\infty(\bar{M}, \pi^*E)$  is a collection

$$\bar{s} = \{\tilde{s}_r\}_{r \in R},$$

where  $\tilde{s}_r \in C_c^\infty(\tilde{M}, E)$  is the zero section for all but finitely many  $r$ 's, and  $\bar{s}([g, \tilde{m}]) = \tilde{s}_r(h\tilde{m})$ , if  $[r, h\tilde{m}] = [g, \tilde{m}] \in G \times_H \tilde{M}$ . Now the lift  $\bar{D}$  of  $\tilde{D}$  to  $\bar{M} = G \times_H \tilde{M}$  satisfies

$$\bar{D}\bar{s} = \left\{ \tilde{D}\tilde{s}_r \right\}_{r \in R}.$$

LEMMA 3.1. *Let  $M$  be a closed Riemannian manifold,  $D$  a pseudo-differential operator on  $M$  and  $\tilde{M}$  a regular cover of  $M$  with countable transformation group  $H$ . Consider an inclusion  $H < G$  and form the regular cover  $\bar{M} = G \times_H \tilde{M}$  of  $M$ . Then for the lifts  $\tilde{D}$  of  $D$  to  $\tilde{M}$  and  $\bar{D}$  of  $\tilde{D}$  to  $\bar{M}$ ,*

$$\text{Index}_H(\tilde{D}) = \text{Index}_G(\bar{D}).$$

*Proof.* It is enough to see that  $S_{\bar{D}} \cong \text{Ind}_H^G(S_{\tilde{D}})$ . Indeed, it is well-known (see [9]) that for a Hilbert  $H$ -module  $A$  one has

$$\dim_H(A) = \dim_G(\text{Ind}_H^G(A)).$$

For  $R$  a fixed set of representatives for  $G/H$ , the map

$$\begin{aligned}\varphi_R: \text{Ind}_H^G(S_{\widetilde{D}}) &\rightarrow S_{\bar{D}} \\ f &\mapsto \{f(r)\}_{r \in R}\end{aligned}$$

is well-defined by  $H$ -equivariance of the elements of  $S_{\widetilde{D}}$  and one checks that it defines a  $G$ -equivariant isometric bijection. Similarly for the adjoint operators.

The following example is a particular case of the previous lemma.

EXAMPLE 3.2. Let us look at the case  $\widetilde{M} = M \times G$ . A section  $\widetilde{s} \in C_c^\infty(\widetilde{M}, \pi^*E)$  is an element  $\widetilde{s} = \{s_g\}_{g \in G}$  where  $s_g \in C^\infty(M, E)$  and  $s_g = 0$  for all but finitely many  $g$ 's. Note that  $L^2(\widetilde{M}, \pi^*E)$  can be identified with  $\ell^2(G) \otimes L^2(M, E)$ . Now

$$\widetilde{D}\widetilde{s} = \{Ds_g\}_{g \in G} \in C_c^\infty(\widetilde{M}, \pi^*F)$$

and hence  $S_{\widetilde{D}}$  may be identified with  $\ell^2(G) \otimes S_D \cong \ell^2(G)^d$ , where  $d = \dim_{\mathbb{C}}(S_D)$ . In this identification the projection  $P$  onto  $S_{\widetilde{D}}$  becomes the identity in  $M_d(\mathcal{N}(G))$  and thus

$$\dim_G(S_{\widetilde{D}}) = \sum_{i=1}^d \langle e, e \rangle = d = \dim_{\mathbb{C}}(S_D).$$

A similar argument for  $D^*$  shows that in this case not only does the  $L^2$ -Index of  $\widetilde{D}$  coincide with the Index of  $D$ , but also the individual terms of the difference correspond to each other. This is not the case in general, see Example 2.2.

#### 4. ON $K$ -HOMOLOGY

Many ideas of this section go back to the seminal article by Baum and Connes [3], which has been circulating for many years and has only recently been published.

An elliptic pseudo-differential operator  $D$  on the closed manifold  $M$  can also be used to define an element  $[D] \in K_0(M)$ , the  $K$ -homology of  $M$ , and according to Baum and Douglas [4], all elements of  $K_0(M)$  are of the form  $[D]$ . The index defined in Section 2 extends to a well-defined

homomorphism (cf. [4])

$$\text{Index}: K_0(M) \rightarrow \mathbf{Z},$$

such that  $\text{Index}([D]) = \text{Index}(D)$ . On the other hand, the projection  $\text{pr}: M \rightarrow \{pt\}$  induces, after identifying  $K_0(\{pt\})$  with  $\mathbf{Z}$ , a homomorphism

$$(*) \quad \text{pr}_*: K_0(M) \rightarrow \mathbf{Z},$$

which, as explained in [4], satisfies

$$\text{pr}_*([D]) = \text{Index}([D]).$$

More generally (cf. [4]), for a not necessarily finite CW-complex  $X$ , every  $x \in K_0(X)$  is of the form  $f_*[D]$  for some  $f: M \rightarrow X$ , and  $K_0(X)$  is obtained as a colimit over  $K_0(M_\alpha)$ , where the  $M_\alpha$  form a directed system consisting of closed Riemannian manifolds (these homology groups  $K_0(X)$  are naturally isomorphic to the ones defined using the Bott spectrum; sometimes, they are referred to as  $K$ -homology groups with *compact supports*). The index map from above extends to a homomorphism

$$\text{Index}: K_0(X) \rightarrow \mathbf{Z},$$

such that  $\text{Index}(x) = \text{Index}([D])$  if  $x = f_*[D]$ , with  $f: M \rightarrow X$ .

We now consider the case of  $X = BG$ , the classifying space of the discrete group  $G$ , and obtain thus for any  $f: M \rightarrow BG$  a commutative diagram

$$\begin{array}{ccc} K_0(M) & \xrightarrow{\text{Index}} & \mathbf{Z} \\ f_* \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}} & \mathbf{Z}. \end{array}$$

Note that  $(*)$  from above implies the following naturality property for the index homomorphism.

LEMMA 4.1. *For any homomorphism  $\varphi: H \rightarrow G$  one has a commutative diagram*

$$\begin{array}{ccc} K_0(BH) & \xrightarrow{\text{Index}} & \mathbf{Z} \\ (B\varphi)_* \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}} & \mathbf{Z}. \end{array} \quad \square$$

We now turn to the  $L^2$ -index of Section 2. It extends to a homomorphism

$$\text{Index}_G: K_0(BG) \rightarrow \mathbf{R}$$

as follows. Each  $x \in K_0(BG)$  is of the form  $f_*(y)$  for some  $y = [D] \in K_0(M)$ ,  $f: M \rightarrow BG$ ,  $M$  a closed smooth manifold and  $D$  an elliptic operator on  $M$ . Let  $\tilde{D}$  be the lifted operator to  $\tilde{M}$ , the  $G$ -covering space induced by  $f: M \rightarrow BG$ . Then put

$$\text{Index}_G(x) := \text{Index}_G(\tilde{D}).$$

One checks that  $\text{Index}_G(x)$  is indeed well-defined, either by direct computation, or by identifying it with  $\tau(x)$ , where  $\tau$  denotes the composite of the assembly map  $K_0(BG) \rightarrow K_0(C_r^*G)$  with the natural trace  $K_0(C_r^*G) \rightarrow \mathbf{R}$  (for this latter point of view, see Higson-Roe [10]; for a discussion of the assembly map see e.g. Kasparov [12], or Valette [14]). The following naturality property of this index map is a consequence of Lemma 3.1.

LEMMA 4.2. *For  $H < G$  the following diagram commutes:*

$$\begin{array}{ccc} K_0(BH) & \xrightarrow{\text{Index}_H} & \mathbf{R} \\ \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}_G} & \mathbf{R}. \quad \square \end{array}$$

Atiyah's  $L^2$ -Index Theorem 2.1 for a given  $G$  can now be expressed as the statement (as already observed in [10])

$$\text{Index}_G = \text{Index}: K_0(BG) \rightarrow \mathbf{R}.$$

## 5. ALGEBRAIC PROOF OF ATIYAH'S $L^2$ -INDEX THEOREM

Recall that a group  $A$  is said to be *acyclic* if  $H_*(BA, \mathbf{Z}) = 0$  for  $* > 0$ . For  $G$  a countable group, there exists an embedding  $G \rightarrow A_G$  into a countable acyclic group  $A_G$ . There are many constructions of such a group  $A_G$  available in the literature, see for instance Kan-Thurston [11, Proposition 3.5], Berrick-Varadarajan [5] or Berrick-Chatterji-Mislin [6]; these different constructions are to be compared in Berrick's forthcoming work [7]. It follows that the suspension  $\Sigma BA_G$  is contractible, and therefore the inclusion  $\{e\} \rightarrow A_G$

induces an isomorphism

$$K_0(B\{e\}) \xrightarrow{\cong} K_0(BA_G).$$

Our strategy is as follows. We show that the Atiyah  $L^2$ -Index Theorem holds in the special case of acyclic groups, and finish the proof combining the above embedding of a group into an acyclic group.

*Proof of Theorem 2.1.* If a group  $A$  is acyclic, the equation  $\text{Index}_A = \text{Index}$  follows from the diagram

$$\begin{array}{ccccc} K_0(BA) & \xrightarrow{\text{Index}_A} & \mathbf{R} & \xleftarrow{\text{Index}} & K_0(BA) \\ \cong \uparrow & & \uparrow & & \cong \uparrow \\ K_0(B\{e\}) & \xrightarrow[\cong]{\text{Index}_{\{e\}}} & \mathbf{Z} & \xleftarrow[\cong]{\text{Index}} & K_0(B\{e\}) \end{array}$$

because  $\text{Index}_{\{e\}} = \text{Index}$  on the bottom line. For a general group  $G$ , consider an embedding into an acyclic group  $A_G$  and complete the proof by using Lemma 3.1, together with Lemmas 4.1 and 4.2.

## REFERENCES

- [1] ATIYAH, M. F. Elliptic operators, discrete groups and von Neumann algebras. *Astérisque* 32–3 (1976), 43–72.
- [2] ATIYAH, M. F. and I. M. SINGER. The index of elliptic operators III. *Ann. of Math.* (2) 87 (1968), 546–604.
- [3] BAUM, P. and A. CONNES.  $K$ -theory for Lie groups and foliations. *L'Enseignement Math.* (2) 46 (2000), 3–42.
- [4] BAUM, P. and R. DOUGLAS.  $K$ -homology and index theory. *Proceedings of Symposia in Pure Mathematics* 38, Part 1 (1982), 117–173.
- [5] BERRICK, A. J. and K. VARADARAJAN. Binate towers of groups. *Arch. Math.* 62 (1994), 97–111.
- [6] BERRICK, A. J., I. CHATTERJI and G. MISLIN. From acyclic groups to the Bass Conjecture for amenable groups. (Submitted for publication 2002.)
- [7] BERRICK, A. J. The acyclic group dichotomy. (Preprint in preparation.)
- [8] DICKS, W. and T. SCHICK. The spectral measure of certain elements of the complex group ring of a wreath product. *Geom. Dedicata* 93 (2002), 121–137.
- [9] ECKMANN, B. Introduction to  $l_2$ -methods in topology: reduced  $l_2$ -homology, harmonic chains,  $l_2$ -Betti numbers. (Notes prepared by Guido Mislin.) *Israel J. Math.* 117 (2000), 183–219.
- [10] HIGSON, N. and J. ROE. *Analytic K-Homology*. Oxford Mathematical Monographs, Oxford University Press, 2000.

- [11] KAN, D. M. and W. P. THURSTON. Every connected space has the homology of a  $K(\pi, 1)$ . *Topology* 15 (1976), 253–258.
- [12] KASPAROV, G.  $K$ -theory, group  $C^*$ -algebras, and higher signatures (Conspicetus). Novikov conjectures, index theorems and rigidity, Vol. 1 (Oberwolfach, 1993), 101–146. London Math. Soc. Lecture Note Ser. 226. Cambridge Univ. Press, 1995.
- [13] SOLOVYOV, Y. P. and E. V. TROITSKY.  *$C^*$ -Algebras and Elliptic Operators in Differential Topology*. (Translated from the 1996 Russian original by Troitsky.) Translations of Mathematical Monographs, 192. Amer. Math. Soc., Providence (R.I.), 2001.
- [14] VALETTE, A. *Introduction to the Baum-Connes Conjecture*. (Notes taken by Indira Chatterji. With an appendix by Guido Mislin.) Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2002.

(Reçu le 15 septembre 2002)

Indira Chatterji

Mathematics Department  
Cornell University  
Ithaca NY 14853  
U. S. A.  
*e-mail* : indira@math.cornell.edu

Guido Mislin

Mathematics Department  
ETHZ  
8092 Zürich  
Switzerland  
*e-mail* : mislin@math.ethz.ch

**vide-leer-empty**