Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ATIYAH'S L^2-INDEX THEOREM
Autor: Chatterji, Indira / Mislin, Guido
Kapitel: 3. Hilbert modules
DOI: https://doi.org/10.5169/seals-66679

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 12.02.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. Hilbert modules

Recall that for $H < G$ and X an H-space, the *induced* G-space is

$$G \times_H X = (G \times X)/H$$

where H acts on $G \times X$ via $h \cdot (g, x) = (gh^{-1}, hx)$ and the left G-action on $G \times_H X$ is given by $g \cdot [k, x] = [gk, x]$ (where $[k, x]$ denotes the class of the pair $(k, x) \in G \times X$ in $G \times_H X$). For $A \subseteq \ell^2(H)^n$ a Hilbert H-module one defines $\mathrm{Ind}_H^G(A)$, the *induced* Hilbert G-module, as follows:

$$\mathrm{Ind}_H^G(A) = \left\{ f\colon G \to A, \quad f(gh) = h^{-1}f(g), \quad \sum_{\gamma \in G/H} \|f(\gamma)\|^2 < \infty \right\}.$$

On $\mathrm{Ind}_H^G(A)$ the action of G is given as follows:

$$(\gamma \cdot f)(\mu) = f(\gamma^{-1}\mu), \qquad \gamma, \mu \in G \text{ and } f \in \mathrm{Ind}_H^G(A).$$

For $\widetilde{M}$ an H-free, cocompact Riemannian manifold and $\widetilde{D}$ an H-equivariant pseudo-differential operator on $\widetilde{M}$, one can express the lift $\overline{D}$ of $\widetilde{D}$ to $\overline{M} = G \times_H \widetilde{M}$ as follows. Fix a set R of representatives for G/H and write $\pi\colon \overline{M} \to \widetilde{M}$ for the projection; a section $\overline{s} \in C_c^\infty(\overline{M}, \pi^*E)$ is a collection

$$\overline{s} = \{\widetilde{s}_r\}_{r \in R},$$

where $\widetilde{s}_r \in C_c^\infty(\widetilde{M}, E)$ is the zero section for all but finitely many r's, and $\overline{s}([g, \widetilde{m}]) = \widetilde{s}_r(h\widetilde{m})$, if $[r, h\widetilde{m}] = [g, \widetilde{m}] \in G \times_H \widetilde{M}$. Now the lift $\overline{D}$ of $\widetilde{D}$ to $\overline{M} = G \times_H \widetilde{M}$ satisfies

$$\overline{D}\,\overline{s} = \left\{ \widetilde{D}\widetilde{s}_r \right\}_{r \in R}.$$

Lemma 3.1. *Let M be a closed Riemannian manifold, D a pseudo-differential operator on M and $\widetilde{M}$ a regular cover of M with countable transformation group H. Consider an inclusion $H < G$ and form the regular cover $\overline{M} = G \times_H \widetilde{M}$ of M. Then for the lifts $\widetilde{D}$ of D to $\widetilde{M}$ and $\overline{D}$ of $\widetilde{D}$ to $\overline{M}$,*

$$\mathrm{Index}_H(\widetilde{D}) = \mathrm{Index}_G(\overline{D}).$$

Proof. It is enough to see that $S_{\overline{D}} \cong \mathrm{Ind}_H^G(S_{\widetilde{D}})$. Indeed, it is well-known (see [9]) that for a Hilbert H-module A one has

$$\dim_H(A) = \dim_G(\mathrm{Ind}_H^G(A)).$$

For R a fixed set of representatives for G/H, the map

$$\varphi_R\colon \operatorname{Ind}_H^G(S_{\widetilde{D}}) \to S_{\overline{D}}$$
$$f \mapsto \{f(r)\}_{r\in R}$$

is well-defined by H-equivariance of the elements of $S_{\widetilde{D}}$ and one checks that it defines a G-equivariant isometric bijection. Similarly for the adjoint operators.

The following example is a particular case of the previous lemma.

EXAMPLE 3.2. Let us look at the case $\widetilde{M} = M \times G$. A section $\widetilde{s} \in C_c^\infty(\widetilde{M}, \pi^*E)$ is an element $\widetilde{s} = \{s_g\}_{g\in G}$ where $s_g \in C^\infty(M,E)$ and $s_g = 0$ for all but finitely many g's. Note that $L^2(\widetilde{M}, \pi^*E)$ can be identified with $\ell^2(G) \otimes L^2(M,E)$. Now

$$\widetilde{D}\widetilde{s} = \{Ds_g\}_{g\in G} \in C_c^\infty(\widetilde{M}, \pi^*F)$$

and hence $S_{\widetilde{D}}$ may be identified with $\ell^2(G) \otimes S_D \cong \ell^2(G)^d$, where $d = \dim_{\mathbf{C}}(S_D)$. In this identification the projection P onto $S_{\widetilde{D}}$ becomes the identity in $M_d(\mathcal{N}(G))$ and thus

$$\dim_G(S_{\widetilde{D}}) = \sum_{i=1}^{d} \langle e, e\rangle = d = \dim_{\mathbf{C}}(S_D)\,.$$

A similar argument for D^* shows that in this case not only does the L^2-Index of $\widetilde{D}$ coincide with the Index of D, but also the individual terms of the difference correspond to each other. This is not the case in general, see Example 2.2.

4. ON K-HOMOLOGY

Many ideas of this section go back to the seminal article by Baum and Connes [3], which has been circulating for many years and has only recently been published.

An elliptic pseudo-differential operator D on the closed manifold M can also be used to define an element $[D] \in K_0(M)$, the K-homology of M, and according to Baum and Douglas [4], all elements of $K_0(M)$ are of the form $[D]$. The index defined in Section 2 extends to a well-defined