

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 49 (2003)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: NOTE ON THE HOPF-STIEFEL FUNCTION
Autor: Eliahou, Shalom / Kervaire, Michel
Kapitel: Introduction
DOI: <https://doi.org/10.5169/seals-66683>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A NOTE ON THE HOPF-STIEFEL FUNCTION

by Shalom ELIAHOU^{*)} and Michel KERVAIRE

INTRODUCTION

In the preceding paper of this volume [P], Alain Plagne gives a formula for the (generalized) Hopf-Stiefel function β_p .

Given a prime number p , and two positive integers r, s , recall that $\beta_p(r, s)$ is defined as the smallest integer n such that $(x+y)^n \in (x^r, y^s)$, where (x^r, y^s) is the ideal generated by x^r and y^s in the polynomial ring $\mathbf{F}_p[x, y]$.

Plagne's theorem reads

THEOREM 1. *Let r, s be positive integers, then $\beta_p(r, s)$ is given by the formula*

$$(1) \quad \beta_p(r, s) = \min_{t \in \mathbb{N}} \left(\left\lceil \frac{r}{p^t} \right\rceil + \left\lceil \frac{s}{p^t} \right\rceil - 1 \right) p^t.$$

In [P], this formula is derived as a corollary of a theorem on Additive Number Theory, Theorem 4, which is the main result of the paper.

Here, we give another proof of Theorem 1 using a purely arithmetical argument.

Recall from [EK, p. 22], where $\beta_p(r, s)$ was introduced, that this function can be described in terms of the p -adic expansions of $r - 1$ and $s - 1$ as follows.

^{*)} During the preparation of this paper, the first author has partially benefited from a research contract with the Fonds National Suisse pour la Recherche Scientifique.

THEOREM 2. Let $r - 1 = \sum_{i \geq 0} a_i p^i$ and $s - 1 = \sum_{i \geq 0} b_i p^i$ be the respective p -adic expansions of $r - 1$ and $s - 1$, with $0 \leq a_i, b_i \leq p - 1$ for all i .

Define the integer k as the largest index for which $a_k + b_k \geq p$, if any exists. Otherwise, that is if $a_i + b_i \leq p - 1$ for all $i \geq 0$, set $k = -1$.

Then, $\beta_p(r, s)$ is determined by

$$(2) \quad \beta_p(r, s) = \left(\left\lfloor \frac{r-1}{p^{k+1}} \right\rfloor + \left\lfloor \frac{s-1}{p^{k+1}} \right\rfloor + 1 \right) p^{k+1}.$$

Although the point of Plagne's paper is to stress the relationship of his formula with Additive Number Theory, it is interesting to note that (1) also admits a direct proof using the above Theorem 2.

This is the content of the next section. In Section 2, we provide a simple proof of Theorem 2.

1. DERIVING THEOREM 1 FROM THEOREM 2

It is very easy to understand the relationship of the floor-function $\lfloor \xi \rfloor$, or integral part of ξ , appearing in Theorem 2, with the ceiling-function $\lceil \xi \rceil$, the smallest integer at least as big as ξ , used in formula (1).

The main object of this section will be to locate the minimum over $\ell \geq 0$ of the expression $\left(\left\lceil \frac{r}{p^\ell} \right\rceil + \left\lceil \frac{s}{p^\ell} \right\rceil - 1 \right) p^\ell$ and to show that this minimum is attained at $\ell = k + 1$ with k as defined in Theorem 2.

For every index $\ell \geq 0$, we have

$$0 < \frac{1 + \sum_{i=0}^{\ell-1} a_i p^i}{p^\ell} \leq \frac{1 + \sum_{i=0}^{\ell-1} (p-1)p^i}{p^\ell} = 1. \quad .$$

Since $r = 1 + \sum_{i \geq 0} a_i p^i$, it follows that

$$\left\lceil \frac{r}{p^\ell} \right\rceil = \sum_{i \geq 0} a_{i+\ell} p^i + 1.$$

Similarly, we have $0 \leq \frac{\sum_{i=0}^{\ell-1} a_i p^i}{p^\ell} \leq \frac{\sum_{i=0}^{\ell-1} (p-1)p^i}{p^\ell} = \frac{p^\ell - 1}{p^\ell} < 1$, and

$$(3) \quad \left\lfloor \frac{r-1}{p^\ell} \right\rfloor = \sum_{i \geq 0} a_{i+\ell} p^i.$$