Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 50 (2004)
Heft: 1-2: L'enseignement mathématique

Artikel: A genus formula for some plane curves
Autor: Kang, Ming-Chang
Bibliographie
DOI: https://doi.org/10.5169/seals-2644

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 22.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

STEP 6. Note that $\bar{A}$ is also the normalization of B. We will compute $\dim(\bar{A}/B)$.

By [Se], p. 59, Formula (3), the question can again be reduced to the complete case. Let D be the completion of B. Then

$$D = k[[w,u]]/\left(\epsilon u^{m-n} - w^n \prod_{1\le i\le m}(1+\lambda_i u)\right).$$

Find $\alpha \in D$ such that $\alpha^{m-n} = \epsilon^{-1}\prod_{1\le i\le m}(1+\lambda_i u)$. Define $U = u/\alpha$. Then $D \simeq k[[w,U]]/(U^{m-n} - w^n)$. Now we can apply Lemma 4 to get $\dim(\bar{D}/D) = \{(n-1)(m-n-1) - 1 + d\}/2$.

STEP 7. Finally we find that $\delta_P = \dim(\bar{A}/A) = \dim(B/A) + \dim(\bar{A}/B) = \{(m-1)(m-n-1) - 1 + d\}/2$. Thus

$$(N-1)(N-2)/2 - \delta_P = \{(m-1)(n-1) + 1 - d\}/2$$

because $N = \max\{m,n\} = m$. This completes the proof of Theorem 2.

REMARKS.

(1) From the above proof, it is clear that Theorem 2 remains valid if $\operatorname{char} k = p > 0$ and p doesn't divide $mn\prod_{1\le i\le l} m_i n_i$.

(2) Similarly, if p is a prime number and the affine curve is defined by $y^p = \prod_{1\le i\le l}(x-\lambda_i)^{m_i}$ such that the λ_i are distinct, $1 \le m_i < p$ and p doesn't divide $\sum_{1\le i\le l} m_i$, then Theorem 2 (and its proof for this case) remains valid no matter what $\operatorname{char} k$ may be. Note that the latter assumption can always be achieved. For, if we denote $\sum_{1\le i\le l} m_i$ by m and suppose that $m = pr$, we may assume that $\lambda_1 = 0$. Divide both sides of the equation by x^m. Consider the new variables $u = 1/x$, $v = y/x^r$.

(3) On the other hand, if we assume that k is a perfect field (such that (i) $p \nmid mn\prod_{1\le i\le l} m_i n_i$ if $\operatorname{char} k = p > 0$, or (ii) p is a prime number and the affine curve is defined by $y^p = \prod_{1\le i\le l}(x-\lambda_i)^{m_i}$ with ...) but not algebraically closed, then Theorem 2 is true because we can extend the constant field k to its algebraic closure at the beginning of the proof without affecting the genus by [Ch], p. 99.

REFERENCES

[BK] BRIESKORN, E. and H. KNÖRRER. *Plane Algebraic Curves*. Birkhäuser, Basel, 1986.

[Ca] CASAS-ALVERO, E. *Singularities of Plane Curves*. London Math. Soc. Lecture Note Series *276*. Cambridge Univ. Press, Cambridge, 2000.

[Ch] CHEVALLEY, C. *Introduction to the Theory of Algebraic Functions of One Variable*. Amer. Math. Soc., Providence, 1951.

[DLS] DAVENPORT, H., D. J. LEWIS and A. SCHINZEL. Equations of the form $f(x) = g(y)$. *Quart. J. Math. Oxford Ser. (2) 12* (1961), 304–312.

[Fe] FEIT, W. Some consequences of the classification of finite simple groups. In: *Proc. Symp. Pure Math. 37* (1980), 175–181.

[Hi] HIRONAKA, H. On the arithmetic genera and the effective genera of algebraic curves. *Mem. Coll. Sci. Univ. Kyoto Ser. A. Math. 30* (1957), 177–195.

[Mi] MIRANDA, R. *Algebraic Curves and Riemann Surfaces*. Graduate Studies in Math. *5*. Amer. Math. Soc., Providence, 1995.

[NZ] NIVEN, I. and H. S. ZUCKERMAN. *An Introduction to the Theory of Numbers*, 2nd ed. Wiley, New York, 1966.

[Pr] PRETZEL, O. *Codes and Algebraic Curves*. Clarendon Press, Oxford, 1998.

[Se] SERRE, J.-P. *Algebraic Groups and Class Fields*. Springer GTM *117*. Springer-Verlag, Berlin, 1988.

(Reçu le 13 novembre 2003)

Ming-Chang Kang

Department of Mathematics
National Taiwan University
Taipei, Taiwan
e-mail: kang@math.ntu.edu.tw