

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 50 (2004)
Heft: 3-4: L'enseignement mathématique

Artikel: On integral solutions of quadratic inequalities
Autor: Bochnak, J. / Jackson, T.

Bibliographie
DOI: <https://doi.org/10.5169/seals-2654>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

must be equivalent to one of a finite number of forms $r_i h_i$. Now let q be the least common denominator of the coefficients of $r_i h_i$. If we could have $f'(x_1, x_2, \dots, x_{n-1}, 0)$ equivalent to $r_i h_i$ for an infinite number of possible values of α_2 , with $0 \leq \alpha_2 < 1$, there would be two allowable values, say β and γ , with $0 < |\beta - \gamma| < \frac{1}{2q}$. Then considering $f'(0, 1, 0, \dots, 0)$, we see that $f'(x_1, x_2, \dots, x_{n-1}, 0)$ represents $k - \beta^2$ for $\alpha_2 = \beta$ and represents $k - \gamma^2$ for $\alpha_2 = \gamma$. However

$$|(k - \beta^2) - (k - \gamma^2)| = |\beta^2 - \gamma^2| < \frac{1}{2q} |\beta + \gamma| < \frac{1}{q}$$

contradicting the fact that distinct values of $r_i h_i$ are never closer than $\frac{1}{q}$. Similar considerations of $f'(0, 0, 1, \dots, 0), \dots, f'(0, 0, \dots, 1, 0)$ show that there are only a finite number of allowable values of $\alpha_3, \dots, \alpha_{n-1}$ for each $r_i h_i$.

Finally, to show that the number of allowable values of α_n in (8) is finite, consider the indefinite $(n-1)$ -ary sections $f'(x_1, x_2, \dots, x_{n-2}, 0, x_n)$, $f'(x_1, x_2, \dots, x_{n-2}, x_n, x_n)$ and $f'(x_1, x_2, \dots, x_{n-2}, 2x_n, x_n)$. At least one of these, called $\psi(x_1, x_2, \dots, x_{n-2}, x_n)$ say, has a non-zero determinant (whose value depends only on k and the coefficients of g'). So, taking $\varepsilon_2 = \varepsilon |D(f)|^{1/n} |D(\psi)|^{-1/n-1} > 0$, ψ will represent a small positive value

$$v_2 < \varepsilon_2 |D(\psi)|^{1/n-1} = \varepsilon |D(f)|^{1/n}$$

unless it is equivalent to a multiple of one of a finite number of forms. Since $N(\psi) = 1$, there will again only be one allowable multiple for each of the finite number of forms. As before, we can also see that for each of the finite number of possibilities for $\alpha_2, \dots, \alpha_{n-1}, k, g'$ there will only be a finite number of allowable values of α_n .

It follows that the number of forms $f \in \mathcal{E}_n$ for which (6) fails (for a given $\varepsilon > 0$) is finite. So the theorem holds for n -ary forms.

REFERENCES

- [1] CASSELS, J. W. S. *An Introduction to Diophantine Approximation*. Camb. Univ. Press, Cambridge, 1957.
- [2] ——— *An Introduction to the Geometry of Numbers*. Springer, Berlin, 1971.
- [3] CASSELS, J. W. S. and H. P. F. SWINNERTON-DYER. On the product of three homogeneous linear forms and indefinite ternary quadratic forms. *Phil. Trans. Roy. Soc. London* 248 (1955), 73–96.
- [4] DANI, S. G. and G. MARGULIS. Values of quadratic forms at integral points: an elementary approach. *L'Enseign. Math.* (2) 36 (1990), 143–174.

- [5] DAVENPORT, H. On indefinite ternary quadratic forms. *Proc. London Math. Soc.* (2) 51 (1950), 145–160.
- [6] JACKSON, T.H. Small positive values of indefinite binary quadratic forms. *J. London Math. Soc.* 43 (1968), 730–738.
- [7] — Small positive values of indefinite quadratic forms. *J. London Math. Soc.* (2) 1 (1969), 643–659.
- [8] OPPENHEIM, A. One-sided inequalities for quadratic forms : II quaternary forms. *Proc. London Math. Soc.* (3) 3 (1953), 418–429.
- [9] — Values of quadratic forms (I), (II). *Quart. J. Math. Oxford Ser.* (2) 4 (1953), 54–66.
- [10] VENKOV, B. A. On the extremal problem of Markov for indefinite ternary quadratic forms. *Izv. Akad. Nauk SSSR* 9 (1945), 429–494.
- [11] VULAKH, L. Y. On minima of indefinite rational quadratic forms. *J. Number Theory* 21 (1985), 275–285.
- [12] WATSON, G. L. One-sided inequalities for integral quadratic forms. *Quart. J. Math. Oxford Ser.* (2) 9 (1958), 99–108.
- [13] — *Integral Quadratic Forms*. Camb. Univ. Press, Cambridge, 1960.
- [14] — Asymmetric inequalities for indefinite quadratic forms. *Proc. London Math. Soc.* (3) 18 (1968), 95–113.

(Reçu le 20 avril 2004)

J. Bochnak

Mathematics Department
Vrije Universiteit
De Boelelaan 1081a
NL-1081 HV Amsterdam
The Netherlands
e-mail: bochnak@cs.vu.nl

T. Jackson

Mathematics Department
University of York
Heslington
GB-York YO10 5DD
England
e-mail: thj1@york.ac.uk

Leere Seite

Blank page

Page vide