

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 52 (2006)
Heft: 3-4: L'enseignement mathématique

Artikel: Hyperbolic polygons and simple closed geodesics
Autor: Parlier, Hugo

Bibliographie
DOI: <https://doi.org/10.5169/seals-2237>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

In [10], A. Yamada proved that $r \geq \ln \frac{2+\sqrt{7}}{\sqrt{3}}$ with equality occurring when S is the thrice punctured sphere (which proves that Yamada's lower bound is sharp). Interestingly, $\ln \frac{2+\sqrt{7}}{\sqrt{3}} = \operatorname{arcsinh}(\frac{2}{\sqrt{3}})$ and $\frac{1}{2} \ln 3 = \operatorname{arccosh}(\frac{2}{\sqrt{3}})$. The link is stronger than this apparent coincidence.

First of all, a thrice punctured sphere can be constructed by pasting two ideal hyperbolic triangles along all three edges. The three punctures are the points at infinity. The value $\frac{1}{2} \ln 3$ was also obtained using this triangle as the maximal radius of an inscribed disk. Theorem 5.6, seen in the light of Theorem 6.1, reads as follows:

THEOREM 6.2. *Let S be a hyperbolic Riemann surface of signature (g, n) . There exists a point $x \in S$ such that $D(x, \frac{1}{2} \ln 3)$ is simply connected. The value $\frac{1}{2} \ln 3$ is sharp. \square*

Notice that in contrast to Theorem 5.6, the value $\frac{1}{2} \ln 3$ is sharp for the set of surfaces with boundary but not for any individual surface. Surfaces with boundary play an important role in a variety of subjects, including the study of Klein surfaces (orientable or non-orientable hyperbolic surfaces). In other words, a Klein surface is either a hyperbolic Riemann surface, or the quotient of a closed hyperbolic Riemann surface by an orientation reversing involution (whose fixed point set is a set of disjoint simple closed geodesics). In terms of Klein surfaces, Theorem 5.6 implies the following corollary, where again the adjective "sharp" means sharp for the set of Klein surfaces.

COROLLARY 6.3. *Let S be a hyperbolic Klein surface. There exists a point $x \in S$ such that $D(x, \frac{1}{2} \ln 3)$ is simply connected. The value $\frac{1}{2} \ln 3$ is sharp.*

ACKNOWLEDGEMENTS. I thank Peter Buser for careful rereading and suggestions. I thank all the people I have bored with my ideas, and in particular Aline Aigon-Dupuy, who also helped with figures, and Gérard Maze.

REFERENCES

- [1] BAVARD, C. Disques extrémaux et surfaces modulaires. *Ann. Fac. Sci. Toulouse Math.* (6) 5 (1996), 191–202.
- [2] BEARDON, A. F. *The Geometry of Discrete Groups*. Springer-Verlag, New York, 1995. (Corrected reprint of the 1983 original.)

- [3] BIRMAN, J. S. and C. SERIES. Geodesics with bounded intersection number on surfaces are sparsely distributed. *Topology* 24 (1985), 217–225.
- [4] BUSER, P. *Geometry and Spectra of Compact Riemann Surfaces*. Birkhäuser, Boston, 1992.
- [5] CANARY, R. D., D. B. A. EPSTEIN and P. GREEN. Notes on notes of Thurston. In: *Analytical and Geometric Aspects of Hyperbolic Space*, Symp. Coventry and Durham/Engl. 1984, 3–92. London Math. Soc. Lect. Note Ser. 111, 1987.
- [6] CASSON, A. J. and S. A. BLEILER. *Automorphisms of Surfaces After Nielsen and Thurston*. Cambridge University Press, Cambridge, 1988.
- [7] FENCHEL, W. and J. NIELSEN. *Discontinuous Groups of Isometries in the Hyperbolic Plane*. Walter de Gruyter, Berlin, 2003.
- [8] KERCKHOFF, S. P. The Nielsen realization problem. *Ann. of Math. (2)*, 117 (1983), 235–265.
- [9] MARDEN, A. Universal properties of Fuchsian groups in the Poincaré metric. In: *Discontinuous Groups and Riemann Surfaces* (Proc. Conf., Univ. Maryland, College Park, Md., 1973), 315–339. Princeton Univ. Press, Princeton, N.J., 1974.
- [10] YAMADA, A. On Marden's universal constant of Fuchsian groups, II. *J. Analyse Math.* 41 (1982), 234–248.

(Reçu le 14 juillet 2006)

Hugo Parlier

Section de Mathématiques
Université de Genève
Case postale 64
CH-1211 Genève 4
Switzerland
e-mail: hugo.parlier@math.unige.ch

Leere Seite
Blank page
Page vide