Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 53 (2007)

Heft: 3-4

Artikel: On a question of Erds and Graham

Autor: Bauer, Mark / Bennett, Michael

DOI: https://doi.org/10.5169/seals-109547

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 26.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

ON A QUESTION OF ERDŐS AND GRAHAM

by Mark BAUER*) and Michael A. BENNETT†)

ABSTRACT. In this note, we sharpen work of Ulas to provide what is, in some sense, the minimal counterexample to a "conjecture" of Erdős and Graham about square values of products of disjoint blocks of consecutive integers.

1. Introduction

A remarkable paper of Erdős and Selfridge [2] completed a long-standing project of Erdős, showing that the product of consecutive integers cannot be a perfect power (and, in particular, a square). Referencing this problem, in Erdős and Graham [1] we find the following quote:

In the same spirit one could ask when the product of two or more disjoint blocks of consecutive integers can be a power. For example, if A_1, \ldots, A_n are disjoint intervals each consisting of at least 4 integers then perhaps the product $\prod_{k=1} \prod_{a_k \in A_k} a_k$ is a nonzero square in only a finite number of cases.

In making such an assertion, one presumes that Erdős and Graham were guided by density arguments (which do indeed suggest finiteness for the integer points on the corresponding hypersurfaces). As noted in Ulas [5], however, these arguments can fail to hold in this situation; if the A_i 's are taken to be blocks of precisely 4 integers, and if the number of such blocks is large enough, then the products take on square values infinitely often. In fact, Ulas suggests that there are likely infinitely many such blocks, in every case where the number of blocks is suitably large relative to the interval lengths. What seems to occur is that we have integral curves lying on our

^{*)} The first author is supported in part by a grant from NSERC.

^{†)} The second author is supported in part by grants from NSERC and the Killam Foundation.

hypersurfaces; to predict or guarantee that such curves do in fact exist appears to be a hard problem.

In this short note, we will search for what might be termed minimal counterexamples to the proposal of Erdős and Graham (as one can observe from the quote above, it is probably unfair to characterize this as a conjecture). Along the way, we will sharpen and generalize the results of [5].

SOME RESULTS

Let us, fixing a positive integer j and the j-tuple (k_1, k_2, \ldots, k_j) of positive integers, consider the equation

(2.1)
$$\prod_{i=1}^{j} \prod_{l=0}^{k_i-1} (x_i + l) = y^2,$$

where the variables x_1, x_2, \dots, x_i are positive integers with the property that

$$(2.2) x_s < x_t implies that x_s + k_s \le x_t.$$

We may clearly suppose, without loss of generality, that j > 1 (else we may appeal to the aforementioned theorem of Erdős-Selfridge) and that

$$2 \leq k_1 \leq k_2 \leq \cdots \leq k_j$$

(if $k_1 = 1$, then (2.1) has, trivially, infinitely many solutions).

Our two results are as follows. The first is a generalization of Theorem 1 of [5]. This result deals with situations omitted from consideration by Erdős and Graham; we include it for completeness.

THEOREM 2.1. If either $k_1 = 2$ or $(k_1, k_2) = (3,3)$ then equation (2.1) has infinitely many solutions with (2.2).

We also prove

THEOREM 2.2. If $j \ge 3$ and $k_i = 4$ for $1 \le i \le j$ then equation (2.1) has infinitely many solutions with (2.2).

This latter result affirms a conjecture of Ulas (who deduced a like statement for j = 4 and $j \ge 6$). The families of examples we construct to show that (2.1) has infinitely many solutions with (2.2) in case j = 3 and $(k_1, k_2, k_3) = (4, 4, 4)$ are, we believe, minimal in j amongst counterexamples to the Erdős-Graham proposal.

3. Proof of Theorem 2.1

Let us begin by supposing that $k_1 = 2$. We choose the x_i 's such that the product

$$\prod_{i=2}^{j} \prod_{l=0}^{k_i - 1} (x_i + l)$$

is of the form $4m_1m_2^2$, where $m_1 > 1$ is squarefree. In particular, if j = 2, $k_2 \in \{2,3\}$, we take $x_2 = 3$, while, otherwise, we may choose the x_i $(i \ge 2)$ such that

$$\prod_{i=2}^{J} \prod_{l=0}^{k_i - 1} (x_i + l) = m!$$

where $m = \sum_{i=2}^{j} k_i$ (note that m! cannot be equal to a square, by Bertrand's Postulate). We thus find that (2.1) is satisfied precisely when there exists an integer y_1 for which

$$x_1(x_1+1) = m_1 y_1^2$$

or, equivalently,

$$(2x_1 + 1)^2 - m_1(2y_1)^2 = 1.$$

Since this equation has, for each squarefree m_1 , infinitely many solutions in positive integers x_1 and y_1 , we conclude that, if $j \ge 2$ and $k_1 = 2$, then (2.1) necessarily has infinitely many solutions.

Next, suppose that $(k_1, k_2) = (3,3)$. If j = 2, as noted by K. R. S. Sastry, we may choose $x_1 = n$, $x_2 = 2n$, where n and m are positive integers satisfying

$$(n+2)(2n+1) = m^2$$

(see Guy [3]). As is well-known, there are infinitely many such solutions. We may therefore suppose that j > 2 and choose the x_i 's such that the product

$$\prod_{i=3}^{j} \prod_{l=0}^{k_i - 1} (x_i + l)$$

is of the form $m_1m_2^2$, where $m_1 \neq 2$ is squarefree. To do this, we may take the x_i such that

$$\prod_{i=3}^{J} \prod_{l=0}^{k_i-1} (x_i + l) = m!$$

with $m = \sum_{i=3}^{j} k_i$,

$$x_2 = 2x_1 + 2$$
 and $x_1 = 3x_0$.

Having a solution to equation (2.1) is thus equivalent to finding positive integers x_1 and y_1 satisfying

$$x_0(2x_0+1) = m! y_1^2$$

and so

$$(4x_0 + 1)^2 - 8m! y_1^2 = 1$$
.

Since m!/2 is not a square (for m > 2), it follows that this equation has infinitely many solutions in integers x_0 and y_1 . This completes our proof.

4. Proof of Theorem 2.2

As noted previously, Ulas derived Theorem 2.2 for j = 4 or $j \ge 6$. He observed a number of solutions in case j = 3 and strongly conjectured that there are infinitely many if j = 5. Since there exists a solution to equation (2.1) with

$$j = 2$$
, $k_1 = k_2 = 4$, $x_1 = 33$, $x_2 = 1680$,

we may conclude as desired by showing that (2.1) has infinitely many solutions with (2.2), if j = 3. In fact, we will provide three infinite families of solutions in this case.

Let us begin by considering the Diophantine equation

$$(4.1) u^2 - 3v^2 = -2.$$

The positive integral values of u that satisfy this equation are given by the recurrence

$$u_1 = 1$$
, $u_2 = 5$, $u_{n+1} = 4u_n - u_{n-1}$ for $n \ge 2$.

There are thus infinitely many such solutions with $u \equiv 1 \mod 4$. If we additionally assume that $u \geq 265$ and set

$$(x_1, x_2, x_3) = \left(\frac{u-5}{4}, \frac{v-3}{2}, \frac{u+1}{2}\right),$$

then

$$x_1(x_3+2)x_2(x_2+3) = \frac{(u^2-25)(v^2-9)}{32} = \frac{(u^2-25)^2}{96}$$

and

$$(x_1+1)x_3(x_2+1)(x_2+2) = \frac{(u^2-1)(v^2-1)}{32} = \frac{(u^2-1)^2}{96}$$

whereby

$$\prod_{i=1}^{3} \prod_{l=0}^{3} (x_i + l) = \left(\frac{(u^2 - 1)(u + 3)(u + 7)(u^2 - 25)}{768} \right)^2.$$

The assumption that $u \ge 265$ guarantees that the x_i 's satisfy (2.2).

Next, we note that (4.1) also has infinitely many solutions with $u \equiv -1 \mod 4$. For such a solution with $u \geq 3691$, we take

$$(x_1,x_2,x_3)=\left(\frac{u-7}{4},\frac{v-3}{2},\frac{u-7}{2}\right).$$

A little work shows that now

$$\prod_{i=1}^{3} \prod_{l=0}^{3} (x_i + l) = \left(\frac{(u^2 - 1)(u - 3)(u - 7)(u^2 - 25)}{768} \right)^2.$$

Our third family is also given by a recurrence. We now consider solutions to the equation

$$(4.2) u^2 - 5v^2 = 4$$

in odd integers u and v (so that $u \equiv 3 \mod 4$). We then take

$$(x_1, x_2, x_3) = \left(\frac{v-3}{2}, \frac{u-7}{4}, \frac{u+1}{2}\right)$$

and find that

$$\prod_{i=1}^{3} \prod_{l=0}^{3} (x_i + l) = \left(\frac{(u^2 - 9)(u+1)(u+5)(u^2 - 49)}{1280} \right)^2.$$

Notice that solutions to equation (4.2) satisfy $(u, v) = (L_{6n\pm 2}, F_{6n\pm 2})$, where L_k and F_k denote the kth Lucas and Fibonacci numbers, respectively.

5. Concluding remarks

It is worth noting that the examples in [5] in case j = 4 (and $k_i = 4$ for $1 \le i \le 4$) grow polynomially (that is, the number of such examples with $\max\{x_i\} < X$ exceeds X^{θ} for some $\theta > 0$), while those constructed here, for j = 3 (and $k_i = 4$) with $\max\{x_i\} < X$, are bounded in number by $c \log X$ for some constant c. It may be that this represents the true state of affairs for solutions to (2.1) in these instances, but it would appear to be most difficult to prove. More generally, it is possible that the behaviour of solutions to (2.1) is

governed in some way by the size of $\sum_{i=1}^{j} 1/k_i$. There is no obvious heuristic that comes to mind to support this, however.

We suspect that the case j=3, $k_1=k_2=k_3=4$ is, in some sense, minimal for (2.1) to have infinitely many solutions with (2.2). Indeed, we would guess that if j=2 and $k_1\geq 4$ then (2.1) has at most finitely many solutions with (2.2). The hypothesis that $k_1\geq 4$ is certainly necessary here (even when we cannot apply Theorem 2.1) as it is easy to show that (2.1) has infinitely many solutions with j=2 and $(k_1,k_2)=(3,4)$ (as before, one can construct at least two families from recurrence sequences). An argument of P.G. Walsh (private communication) provides reasonable support (via the ABC conjecture) for the belief that the number of solutions to (2.1) with (2.2) if $j=2, k_1=k_2=4$ is finite.

REFERENCES

- [1] ERDŐS, P. and R. L. GRAHAM. Old and New Problems and Results in Combinatorial Number Theory. Monograph L'Enseign. Math. 28, Geneva, 1980.
- [2] ERDŐS, P. AND J. L. SELFRIDGE. The product of consecutive integers is never a power. *Illinois J. Math.* 19 (1975), 292–301.
- [3] GUY, R. K. Unsolved Problems in Number Theory. 3rd ed. Springer, 2004.
- [4] SKAŁBA, M. Products of disjoint blocks of consecutive integers which are powers. *Collog. Math.* 98 (2003), 1–3.
- [5] ULAS, M. On products of disjoint blocks of consecutive integers. L'Enseign. Math. (2) 51 (2005), 331–334.

(Reçu le 6 février 2007)

Mark Bauer

University of Calgary
Calgary, Alberta
Canada

e-mail: mbauer@math.ucalgary.ca

Michael A. Bennett

University of British Columbia Vancouver, B.C. Canada

e-mail: bennett@math.ubc.ca