

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 53 (2007)
Heft: 3-4

Artikel: On a question of Erdős and Graham
Autor: Bauer, Mark / Bennett, Michael

Bibliographie

DOI: <https://doi.org/10.5169/seals-109547>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Siehe Rechtliche Hinweise.

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. Voir Informations légales.

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. See Legal notice.

Download PDF: 17.05.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

governed in some way by the size of $\sum_{i=1}^j 1/k_i$. There is no obvious heuristic that comes to mind to support this, however.

We suspect that the case $j = 3$, $k_1 = k_2 = k_3 = 4$ is, in some sense, minimal for (2.1) to have infinitely many solutions with (2.2). Indeed, we would guess that if $j = 2$ and $k_1 \geq 4$ then (2.1) has at most finitely many solutions with (2.2). The hypothesis that $k_1 \geq 4$ is certainly necessary here (even when we cannot apply Theorem 2.1) as it is easy to show that (2.1) has infinitely many solutions with $j = 2$ and $(k_1, k_2) = (3, 4)$ (as before, one can construct at least two families from recurrence sequences). An argument of P.G. Walsh (private communication) provides reasonable support (via the ABC conjecture) for the belief that the number of solutions to (2.1) with (2.2) if $j = 2, k_1 = k_2 = 4$ is finite.

REFERENCES

- [1] ERDŐS, P. and R.L. GRAHAM. *Old and New Problems and Results in Combinatorial Number Theory*. Monograph *L'Enseign. Math.* 28, Geneva, 1980.
- [2] ERDŐS, P. AND J. L. SELFRIDGE. The product of consecutive integers is never a power. *Illinois J. Math.* 19 (1975), 292–301.
- [3] GUY, R.K. *Unsolved Problems in Number Theory*. 3rd ed. Springer, 2004.
- [4] SKAŁBA, M. Products of disjoint blocks of consecutive integers which are powers. *Colloq. Math.* 98 (2003), 1–3.
- [5] ULAS, M. On products of disjoint blocks of consecutive integers. *L'Enseign. Math.* (2) 51 (2005), 331–334.

(Reçu le 6 février 2007)

Mark Bauer

University of Calgary
Calgary, Alberta
Canada
e-mail : mbauer@math.ucalgary.ca

Michael A. Bennett

University of British Columbia
Vancouver, B.C.
Canada
e-mail : bennett@math.ubc.ca