

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: Automorphism groups of right-angled Artin groups
Autor: Charney, Ruth / Vogtmann, Karen
DOI: <https://doi.org/10.5169/seals-109892>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

23

AUTOMORPHISM GROUPS OF RIGHT-ANGLED ARTIN GROUPS

by Ruth CHARNEY and Karen VOGTMANN

We propose some problems concerning outer automorphism groups of right-angled Artin groups (“RAAG’s”). Since F_n and \mathbf{Z}^n are examples of RAAG’s, it is tempting to view outer automorphism groups of general right-angled Artin groups as interpolating between $Out(F_n)$ and $GL(n, \mathbf{Z})$, and to ask to what extent properties common to both $Out(F_n)$ and $GL(n, \mathbf{Z})$ are shared by all of these groups.

Recall that a RAAG A_Γ based on a simplicial graph Γ is generated by the vertices of Γ , and the only relations are that v commutes with w if v and w are joined by an edge of Γ . Servatius [4] and Laurence [3] gave generators for $Aut(A_\Gamma)$, but not much else is known about this group except in the cases when Γ has no edges (so A_Γ is free) and Γ is the complete graph (so A_Γ is free abelian). Laurence’s generators are of four types: (1) inner automorphisms, (2) symmetries of Γ and inversions of the generators v , (3) *partial conjugations*, which conjugate everything in some connected component of $\Gamma - st(v)$ by v , and (4) *transvections*, which multiply v by w (on the right or left) if $lk(v) \subseteq st(w)$. Thus every element of $Out(A_\Gamma)$ lifts to an automorphism of the free group on the vertices of Γ , but the natural map from $Out(A_\Gamma)$ to $GL(n, \mathbf{Z})$ is usually not surjective.

There is a CAT(0) cube complex associated to any RAAG, whose 1-skeleton is the Cayley graph of the group, and which has a k -dimensional cube whenever the 1-skeleton of the cube appears (kind of a “cube-flag” complex, see [2]). The RAAG acts freely on this; the quotient has a loop for each generator and a k -torus for each complete subgraph on k vertices in Γ . This cube complex is 2-dimensional if and only if the graph Γ has no triangles. In this case, we have constructed an “outer space” for $Out(A_\Gamma)$, which is a contractible space on which $Out(A_\Gamma)$ acts with finite stabilizers [1]. Points in this outer space are morally actions of A_Γ on 2-dimensional CAT(0) complexes,

though the actual description is in terms of products of trees. This outer space is finite-dimensional, and we obtain :

THEOREM 23.1. *If Γ is connected and triangle-free then $\text{Out}(A_\Gamma)$ has a torsionfree subgroup of finite index and it has finite virtual cohomological dimension.*

Although we obtain both upper and lower bounds on the virtual cohomological dimension, these bounds in general do not match, so we ask :

QUESTION 23.2. *What is the exact virtual cohomological dimension of the outer automorphism group of a 2-dimensional right-angled Artin group ?*

The no-triangles condition on Γ was very convenient, but of course we would like to know what happens for any Γ :

QUESTION 23.3. *Do the outer automorphism groups of all right-angled Artin groups have torsion-free subgroups of finite index ?*

QUESTION 23.4. *Calculate the virtual cohomological dimension of the automorphism group of any right-angled Artin group.*

ADDED IN PROOF. For progresses on these problems see : R. CHARNEY and K. VOGTMANN. ‘Finiteness properties of automorphism groups of right-angled Artin groups’, submitted. And also : BUX, K.-U., R. CHARNEY and K. VOGTMANN. ‘Automorphisms of two-dimensional RAAGS and partially symmetric automorphisms of free groups’, submitted.

REFERENCES

- [1] CHARNEY, R., J. CRISP and K. VOGTMANN. Automorphisms of 2-dimensional right-angled Artin groups. *Geom. Topol.* 11 (2007), 2227–2264.
- [2] CHARNEY, R. and M. DAVIS. Finite $K(\pi, 1)$ s for Artin groups. In: *Prospects in Topology (Princeton, NJ, 1994)*. Annals of Mathematics Studies 138, 110–124. Princeton University Press, Princeton, NJ, 1995.
- [3] LAURENCE, M.R. A generating set for the automorphism group of a graph group. *J. London Math. Soc. (2)* 52 (1995), 318–334.
- [4] SERVATIUS, H. Automorphisms of graph groups. *J. Algebra* 126 (1989), 34–60.

R. Charney

Brandeis University
304 Goldsmith
Waltham, MA 02454-9110
USA
e-mail: charney@brandeis.edu

K. Vogtmann

Cornell University
555 Malott Hall
Ithaca, NY 14853-4201
USA
e-mail: vogtmann@math.cornell.edu