Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 54 (2008)

Heft: 1-2

Artikel: The homotopy exponent conjecture of p-completed classifying spaces

of finite groups

Autor: Cohen, Fred / Levi, Ran

DOI: https://doi.org/10.5169/seals-109894

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 27.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

THE HOMOTOPY EXPONENT CONJECTURE FOR p-COMPLETED CLASSIFYING SPACES OF FINITE GROUPS

by Fred COHEN and Ran LEVI

Conjecture 25.1. Let π be a finite group, and let $B\pi_p^{\wedge}$ denote its p-completed classifying space. Then the homotopy groups of $B\pi_p^{\wedge}$ have a bounded exponent, i.e., there exists an integer r such that

$$p^r \cdot \pi_*(B\pi_p^\wedge) = \{0\} .$$

Since π is finite, the fundamental group of $B\pi_p^{\wedge}$ is the finite p-group given by the quotient of π by its minimal normal subgroup of p-power index, denoted by $O^p(\pi)$. If the order of $O^p(\pi)$ is not divisible by p, then $B\pi_p^{\wedge}$ is homotopy equivalent to $B(\pi/O^p(\pi))$, and the conjecture reduces to a triviality. If p divides the order of $O^p(\pi)$, then $B\pi_p^{\wedge}$ has infinitely many non-vanishing homotopy groups, all of which are finite p-groups. It is therefore natural to ask whether there is an upper bound on the exponent of these homotopy groups.

Our conjecture is related to the Moore finite exponent conjecture. Recall that a group G is said to have exponent m if every element $x \in G$ of finite order has order dividing m. The Moore conjecture states that for a simply-connected space X of the homotopy type of a finite CW-complex, the graded group $\pi_*(X) \otimes \mathbf{Q}$ is a finite dimensional rational vector space if and only if for every prime p, the graded group $\pi_*(X) \otimes \mathbf{Z}_{(p)}$ has an exponent.

In particular, if X is a finite simply connected CW-complex whose homotopy groups are all finite, then the Moore conjecture implies that $\pi_*(X)$ has an exponent. One can show that if π is a finite group, then the component of the constant loop in $\Omega B\pi_p^\wedge$ is a retract of the loop space of a finite simply connected torsion complex. Therefore our conjecture would follow at once if the much stronger Moore conjecture were true.

For any finite group π , $\pi_i(B\pi_p^{\wedge}) \cong \pi_i(BO^p(\pi)_p^{\wedge})$ for all $i \geq 2$. Furthermore, for each $i \geq 3$ $\pi_i(B\pi_p^{\wedge}) \cong \pi_i(BU^p(\pi)_p^{\wedge})$, where $U^p(\pi)$ is the p-universal central extension of $O^p(\pi)$. In all known examples for the conjecture, the order of the Sylow p-subgroup of $U^p(\pi)$ is an upper bound for the order of torsion in $\pi_*(B\pi_p^{\wedge})$. There are examples where this bound is sharp.

It is known that for any finite group π , the p-torsion in the homology of the loop space $\Omega B\pi_p^{\wedge}$ is bounded above by the order of the Sylow p-subgroup of $O^p(\pi)$.

EXAMPLE 25.2. Some examples are known. A few are given by

$$\pi = A_5, A_6, A_7, J_1, M_{11}$$

at p=2. A few more at the prime 2 are given by those finite simple groups of 2-rank 2 (including M_{11}) with the possible exception of $U(3, \mathbf{F}_4)$. The finite simple groups of classical Lie type over the field \mathbf{F}_q , where q is a power of a prime different from p, provide a large family of examples at the prime p.

Finite simple groups of Lie type at the defining characteristic. Almost no examples are known of the behavior of $\pi_i(BG(\mathbf{F}_{p^k})_p^{\wedge})$, where G is a finite simple groups of Lie type.

Finite groups with Abelian Sylow p-subgroups. For a finite group π with cyclic Sylow p-subgroup and p > 2, $B\pi_p^{\wedge}$ is known to have a homotopy exponent. Furthermore, the best possible upper bound for this exponent is given by the order of the Sylow p-subgroup. On the other hand, for a finite group π with an abelian Sylow p-subgroup of rank larger than 1 neither one of the above statements is known to hold.

Alternating groups. Further natural open cases are A_n with n > 7 at the prime 2, and at all primes p in the cases where the Sylow p-subgroup is not cyclic.

The examples mentioned above are obtained by considering the structure of the loop space of $B\pi_p^{\wedge}$. This space sometimes admits a non-trivial product decomposition or is finitely resolvable by fibrations involving more recognizable spaces, which are known to have homotopy exponents by the work of Cohen–Moore–Neisendorfer [2] and [3].

The largest and most comprehensive reference on this subject is [4], where some other theoretical results are established, and many examples are given.

This paper is motivated by the spherical resolvability conjecture for ΩBG_p^{\wedge} due to the first author. This conjecture, if it was true would imply the conjecture proposed here. Unfortunately, as demonstrated in [5], the spherical resolvability conjecture is false, but at the same time some of the examples that prove it wrong do satisfy the finite exponent conjecture ([1], Cor. 5.4). A survey of many of the known results on spaces of type $B\pi_p^{\wedge}$ for π finite, including examples of homotopy exponents, is the paper [1].

REFERENCES

- [1] COHEN, F.R. and R. LEVI. On the homotopy theory of *p*-completed classifying spaces. In: *Group Representations: Cohomology, Group Actions and Topology, (Seattle, WA, 1996)*, 157–182. Proc. Sympos. Pure Math. 63. Amer. Math. Soc., Providence, RI, 1998.
- [2] COHEN, F.R., J.C. MOORE and J.A. NEISENDORFER. Torsion in homotopy groups. *Ann. of Math.* (2) 109 (1979), 121–168.
- [3] COHEN, F.R., J.C. MOORE and J.A. NEISENDORFER. Decompositions of loop spaces and applications to exponents. In: *Algebraic Topology, Aarhus 1978 (Proc. Sympos., Univ. Aarhus, Aarhus, 1978)*, 1–12. Lecture Notes in Mathematics 763. Springer-Verlag, Berlin, 1979.
- [4] LEVI, R. On Finite Groups and Homotopy Theory. Mem. Amer. Math. Soc. 118, no. 567 (1995).
- [5] LEVI, R. On homological rate of growth and the homotopy type of ΩBG_p^{\wedge} . *Math. Z.* 226 (1997), 429–444.

Fred Cohen

Mathematics Department University of Rochester Rochester, N.Y. 14620 USA

e-mail: cohf@math.rochester.edu

Ran Levi

Department of Mathematical Sciences University of Aberdeen Aberdeen AB24 3UE United Kingdom *e-mail*: ran@maths.abdn.ac.uk