

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 54 (2008)
Heft: 1-2

Artikel: The Friedlander-Milnor conjecture
Autor: Friedlander, Eric M.
DOI: <https://doi.org/10.5169/seals-109901>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

32

THE FRIEDLANDER–MILNOR CONJECTURE

by Eric M. FRIEDLANDER

The conjecture of the title of this note has resisted 40 years of effort and remains not only unsolved but also lacking in a plausible means of either proof or counter-example.

The original form of this conjecture is one I struggled with during my days at Princeton in the early 1970's:

CONJECTURE 32.1. *Let $G(\mathbf{C})$ be a complex reductive algebraic group and let $G(\mathbf{C})^\delta$ denote this group viewed as a discrete group. Then the map on classifying spaces of the continuous (identity) group homomorphism*

$$i: G(\mathbf{C})^\delta \rightarrow G(\mathbf{C})$$

induces an isomorphism in cohomology with finite coefficients \mathbf{Z}/n for any $n \geq 0$:

$$i^*: H^*(BG(\mathbf{C}), \mathbf{Z}/n) \xrightarrow{\simeq} H^*(G(\mathbf{C})^\delta, \mathbf{Z}/n).$$

Conjecture 32.1 is easily seen to be true for a torus (i.e., $G = \mathbf{G}_m^{\times r}$ for some $r > 0$), but even the simplest non-trivial case (that of $G = \mathrm{SL}_2$) remains inaccessible.

Guido and I published 5 papers together, all in some sense connected with this conjecture. We used the integral form $G_{\mathbf{Z}}$ of a complex reductive algebraic group (which is a group scheme over $\mathrm{Spec} \mathbf{Z}$) in order to form the group $G(F)$ of points of G with values in a field F . Most of our joint work investigated various relations between $G(\mathbf{C})$ and $G(F)$, the case $F = \overline{\mathbf{F}}_p$ (the algebraic closure of a prime field \mathbf{F}_p) being of special interest.

One knows from considerations of étale cohomology that the cohomology of $BG(\mathbf{C})$ with \mathbf{Z}/n coefficients is naturally isomorphic to that of the étale

homotopy classifying space of the algebraic group G_F for F algebraically closed of characteristic $p \geq 0$:

$$H^*(BG(\mathbf{C}), \mathbf{Z}/n) \simeq H^*((BG_F)_{\text{et}}, \mathbf{Z}/n), \quad \text{provided that } (p, n) = 1.$$

This enables one to construct a map $H^*(BG(\mathbf{C}), \mathbf{Z}/n) \rightarrow H^*(G(F), \mathbf{Z}/n)$ relating the cohomology with mod- n coefficients of the classifying space of $G(\mathbf{C})$ with the cohomology with mod- n coefficients of the discrete group $G(F)$ for any field F .

The following is a generalization of Conjecture 32.1, one that appears likely to be true if and only if Conjecture 32.1 is valid.

CONJECTURE 32.2. *Let $G(\mathbf{C})$ be a complex reductive algebraic group, let $n > 0$ be a positive integer, and let p denote either 0 or a prime which does not divide n . Then for any algebraically closed field F of characteristic p , the comparison of the cohomology of $BG(\mathbf{C})$ and $G(F)$ determines an isomorphism*

$$H^*(G(F), \mathbf{Z}/n) \simeq H^*(BG(\mathbf{C}), \mathbf{Z}/n).$$

In our first paper together [1], Guido and I began our investigation of “locally finite approximations” of Lie groups. We also formulated the following conjecture and proved it equivalent to Conjecture 32.2.

CONJECTURE 32.3. *Let F be an algebraically closed field of characteristic $p \geq 0$ and let $n > 0$ be a positive integer not divisible by p if $p > 0$. Then Conjecture 32.2 is valid for $G(F)$ if and only for every $0 \neq x \in H^*(G(F), \mathbf{Z}/n)$, there exists some finite subgroup $\pi \subset G(F)$ such that x restricts non-trivially to $H^*(\pi, \mathbf{Z}/n)$.*

The most familiar form of the “Friedlander–Milnor Conjecture” is that formulated by John Milnor in [2]. In that paper, Milnor verifies this conjecture for solvable groups.

CONJECTURE 32.4. *Let G be a Lie group with finitely many components and let G^δ denote the same group now viewed as a discrete group. Then for any integer $n > 0$, the continuous (identity) map $i: G^\delta \rightarrow G$ induces an isomorphism on cohomology with mod- n coefficients:*

$$i^*: H^*(BG, \mathbf{Z}/n) \xrightarrow{i^*} H^*(G^\delta, \mathbf{Z}/n).$$

We remark that the most substantial progress to date on these conjectures is due to Andrei Suslin, who proves a “stable” version of Conjectures 32.1 and 32.2 in [3].

REFERENCES

- [1] FRIEDLANDER, E. and G. MISLIN. Cohomology of classifying spaces of complex Lie groups and related discrete groups. *Comment. Math. Helv.* 59 (1984), 347–361.
- [2] MILNOR, J. On the homology of Lie groups made discrete. *Comment. Math. Helv.* 58 (1983), 72–85.
- [3] SUSLIN, A. On the K -theory of local fields. *J. Pure Appl. Algebra* 34 (1984), 301–318.

E. Friedlander

Department of Mathematics
Northwestern University
Evanston, IL 60208
USA
e-mail: eric@math.northwestern.edu